



## SIR PRATEEK JAIN

- . Founder @ Physicsaholics
- . Top Physics Faculty on Unacademy (IIT JEE & NEET)
- . 8+ years of teaching experience in top institutes like FIITJEE (Delhi, Indore) , CP (KOTA) etc.
- . Produced multiple Top ranks.
- . Research work with HC Verma sir at IIT Kanpur
- . Interviewed by International media.



## NEET UG subscription

PLUS

ICONIC \*\*

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months ₹2,100/mo  
No cost EMI +10% OFF ₹50,400 >

18 months ₹2,363/mo  
No cost EMI +10% OFF ₹42,525 >

12 months ₹2,888/mo  
No cost EMI +10% OFF ₹34,650 >

6 months ₹4,200/mo  
No cost EMI +10% OFF ₹25,200 >

To be paid as a one-time payment

[View all plans](#)



Awesome! **PHYSICSLIVE** code applied



# PHYSICSLIVE

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS and learn from India's Top Faculties.



## NEET UG subscription

PLUS

ICONIC \*\*

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months ₹2,100/mo  
No cost EMI +10% OFF ₹50,400 >

18 months ₹2,363/mo  
No cost EMI +10% OFF ₹42,525 >

12 months ₹2,888/mo  
No cost EMI +10% OFF ₹34,650 >

6 months ₹4,200/mo  
No cost EMI +10% OFF ₹25,200 >

To be paid as a one-time payment

[View all plans](#)



Awesome! **PHYSICSLIVE** code applied



For Video Solution of this DPP, Click on below link

Video Solution  
on Website:-

<https://physicsaholics.com/home/courseDetails/53>

Video Solution  
on YouTube:-

[https://youtu.be/5hJdZ\\_raBLQ](https://youtu.be/5hJdZ_raBLQ)

# **JEE Main & Advanced, NSEP, INPhO, IPhO** **Physics DPP**

**DPP-5 NLM: Spring Force**  
**By Physicsaholics Team**


Q) A bead of mass  $m$  is attached to one end of a spring of natural length  $R$  and spring constant  $k = \frac{(\sqrt{3}+1)mg}{R}$ . The other end of the spring is fixed at point A on a smooth vertical ring of radius  $R$  as shown in figure. The normal reaction at B just after it is released to move is:

(a)  $\frac{mg}{2}$

(b)  $\sqrt{3}mg$

(c)  $3\sqrt{3}mg$


(d)  $\frac{3\sqrt{3}mg}{2}$



Ans. d



Solution:




length of spring  
 $= 2R \cos 30^\circ = R\sqrt{3}$

elongation in spring  $= R\sqrt{3} - R = R(\sqrt{3} - 1)$


spring force  $Kx = \frac{(\sqrt{3} + 1)mg}{R} \times R(\sqrt{3} - 1)$   
 $= 2mg$

$N = mg \cos 30^\circ + Kx \cos 30^\circ = \frac{mg\sqrt{3}}{2} + 2mg \frac{\sqrt{3}}{2}$   
 $= \frac{3mg\sqrt{3}}{2}$   
 (D)



Q) In the figure, the ball A is released from rest when the spring is at its natural (unstretched) length. For the block B, of mass  $M$  to leave contact with the ground at some stage, the minimum mass of A must be

- (a)  $2M$
- (b)  $M$
- (c)  $M/2$
- (d) a function of  $M$  and the force constant of the spring



Join Unacademy PLUS Referral Code :

Physicslive



Ans. c

Solution:

maximum elongation  
in spring

$\hat{=}$  maximum Displacement  
of A


$$= 2mg/k$$

maximum spring force

$$= k \times \frac{2mg}{k} = 2mg$$

If this force is greater  
than or equal to  $Mg$ ,  
it will lift.


$$2mg \geq Mg \Rightarrow m \geq M/2$$



(c)

Q) In the system shown  $m_1 > m_2$ . System is held at rest by thread BC. Just after lower thread is burnt, C

- (a) Acceleration of  $m_2$  is upwards
- (b) Magnitude of acceleration of both blocks will be  $\left( \frac{m_1 - m_2}{m_1 + m_2} \right) g$
- (c) Acceleration of  $m_1$  will be equal to zero
- (d) Magnitudes of acceleration of two blocks will be non-zero & unequal



Join Unacademy PLUS Referral Code :

Physicslive

Ans. a, c

Solution:

Since spring is directly connected to string.


$$T = F_s = m_1 g$$

After burning string C,  $T_1 = 0$  but all other forces will remain same.

$\Rightarrow$  acceleration of  $m_1 = 0$

acceleration of  $m_2$

$$= \frac{m_1 g - m_2 g}{m_2} \uparrow$$



(A, c)




Q) From the fixed pulley, masses 2 kg, 1 kg and 3 kg are suspended as shown in figure. Find the extension in the spring when acceleration of 3kg and 1kg is same if spring constant of the spring  $k = 100 \text{ N/m}$ . ( $g = 10 \text{ m/s}^2$ )

(a) 10 cm

(b) 20 cm

(c) 30 cm

(d) 25 cm



Join Unacademy PLUS Referral Code :

Physicslive

Ans. **b**

Solution:

$$a = \frac{\text{supporting} - \text{opposing}}{\text{total mass}}$$

$$a = \frac{4g - 2g}{6} = g/3$$


from F.B.D of 3kg

$$3g - F_s = 3a$$

$$3g - 100x = g$$

$$x = \frac{2g}{100} = \frac{1}{5} \text{ m}$$


$$x = 20 \text{ cm}$$



(3)

Q) The acceleration of masses  $m_1$  and  $m_2$  and  $m_3$  shown in figure just after string is cut at point A is given by  $a_1$ ,  $a_2$  and  $a_3$  choose the correct answer

- (a)  $a_1 = g, a_2 = g/2, a_3 = 0$
- (b)  $a_1 = \left(1 + \frac{m_2}{m_1}\right)g, a_2 = 0, a_3 = g$
- (c)  $a_1 = g, a_2 = \left(1 + \frac{m_1}{m_2}\right)g, a_3 = 0$
- (d)  $a_1 = 0, a_2 = \left(1 + \frac{m_1}{m_2}\right)g, a_3 = g$



Join Unacademy PLUS Referral Code :

Physicslive

Ans. d




Solution:

Just after cutting  
A,  $T=0$  &  $T_1=0$

$m_3$  will fall freely.


$m_1$  will be in  
equilibrium

F.B.D of  $m_2$



A free body diagram of mass  $m_2$  showing two downward arrows: one labeled  $m_2g$  and the other labeled  $F_s$ .

$$a_2 = \frac{m_2g + m_1g}{m_2}$$



Q) Two identical bars of mass  $m$  each are connected by a weightless spring of stiffness  $x$  and length (in the non-deformed state)  $l_0$  rest on a horizontal plane. A constant horizontal force  $F$  starts acting on one of the bar. Find the maximum elongation in spring during the subsequent motion of the system?

(a)  $F/k$

(b)  $F/2k$

(c)  $2F/k$


(d)  $3F/2k$

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. a

Solution:



elongation  $x$

$$a_1 = \frac{F - kx}{m}, \quad a_2 = \frac{kx}{m}$$

relative acceleration  $a = a_1 - a_2 = \frac{F - 2kx}{m}$

$\Rightarrow V \frac{dv}{dx} = \frac{F - 2kx}{m}$  where  $V$  is relative velocity

at maximum elongation  $V = 0$

$$\Rightarrow \int_0^0 V dv = \int_0^x \frac{F - 2kx}{m} dx$$

$$\Rightarrow Fx - kx^2 = 0 \Rightarrow x = \frac{F}{k}$$


Q) A wedge of mass 'M' and angle of inclination ' $\theta$ ' and of mass 'm' is arranged in a manner shown in the figure. The spring of force constant 'k' attached to the wedge. Assuming the pulleys to be massless and all surfaces to be frictionless. Find the compression of the spring under equilibrium condition.

(a)  $\frac{mg \sin \theta}{k}$

(b)  $\frac{2mg \sin \theta}{k}$

(c)  $\frac{mg \sin \theta}{2k}$

(d) none of these





Ans. a

Solution:

$$T \cos \theta + kx = N \sin \theta + T$$



$$\Rightarrow N \sin \theta + T(1 - \cos \theta) = kx \quad \text{--- (i)}$$

$$T = mg \sin \theta \quad \text{--- (ii)}$$

$$N = mg \cos \theta \quad \text{--- (iii)}$$

$$\Rightarrow mg \sin \theta \cos \theta + mg \sin \theta (1 - \cos \theta) = kx$$

$$\Rightarrow x = \frac{mg \sin \theta}{k}$$




Q) Find acceleration of mass  $2m$  just after burning string A ? Initially system was in equilibrium .

(a)  $g$

(b)  $g/2$

(c)  $g/3$

(d)  $g/4$



Join Unacademy PLUS Referral Code :

Physicslive

Ans. a


## Solution

Just after burning  
String A,  $T' = 0$

$T$  will change  
& block  $2m$   
& upper block  
 $m$  move together

acceleration of  
 $2m$

$$= \frac{2mg + F}{3m}$$
$$= g \downarrow$$



(A)




Q) Find relative acceleration of block  $m$  and  $2m$  just after burning string ? Initially system was in equilibrium and tension in string was  $2mg$ .

(a)  $g$

(b)  $2g$

(c)  $3g$

(d)  $4g$




Join Unacademy PLUS Referral Code :

Physicslive

Ans. c

Solution:



acceleration of m

$$= \frac{2mg}{m} = 2g \downarrow$$

acceleration of 2m

$$= \frac{2mg}{2m} = g \uparrow$$

relative acceleration

$$= 3g$$

(c)

Q) Two springs of stiffness  $k$  and  $2k$  are connected in series. Free end of first spring (stiffness  $k$ ) is fixed and free end of second spring is pulled by an external agent with constant velocity  $v$ . Find velocity of joint of springs ?

(a)  $v/3$

(b)  $2v/3$

(c)  $v$


(d) none of these

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. b

Solution:



Let elongation in  $2K$  is  $x$  & that of  $K$  is  $x_1$ .

Since net force on joint is zero.

$$Kx_1 = 2Kx \Rightarrow x_1 = 2x$$

$$V = \frac{d}{dt}(3x) = 3 \frac{dx}{dt}$$

$$V_1 = \frac{d}{dt}(2x) = 2 \frac{dx}{dt} = \frac{2V}{3} \quad (b)$$



Q) Two ends of a spring of natural length  $l$  and stiffness  $k$  are being pulled apart by external agents with constant velocities  $v$  and  $2v$ . Find velocity of mid point of spring ?

(a)  $v$

(b)  $v/2$

(c)  $3v/2$



(d) zero

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. b

Solution:



\* all parts of spring elongate uniformly.

$$V_1 + V = \frac{dx}{dt}$$

$$V_1 + V = 3V/2$$

$$V_1 = V/2$$

$$3V = \frac{d}{dt}(2x) = 2 \frac{dx}{dt}$$

$$\Rightarrow \frac{dx}{dt} = \frac{3V}{2}$$

For Video Solution of this DPP, Click on below link

Video Solution  
on Website:-

<https://physicsaholics.com/home/courseDetails/53>

Video Solution  
on YouTube:-

[https://youtu.be/5hJdZ\\_raBLQ](https://youtu.be/5hJdZ_raBLQ)

Chalo Niklo