

# Notes on Abbot's Introduction to Analysis

Zaid Khan

July 24, 2015

# Contents

<b>1</b>	<b>TODO</b>	<b>3</b>
<b>2</b>	<b>Sequences and Series</b>	<b>3</b>
2.1	The Limit of Sequence . . . . .	3
2.1.1	Definition of convergence for a sequence . . . . .	3
2.1.2	Outline of a Proof of Convergence . . . . .	3
2.2	The Monotone Convergence Theorem . . . . .	3
2.2.1	Monotone Convergence Theorem . . . . .	3
2.3	Subsequences and the Bolzano-Weierstrass Theorem . . . . .	3
2.3.1	Subsequence Convergence Theorem . . . . .	3
2.3.2	Bolzano-Weierstrass Theorem . . . . .	3
2.4	The Cauchy Criterion . . . . .	4
2.4.1	Cauchy Criterion Definition . . . . .	4
2.4.2	Cauchy Convergence . . . . .	4
2.4.3	Effects of Completeness on Convergence . . . . .	4
2.5	Properties of Infinite Series . . . . .	4
2.5.1	Convergence of an Infinite Series . . . . .	4
2.5.2	Cauchy Criterion for Series . . . . .	4
2.5.3	Comparison Test . . . . .	4
2.5.4	Absolute Convergence Test . . . . .	5
2.5.5	Alternating Series Test . . . . .	5
2.5.6	Absolute Convergent Rearrangement Theorem For Series . . . . .	5
<b>3</b>	<b>Topology of the Reals</b>	<b>5</b>
3.1	Open and Closed Sets . . . . .	5
3.1.1	Epsilon Neighborhoods . . . . .	5
3.1.2	Definition of an Open Set . . . . .	5
3.1.3	Union and Intersection of Open Sets . . . . .	5
3.1.4	Limit Points . . . . .	5
3.1.5	Limit Point Sequence Convergence . . . . .	5
3.1.6	Definition of an Isolated Point . . . . .	6
3.1.7	Definition of a Closed Set . . . . .	6
3.1.8	Closed Sets and Cauchy Sequences . . . . .	6
3.1.9	Density of $\mathbb{Q}$ in $\mathbb{R}$ . . . . .	6
3.1.10	Closure . . . . .	6
3.1.11	Complements and Open/Closed Sets . . . . .	6
3.2	Compact Sets . . . . .	6
3.2.1	Compactness . . . . .	6
3.2.2	Characterization of Compactness in $\mathbb{R}$ . . . . .	6
3.2.3	Nested Compact Set Property . . . . .	6
3.2.4	Open Covers . . . . .	7
3.2.5	Heine-Borel Theorem . . . . .	7

## 1 TODO

## 2 Sequences and Series

### 2.1 The Limit of Sequence

#### 2.1.1 Definition of convergence for a sequence

A sequence converges to some number  $a$  if  $\forall \epsilon > 0 \quad \exists N \in \mathbb{N}$  such that  $\forall n \geq N \implies |a_n - a| < \epsilon$ .

#### 2.1.2 Outline of a Proof of Convergence

- Let  $\epsilon > 0$  be arbitrary.
- Demonstrate a choice of  $N \in \mathbb{N}$ . This essentially means solve for  $N$  in terms of  $\epsilon > 0$ .
- Show that  $N$  actually works. This is simply the inequality that you derived in the previous item.
- Assume  $n \geq N$ .
- Derive the inequality  $|a_n - a| < \epsilon$ . Again, this is simply equivalent to solving for  $\epsilon$  in an inequality.

### 2.2 The Monotone Convergence Theorem

A sequence is **monotone** if it is either increasing or decreasing.

#### 2.2.1 Monotone Convergence Theorem

If a sequence is monotone and bounded, then it converges.

An infinite series is said to be converging if the sequence of partial sums is converging. Note how the definition of convergence of a series is linked back to the notion of convergence for a sequence.

### 2.3 Subsequences and the Bolzano-Weierstrass Theorem

#### 2.3.1 Subsequence Convergence Theorem

The subsequences of a convergent sequence converge to the same limit as the original sequence.

#### 2.3.2 Bolzano-Weierstrass Theorem

There exists a convergent subsequence within every bounded sequence.

## 2.4 The Cauchy Criterion

### 2.4.1 Cauchy Criterion Definition

A sequence  $a_n$  is called a *Cauchy sequence* if  $\forall \epsilon > 0 \quad \exists N \in \mathbb{N} : m, n \geq N \implies |a_n - a_m| < \epsilon$ . To simplify, a sequence is cauchy if there exists a point in the sequence after which all the numbers are closer to each other than to any arbitrary epsilon.

### 2.4.2 Cauchy Convergence

- Every convergent sequence is a Cauchy sequence.
- Cauchy sequences are bounded.
- A sequence converges if and only if it is a Cauchy Sequence.

### 2.4.3 Effects of Completeness on Convergence

Completeness is a necessary condition in order for the above definitions of convergence to work correctly. In particular, we need a way to guarantee that when a sequence is converging, the number that it is converging to is actually there. This is part of the reason why the real numbers were introduced - the rationals have holes in them. The Axiom of Completeness, Nested Interval Property, Bolzano-Weierstrauss, Cauchy Criterion, and Monotone Convergence Theorem are all equivalent.

## 2.5 Properties of Infinite Series

### 2.5.1 Convergence of an Infinite Series

The convergence of an infinite series  $a_k$  is defined in terms of the sequence of its partial sums  $s_n$ .  $\sum_{k=1}^{\infty} a_k = A$  means that  $\lim(s_n) = A$ .

### 2.5.2 Cauchy Criterion for Series

- The series  $\sum_{k=1}^{\infty} a_k$  converges if and only if, given  $\epsilon > 0 \quad \exists N \in \mathbb{N} : n > m \geq N \implies |a_{m+1} + a_{m+2} + \dots + a_n| < \epsilon$ .
- **Corollary:** If the series  $\sum_{k=1}^{\infty} a_k$  converges, then  $a_k \rightarrow 0$ .

### 2.5.3 Comparison Test

Assume  $(a_k)$  and  $(b_k)$  are sequences satisfying  $0 \leq a_k \leq b_k \quad \forall k \in \mathbb{N}$ .

- If  $\sum b_k$  converges, then  $\sum a_k$  also converges.
- If  $\sum a_k$  diverges, then  $\sum b_k$  also diverges.

### 2.5.4 Absolute Convergence Test

If the series  $\sum |a_k|$  converges, then  $\sum a_k$  converges as well. A sequence is said to converge absolutely if for a sequence  $a_k$ ,  $\sum_{k=1}^{\infty} |a_k|$  also converges. Otherwise, the sequence is said to converge conditionally.

### 2.5.5 Alternating Series Test

Let  $(a_n)$  be a sequence that is monotonically decreasing and has a limit of 0. Then, the alternating series  $\sum_{n=1}^{\infty} (-1)^{n+1} a_n$  converges.

### 2.5.6 Absolute Convergent Rearrangement Theorem For Series

If a series converges absolutely, then any rearrangement of the series converges to the same limit.

## 3 Topology of the Reals

### 3.1 Open and Closed Sets

#### 3.1.1 Epsilon Neighborhoods

An  $\epsilon$ -neighborhood is a set  $V_{\epsilon}(x) = \{x \in \mathbb{R} : |x - a| < \epsilon\}$ . Note that the interval is open - it does not contain the endpoints since the distance must be less than epsilon.

#### 3.1.2 Definition of an Open Set

A set  $O \subseteq \mathbb{R}$  is *open* if  $\forall a \in O$  there exists an  $\epsilon$ -neighborhood  $V_{\epsilon}(a) \subseteq O$ . Recall that an  $\epsilon$ -neighborhood is the set of points surrounding a point that are less than  $\epsilon$  away from it. This set particular set does not contain the endpoints.

#### 3.1.3 Union and Intersection of Open Sets

The union of an arbitrary collection of open sets is open. The intersection of a finite collection of open sets is open.

#### 3.1.4 Limit Points

A point  $x$  is a *limit point* of a set  $A$  if every  $\epsilon$ -neighborhood  $V_{\epsilon}(x)$  of  $x$  intersects  $A$  at some point other than  $x$ . Alternate interpretation: a limit point is the limit of a sequence at the edge of the set.

#### 3.1.5 Limit Point Sequence Convergence

A point  $x$  is a limit point of a set  $A$  if and only if  $x = \lim a_n$  for some sequence  $(a_n)$  contained in  $A$  satisfying  $a_n \neq x \quad \forall n \in \mathbb{N}$ .

### 3.1.6 Definition of an Isolated Point

A point  $a \in A$  is an isolated point of  $A$  if it is not a limit point of  $A$ .

### 3.1.7 Definition of a Closed Set

A set  $F \subseteq \mathbb{R}$  is closed if it contains its limit points.

### 3.1.8 Closed Sets and Cauchy Sequences

A set  $F \subseteq \mathbb{R}$  is closed if and only if every Cauchy sequence contained in  $F$  has a limit that is also an element of  $F$ .

### 3.1.9 Density of $\mathbb{Q}$ in $\mathbb{R}$

For every  $y \in \mathbb{R}$ , there exists a sequence of rational numbers that converges to  $y$ .

### 3.1.10 Closure

Given a set  $A \subseteq \mathbb{R}$ , let  $L$  be the set of all limit points of  $A$ . The set  $A \cup L$  is defined to be the closure of  $A$ .

For any  $A \subseteq \mathbb{R}$ , the closure  $\overline{A}$  is a closed set and the smallest closed set containing  $A$ .

### 3.1.11 Complements and Open/Closed Sets

A set  $O$  is open if and only if  $O^c$  is closed. Likewise, a set  $F$  is closed if and only if  $F^c$  is open.

- The union of a finite collection of finite sets is closed.
- The intersection of an arbitrary collection of arbitrary sets is closed.

## 3.2 Compact Sets

### 3.2.1 Compactness

A set  $K \subseteq \mathbb{R}$  is compact if every sequence in  $K$  has a subsequence that converges to a limit which is also in  $K$ .

### 3.2.2 Characterization of Compactness in $\mathbb{R}$

A set  $K \subseteq \mathbb{R}$  is compact if and only if it is closed and bounded. A set  $A$  is bounded if  $\exists M : |a| \leq M \forall a \in A$ .

### 3.2.3 Nested Compact Set Property

If  $K_1 \supseteq K_2 \supseteq K_3 \supseteq \dots K_n$  is a nested sequence of nonempty compact sets, then the intersection  $\bigcap_{n=1}^{\infty} K_n$  is not empty.

### 3.2.4 Open Covers

Let  $A \subseteq \mathbb{R}$ . An open cover for  $A$  is a possibly infinite collection of open sets whose union contains  $A$ . A finite subcover is finite collection of open sets from the original open cover whose union still completely contains  $A$ .

### 3.2.5 Heine-Borel Theorem

Let  $K$  be a subset of  $\mathbb{R}$ . All of the following statements are equivalent in the sense that any one of them implies the two others.

- $K$  is compact.
- $K$  is closed and bounded.
- Every open cover for  $K$  has a finite subcover.