



2026 BASIC ABSTRACT ALGEBRA STUDY

# LECTURE 1 HOMEWORKS

## SOLUTION - Q11

**Version**

v0.0.0

**Instructor**

Abiria

**Published on**

2026-05-18

**Hosted by**

Coding Lab

**Read this carefully before moving on.**

Please, clearly specify whether the statement in question is *true* or *false*, for the *prove* or *disprove* type of questions. As always, it is highly encouraged for you to write each step clearly with full sentences.

As a demonstration of a proof answer, consider the following example.

The above statement is **true**.

*Proof.* ...



As a demonstration of a disproof answer, consider the following example.

The above statement is **false**.

*Proof.* ...

**Note 1 (Notations)**

The following notations will be used throughout the following questions.

- $\mathbb{Z}$ : A set of all integers.
- $\mathbb{Z}^+$ : A set of all positive integers. i.e.,  $\mathbb{Z}^+ = \{1, 2, 3, \dots\}$ .
- $\mathbb{Z}_n$ : A set of non-negative integers from 0 to  $n - 1$  for some positive integer  $n$ . For instance,  $\mathbb{Z}_4 = \{0, 1, 2, 3\}$ .
- $n\mathbb{Z}$ : A set given as  $\{nx \mid x \in \mathbb{Z}\}$  for an integer  $n$ . i.e., the set of all multiples of  $n$  in  $\mathbb{Z}$ . For instance,  $5\mathbb{Z} = \{\dots, -10, -5, 0, 5, 10, \dots\}$ .
- $\mathbb{Q}$ : A set of all rational numbers.
- $\mathbb{R}$ : A set of all real numbers.
- $\mathbb{R}^+$ : A set of all positive real numbers.
- $\mathbb{C}$ : A set of all complex numbers.
- $A + B$ : A set given as  $\{a + b \mid a \in A, b \in B\}$ . For instance,  $\mathbb{Z}_2 + \mathbb{Z}_3 = \{(0 + 0), (0 + 1), (0 + 2), (1 + 0), (1 + 1), (1 + 2)\} = \{0, 1, 2, 3\}$

**Definition 2 (Subset)**

Given two sets  $A$  and  $B$ ,  $A$  is said to be a subset of  $B$  if the following condition holds.

If  $x$  is an element of  $A$ , then  $x$  is an element of  $B$ .

**Definition 3 (Extensionality)**

Two sets  $A$  and  $B$  are said to be equal if and only if  $A$  is a subset of  $B$  and  $B$  is a subset of  $A$ . This is called the *axiom of extensionality*. If two sets  $A$  and  $B$  are equal, we write  $A = B$ .

**Q1.**

EASY

Which of the following is distinct from the others?

Let  $A$  and  $B$  be two sets. Let  $A(x)$  mean  $x$  is an element of  $A$ , and  $B(x)$  mean  $x$  is an element of  $B$ .

Which of the following is equivalent to saying that  $A \subset B$ ?

- ☐ 1.  $B(x) \rightarrow A(x)$ 
☐ 2.  $\text{not } A(x) \rightarrow B(x)$ 
☐ 3.  $\text{not } A(x) \rightarrow \text{not } B(x)$   
☐ 4.  $A(x) \rightarrow \text{not } B(x)$ 
☐ 5.  $\text{not } B(x) \rightarrow \text{not } A(x)$

#### Definition 4 (Roots of unity)

Let  $\mu_n$  denote the set of all complex solutions to an equation of form  $x^n = 1$  for a positive integer  $n$ . Elements of such a set are called  $n$ -th roots of unity.

For instance,  $\mu_2 = \{1, -1\}$ .

Q2.

EASY

Show that  $\mu_2 \subset \mu_4$ .

Q3.

EASY

Prove or disprove that  $\mathbb{Z} \subset n\mathbb{Z} + m\mathbb{Z}$  for any two distinct positive integers  $n$  and  $m$ .

**Hint 5.** Provide concrete  $n$  and  $m$  that make up a counterexample. Try justifying your answer by using the definition of set builder notation.

Q4.

HARD

Prove or disprove that  $2\mathbb{Z} + 3\mathbb{Z} = \mathbb{Z}$ .

**Hint 6.** cf. lecture note.

Q5.

HARD

Show using [Definition 2](#) that for any two positive integers  $n$  and  $m$ ,  $\mu_n \subset \mu_{nm}$ .

**Hint 7.** Of course, it is a generalization of [Q2](#).

Q6.

HARD

Show by [Definition 3](#) that  $\mathbb{Z}_n + \mathbb{Z}_n = \mathbb{Z}_{2n-1}$  for any positive integer  $n$ .

**Hint 8.** cf. PCSA Q4.

HARD

Q7.

Prove or disprove that if  $\mathbb{Z} = n\mathbb{Z}$  for an integer  $n$ , then either  $n = 1$  or  $n = -1$ .

**Hint 9.** What are the factors of 1 and  $-1$ ?

### Definition 10 (Symmetric difference)

$A \triangle B$  is a binary set operation given by  $(A \setminus B) \cup (B \setminus A)$  and is called a *symmetric difference* between  $A$  and  $B$ .

Q8.

LEVI

Problem by @escentia07.

Prove or disprove that  $A \triangle B \subset C$  if and only if  $A \cup B \subset (A \cap B) \cup C$ .

Q9.

LEVI

For an integer  $n$ , let  $M(k)$  denote the set  $\{x \mid x \equiv k \pmod{n}\}$ . Let  $P$  the set given by  $P = \{M(k) \mid k \in \mathbb{Z}\}$ .

(a) Show that every two distinct elements of  $P$  are disjoint.

**Hint 11.** Use Definition 3 and *contraposition*.

(b) Show that the following holds for any two integers  $a$  and  $b$  by Definition 3.

$$M(a) + M(b) = M(a + b)$$

Q10.

LEVI

Problem by @escentia07.

Let  $f(x) = x^{2121} + x - 1$

How many real solutions does the equation  $f^{2026}(x) = x$  have? Justify your answer.

Answer: \_\_\_\_\_

Q11.

LEVI

Let  $T$  be the set  $\{z \in \mathbb{C} \mid |z| = 1\}$  and  $\mu_\infty$  be the set of all roots of unity as follows.

$$\mu_\infty = \bigcup_{n=1}^{\infty} \mu_n = \mu_1 \cup \mu_2 \cup \dots$$

Prove or disprove that  $T = \mu_\infty$ .

The statement is **× false**.



If you have learned set theory, you might have immediately realized that  $T$  is uncountable while  $\mu_\infty$  is countable.

Here goes the proof!

*Proof.* Let's first ensure that  $\mu_\infty$  is countable. We know that each  $\mu_n$  is finite as  $|\mu_n| = n$  and not empty as they always include 1, making  $\mu_\infty$  not empty. Now take any function  $f : \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mu_\infty$  such that if  $m \leq n$ , then  $f(n, m)$  is the  $m$ -th element of  $\mu_n$  when ordered by arguments. As such  $f$  always exists and is clearly surjective, we have  $|\mu_\infty| \leq |\mathbb{Z}^+ \times \mathbb{Z}^+| = \aleph_0$ . Hence,  $\mu_\infty$  is countable. On the other hand,  $T$  is uncountable as the argument  $\arg : T \rightarrow [0, 2\pi)$  is bijective, and  $[0, 2\pi)$  is uncountable. Since one is countable while the other is uncountable, they cannot be equal.  $\square$

Note that the above proof is perfectly valid even though it doesn't provide any counterexample. Nevertheless, I provide the below 'counterexample-based' disproof as well.

*Proof.* Let  $x = \frac{4}{5} + \frac{3}{5}i$ . It is clear that  $|x| = 1$  hence  $x \in T$ . You may have already realized that this counterexample is from Pythagorean numbers.

If  $x$  is an element of  $\mu_\infty$ , there must exist a positive integer  $n$  such that  $x^n = 1$ , otherwise put,  $(4 + 3i)^n = 5^n$ . Now, let  $(4 + 3i)^k = A_k + B_k i$  for some  $k \in \mathbb{Z}^+$  and find the next power.

$$(4 + 3i)^{k+1} = (A_k + B_k i)(4 + 3i) = (4A_k - 3B_k) + (3A_k + 4B_k)i$$

Thus,  $A_{k+1} = 4A_k - 3B_k$  and  $B_{k+1} = 3A_k + 4B_k$ . As  $A_1 = 4$  and  $B_1 = 3$ , we can ensure that every  $A_k, B_k$  are integers by induction. From here, it is not hard to show that  $A_k \equiv 3B_k \pmod{5}$  and  $B_{k+1} \equiv 3B_k \pmod{5}$ . As  $B_1 = 3$ , we can conclude that every  $B_k$  is essentially congruent to a power of 3, namely  $B_k \equiv 3^k \pmod{5}$ .

Since  $5^n$  is a real number, its imaginary part,  $B_n$ , must be 0 when  $k = n$ . However, the congruence equation  $3^n \equiv 0 \pmod{5}$  does not have a solution, as  $3 \not\equiv 0 \pmod{5}$ . Thus, such positive integer  $n$  does not exist, contradicting with the assumption. Therefore,  $x$  cannot be an  $n$ -th root of unity for any positive integer  $n$ .  $\square$

Keep in mind that multiplication of a complex number is a *rotation* on the complex plane. Thus, what we've just shown here is that the angle (*argument*) of  $x$  does not divide the circle evenly. Observe, or at least, feel how the problem of evenly dividing a circle turns into a problem of solving a solution to a

modular congruence. The roots of unity can be visualized as an exact rotations of a circle, justifying why they are called *cyclic groups*.

Of course, there is a shortcut to the above result. A corollary of the [Theorem 12](#) guarantees that for a rational number other than 0, 1, and  $-1$ , its arctan is always an irrational multiple of  $\pi$ . In the above case,  $\arctan(\frac{3}{4})$  should be an irrational multiple of  $\pi$ , which makes it never reaches 1 no matter how many times it rotates.

### Theorem 12 (Niven)

If  $\theta \in [0, \frac{\pi}{2}]$  and both  $\frac{\theta}{\pi}$  and  $\sin \theta$  are rational, then  $\sin \theta$  is either 0, 1, or  $\frac{1}{2}$ .

There is also a way more ‘algebraic’ proof as well.

*Proof.* Let  $x = \frac{4}{5} + \frac{3}{5}i$  and claim it’s an  $n$ -th root of unity for some  $n \in \mathbb{Z}^+$ . Hence we get  $(4 + 3i)^n = 5^n$ . As 5 can be factored into Gaussian integers  $(2 + i)(2 - i)$  and so on, we have the following prime factorization.

$$i^n(2 - i)^{2n} = (2 + i)^n(2 - i)^n$$

From here, the right hand side has  $(2 + i)$  as a prime  $n$  times, while the left hand side does not contain it. Recall that a prime factorizations by Gaussian integers must be unique up to units. As  $n \neq 0$  by the [Definition 4](#), this is a contradiction. Therefore,  $x$  is not an  $n$ -th root of unity for any positive integer  $n$ .  $\square$

From here, we have two different sets  $T$  and  $\mu_\infty$ . However, as you might have already guessed,  $\mu_\infty$  is a subset of  $T$ . If  $x$  is a solution to  $x^n = 1$  for some positive integer  $n$ , we have  $|x^n| = |1|$  by taking absoluates on each side, from which we get  $|x^n| = |x|^n = 1$ . The only real solution to it is  $|x| = 1$ .

In fact,  $\mu_\infty$  forms a *subgroup* of  $T$  under multiplication. Moreover, every  $\mu_n$  is itself a subgroups of  $\mu_\infty$  and called *cyclic* as every element of  $\mu_n$  can be written as an exponentiation of a *primitive  $n$ -th root of identity*. Finally,  $T$ , as you might have imagined, is called the *circle group*, and is also a subgroup of a much larger group  $U(\mathbb{C})$ , called the *unit group of the complex ring*.

$$\mu_n \leq \mu_\infty \leq T \leq U(\mathbb{C})$$

End.

You may now submit your work.

## Optional evaluation sheet

Unlike PCSA, submitting the sheet is **optional** for homeworks.

For each question so far, you can evaluate it based on the following criteria:

- If you failed to understand the question, mark  $\times$  in the Evaluation column.
- If you understood the question, but didn't know how to solve it, mark  $?$  in the Evaluation column.
- If you understood the question and are sure you have solved it, mark  $\circ$  in the Evaluation column.

| No. | Evaluation | No. | Evaluation |
|-----|------------|-----|------------|
| Q1  |            | Q7  |            |
| Q2  |            | Q8  |            |
| Q3  |            | Q9  |            |
| Q4  |            | Q10 |            |
| Q5  |            | Q11 |            |
| Q6  |            |     |            |

There are these additional questions as well; feel free to answer them.

- (a) Which was the hardest for you, among those you have solved? \_\_\_\_\_
- (b) Which seemed to be the hardest for you, among those you haven't solved? \_\_\_\_\_
- (c) Which was your favorite question, if you had to choose one? \_\_\_\_\_