

Math 351: Homework 10 Due Friday December 7

Jack Ellert-Beck

Chopping

Use Lemma 3.3.4 to complete the proof of the chopping lemma, Lemma 3.2.1

[Use angle difference formulas. Argue that the above argument for sin works for cos as well. Create a period of $2\pi/k$ with some error that disappears as $k \rightarrow \infty$ (that part might not quite be right)]

0.1 Jensen's

Exercise 5.1: Prove lemma 5.2.2.

Lemma: If ρ is convex on I , and $x_1, x_2 \in I$, and equality holds

$$\rho(wx_1 + (1-w)x_2) \leq w\rho(x_1) + (1-w)\rho(x_2)$$

for some $0 < w < 1$, then either $x_1 = x_2$ or ρ is linear on $[x_1, x_2]$.

[BY CONTRADICTION? CONFUSING]

Exercise 5.2: Prove lemma 5.2.3.

Lemma: If ρ is a convex function on an interval I , $x_1, \dots, x_n \in I$, and w_1, \dots, w_n are non-negative real numbers with $\sum w_i = 1$, then

$$\rho\left(\sum w_i x_i\right) \leq \sum w_i \rho(x_i)$$

with equality holding iff ρ is linear on an interval containing the x_i 's, or all x_i 's are the same.

We proceed by induction. The base case is where $n = 2$, which is true by Lemma 5.2.2.

[FINISH THIS ONE BOY]

Exercise 5.3: Prove that convex functions are continuous.

We will show that if a function $\rho : I \rightarrow \mathbb{R}$ (where $I = [a, b]$) has the property that, for $x_1, x_2 \in I$ and all $0 \leq w \leq 1$, $\rho(wx_1 + (1-w)x_2) \leq w\rho(x_1) + (1-w)\rho(x_2)$, then it is continuous on (a, b) , meaning that, at any $c \in I$, $\forall \varepsilon > 0 \exists \delta$ such that $|x - a| < \delta \implies |\rho(x) - \rho(c)| < \varepsilon$.

Pick any $\varepsilon > 0$. If $\varepsilon \leq \max(|\rho(x) - \rho(b)|, |\rho(a) - \rho(x)|)$, then any choice of δ satisfies the definition of continuity. So, we now consider only cases where $\varepsilon < \max(|\rho(x) - \rho(b)|, |\rho(a) - \rho(x)|)$.

0.2 Euler-Lagrange

Exercise 4.1: Use Euler-Lagrange to find the extremals for:

a) $\int_1^2 y'^2/x^3 dx$ with $y(1) = 2, y(2) = 17$

$$\begin{aligned} g(x, y, y') &= y'^2/x^3 \quad \text{and} \quad \frac{\partial g}{\partial y} - \frac{d}{dx} \left(\frac{\partial g}{\partial y'} \right) = 0 \\ &\implies 0 - \frac{d}{dx} \left(\frac{2y'}{x^3} \right) = 0 \\ &\implies \frac{2y'}{x^3} = C_0 \text{ for some constant } C_0 \\ &\implies y' = \frac{C_0}{2} x^3 \\ &\implies y = \int \frac{C_0}{2} x^3 dx = C_1 x^4 + C_2 \end{aligned}$$

so $1C_1 + C_2 = 2$ and $16C_1 + C_2 = 17 \implies C_1 = 1, C_2 = 1$

$$\implies y = x^4 + 1$$

b) $\int_0^{\pi/2} y^2 - y'^2 - 2y \sin(x) dx$ with $y(0) = 1, y(\pi/2) = 1$

$$\begin{aligned} g(x, y, y') &= y^2 - y'^2 - 2y \sin(x) \quad \text{and} \quad \frac{\partial g}{\partial y} - \frac{d}{dx} \left(\frac{\partial g}{\partial y'} \right) = 0 \\ &\implies (2y - 2 \sin(x)) - \frac{d}{dx} (-2y') = 0 \\ &\implies y'' = \sin(x) - y \\ &\implies y = C_2 \sin(x) + C_1 \cos(x) - \frac{1}{2} x \cos(x) \\ &\quad \text{(from Mathematica)} \end{aligned}$$

so $0C_2 + 1C_1 - 0 = 1$ and $1C_2 + 0C_1 - 0 = 1 \implies C_1 = 1, C_2 = 1$

$$\implies y = \sin(x) + \cos(x) - \frac{1}{2} x \cos(x)$$

$$\text{c) } \int_0^\pi y'^2 + 2y \sin(x) dx \text{ with } y(0) = 0, y(\pi) = 0$$

$$g(x, y, y') = y'^2 + 2y \sin(x) \quad \text{and} \quad \frac{\partial g}{\partial y} - \frac{d}{dx} \left(\frac{\partial g}{\partial y'} \right) = 0$$

$$\implies 2 \sin(x) - \frac{d}{dx} (2y') = 0$$

$$\implies y'' = \sin(x)$$

$$\implies y = -\sin(x) + C_1 x + C_2$$

$$\text{(from Mathematica)}$$

$$\text{so } 0C_1 + C_2 = 0 \text{ and } \pi C_1 + C_2 = 0 \implies C_1 = 0, C_2 = 0$$

$$\implies y = -\sin(x)$$