

CRIB parameter sensitivity: difference from the primary estimate

Group	N	70% required sleep				80% required sleep				90% required sleep			
		20 min	30 min	45 min	60 min	20 min	30 min	45 min	60 min	20 min	30 min	45 min	60 min
All adults	9,308	-0.28 (< 0.001)	-0.06 (< 0.001)	+0.22 (< 0.001)	+0.47 (< 0.001)	-0.22 (< 0.001)	0.00 (Ref.)	+0.27 (< 0.001)	+0.51 (< 0.001)	+0.15 (< 0.001)	+0.45 (< 0.001)	+0.79 (< 0.001)	+1.09 (< 0.001)
Sex													
Male	4,465	-0.30 (< 0.001)	-0.07 (< 0.001)	+0.23 (< 0.001)	+0.51 (< 0.001)	-0.23 (< 0.001)	0.00 (Ref.)	+0.29 (< 0.001)	+0.55 (< 0.001)	+0.19 (< 0.001)	+0.51 (< 0.001)	+0.87 (< 0.001)	+1.20 (< 0.001)
Female	4,843	-0.25 (< 0.001)	-0.05 (< 0.001)	+0.21 (< 0.001)	+0.44 (< 0.001)	-0.20 (< 0.001)	0.00 (Ref.)	+0.25 (< 0.001)	+0.47 (< 0.001)	+0.12 (< 0.001)	+0.40 (< 0.001)	+0.72 (< 0.001)	+1.00 (< 0.001)
Age													
18-34	2,643	-0.23 (< 0.001)	-0.05 (< 0.001)	+0.18 (< 0.001)	+0.39 (< 0.001)	-0.18 (< 0.001)	0.00 (Ref.)	+0.22 (< 0.001)	+0.42 (< 0.001)	+0.19 (< 0.001)	+0.46 (< 0.001)	+0.78 (< 0.001)	+1.06 (< 0.001)
35-44	1,475	-0.23 (< 0.001)	-0.05 (< 0.001)	+0.18 (< 0.001)	+0.39 (< 0.001)	-0.18 (< 0.001)	0.00 (Ref.)	+0.22 (< 0.001)	+0.41 (< 0.001)	+0.15 (< 0.001)	+0.41 (< 0.001)	+0.71 (< 0.001)	+0.98 (< 0.001)
45-64	3,102	-0.28 (< 0.001)	-0.06 (< 0.001)	+0.22 (< 0.001)	+0.47 (< 0.001)	-0.22 (< 0.001)	0.00 (Ref.)	+0.27 (< 0.001)	+0.51 (< 0.001)	+0.14 (< 0.001)	+0.43 (< 0.001)	+0.77 (< 0.001)	+1.06 (< 0.001)
≥65	2,088	-0.38 (< 0.001)	-0.08 (< 0.001)	+0.32 (< 0.001)	+0.67 (< 0.001)	-0.30 (< 0.001)	0.00 (Ref.)	+0.38 (< 0.001)	+0.71 (< 0.001)	+0.14 (< 0.001)	+0.51 (< 0.001)	+0.94 (< 0.001)	+1.31 (< 0.001)
BMI													
Underweight	187	-0.21 (< 0.001)	-0.04 (< 0.001)	+0.23 (< 0.001)	+0.53 (< 0.001)	-0.18 (< 0.001)	0.00 (Ref.)	+0.26 (< 0.001)	+0.54 (< 0.001)	+0.19 (< 0.001)	+0.47 (< 0.001)	+0.83 (< 0.001)	+1.16 (< 0.001)
Normal weight	2,713	-0.25 (< 0.001)	-0.05 (< 0.001)	+0.21 (< 0.001)	+0.44 (< 0.001)	-0.20 (< 0.001)	0.00 (Ref.)	+0.25 (< 0.001)	+0.47 (< 0.001)	+0.14 (< 0.001)	+0.41 (< 0.001)	+0.73 (< 0.001)	+1.00 (< 0.001)
Overweight	2,899	-0.27 (< 0.001)	-0.06 (< 0.001)	+0.22 (< 0.001)	+0.45 (< 0.001)	-0.21 (< 0.001)	0.00 (Ref.)	+0.26 (< 0.001)	+0.48 (< 0.001)	+0.15 (< 0.001)	+0.44 (< 0.001)	+0.77 (< 0.001)	+1.06 (< 0.001)
Obesity	3,407	-0.30 (< 0.001)	-0.07 (< 0.001)	+0.24 (< 0.001)	+0.51 (< 0.001)	-0.24 (< 0.001)	0.00 (Ref.)	+0.29 (< 0.001)	+0.55 (< 0.001)	+0.17 (< 0.001)	+0.49 (< 0.001)	+0.86 (< 0.001)	+1.18 (< 0.001)
Race/ethnicity													
Mexican American	1,135	-0.23 (< 0.001)	-0.05 (< 0.001)	+0.20 (< 0.001)	+0.43 (< 0.001)	-0.19 (< 0.001)	0.00 (Ref.)	+0.24 (< 0.001)	+0.46 (< 0.001)	+0.18 (< 0.001)	+0.47 (< 0.001)	+0.82 (< 0.001)	+1.12 (< 0.001)
Other Hispanic	892	-0.25 (< 0.001)	-0.06 (< 0.001)	+0.18 (< 0.001)	+0.40 (< 0.001)	-0.19 (< 0.001)	0.00 (Ref.)	+0.23 (< 0.001)	+0.45 (< 0.001)	+0.16 (< 0.001)	+0.43 (< 0.001)	+0.76 (< 0.001)	+1.04 (< 0.001)
NH White	3,720	-0.28 (< 0.001)	-0.06 (< 0.001)	+0.23 (< 0.001)	+0.48 (< 0.001)	-0.22 (< 0.001)	0.00 (Ref.)	+0.27 (< 0.001)	+0.51 (< 0.001)	+0.13 (< 0.001)	+0.43 (< 0.001)	+0.76 (< 0.001)	+1.04 (< 0.001)
NH Black	2,217	-0.29 (< 0.001)	-0.07 (< 0.001)	+0.22 (< 0.001)	+0.51 (< 0.001)	-0.22 (< 0.001)	0.00 (Ref.)	+0.28 (< 0.001)	+0.55 (< 0.001)	+0.23 (< 0.001)	+0.57 (< 0.001)	+0.97 (< 0.001)	+1.33 (< 0.001)
NH Asian	1,072	-0.29 (< 0.001)	-0.07 (< 0.001)	+0.22 (< 0.001)	+0.49 (< 0.001)	-0.23 (< 0.001)	0.00 (Ref.)	+0.27 (< 0.001)	+0.53 (< 0.001)	+0.21 (< 0.001)	+0.54 (< 0.001)	+0.92 (< 0.001)	+1.26 (< 0.001)
Other/Multi	272	-0.27 (< 0.001)	-0.07 (< 0.001)	+0.20 (< 0.001)	+0.47 (< 0.001)	-0.21 (< 0.001)	0.00 (Ref.)	+0.25 (< 0.001)	+0.52 (< 0.001)	+0.14 (< 0.001)	+0.44 (< 0.001)	+0.79 (< 0.001)	+1.09 (< 0.001)

Cells show the survey-weighted paired difference in waking sedentary hours/day (two-sided p-value), alternative setting minus the primary 80%/30-minute setting. Positive values indicate more estimated sedentary time. All 12 settings use the same 9,308 adults and 55,722 primary-eligible days; days are averaged within participant before survey estimation. N is the unweighted subgroup count. P-values use the survey-design t distribution for the mean within-participant difference and are unadjusted for multiple comparisons. Ref. denotes the identical reference setting, for which no test is performed. BMI is missing for 102 participants. Eligibility is fixed to the primary analysis: alternative settings do not select a new cohort.