

Exercise Solutions for Math 20

Functions, Graphs, Symmetry

Nile Jocson <novoseiversia@gmail.com>

November 20, 2024

1

1.1 Transform each of the following real-valued functions to a functional notation $f(x)$; find its domain and range; and compute $f(1)$.

1.1.a f assigns the number 10 to any one-digit integer.

$$\Rightarrow f(x) = 10$$

Final answer.

$$\Rightarrow \text{dom}(f) = [-9, 9]$$

The domain is the set of one digit integers.

$$\Rightarrow \text{ran}(f) = 10$$

The function can only equate to 10.

$$\Rightarrow f(1) = 10$$



1.1.b f maps a given real number to the real number that is 4 more than its square root.

$$\Rightarrow f(x) = \sqrt{x} + 4$$

Final answer. Assuming that only the principal root is needed.

$$\Rightarrow \text{dom}(f) = [0, +\infty)$$

The square root of a negative number cannot be real.

$$\Rightarrow \text{ran}(f) = [4, +\infty)$$

$$\Rightarrow f(1) = 5$$



1.1.c $\{(x, y) \mid y = \frac{1}{6-x}\}$

$$\Rightarrow f(x) = \frac{1}{6-x}$$

Final answer.

$$\Rightarrow \text{dom}(f) = \mathbb{R} \setminus \{6\}$$

$x = 6$ is an undefined point.

$$\Rightarrow \text{ran}(f) = \mathbb{R}$$

$$\Rightarrow f(1) = \frac{1}{5}$$



1.1.d $\{(x, y) \mid y = \sqrt{x^2 - 5x + 6}\}$

$$\Rightarrow (x - 2)(x - 3) \geq 0$$

Find the domain by solving $x^2 - 5x + 6 \geq 0$; factor by grouping.

Create a table of signs.

	2	3	
$x - 2$	−	+	+
$x - 3$	−	−	+
$(x - 2)(x - 3)$	+	−	+

$$\Rightarrow x \in (-\infty, 2] \cup [3, +\infty)$$

$$\Rightarrow f(x) = \sqrt{x^2 - 5x + 6}$$

Final answer.

Continued on next page

$$\Rightarrow \text{dom}(f) = (-\infty, 2] \cup [3, +\infty)$$

$$\Rightarrow \text{ran}(f) = [0, +\infty)$$

$$\Rightarrow f(1) = \sqrt{2}$$

■

1.1.e $\{(x, y) \mid y = \frac{1}{\sqrt[3]{x^2-1}}\}$

$$\Rightarrow (x-1)(x+1) = 0$$

Find undefined points by solving $x^2 - 1 = 0$;
factor using difference of two squares.

$$\Rightarrow x \in \{-1, 1\}$$

$$\Rightarrow x = \frac{1}{\sqrt[3]{y^2-1}}$$

Find the inverse function.

$$\Rightarrow \frac{1}{x} = \sqrt[3]{y^2-1}$$

$$\Rightarrow \frac{1}{x^3} = y^2 - 1$$

$$\Rightarrow y^2 = \frac{1}{x^3} + 1$$

$$\Rightarrow y^2 = \frac{1}{x^3} + \frac{x^3}{x^3}$$

$$\Rightarrow y^2 = \frac{1+x^3}{x^3}$$

$$\Rightarrow f'(x) = \pm \sqrt{\frac{1+x^3}{x^3}}$$

$$\Rightarrow \sqrt{\frac{1+x^3}{x^3}} > 0$$

Solve for the domain. Note that $x = 0$ is an
undefined point.

Create a table of signs.

	-1	0	
$1 + x^3$	-	+	+
x^3	-	-	+
$\sqrt{\frac{1+x^3}{x^3}}$	+	-	+

$$\Rightarrow \text{dom}(f') = (-\infty, -1] \cup (0, +\infty)$$

$$\Rightarrow f(x) = \frac{1}{\sqrt[3]{x^2-1}}$$

Final answer.

$$\Rightarrow \text{dom}(f) = \mathbb{R} \setminus \{-1, 1\}$$

$$\Rightarrow \text{ran}(f) = (-\infty, -1] \cup (0, +\infty)$$

$$\Rightarrow f(1) \text{ is undefined.}$$

■