Interpretation of Liver Stiffness in kPa[1-5]					
Liver Stiffness for healthy individual without liver disease (95% of ND)[1]		2.3 kPa - 5.9 kPa*			
		No - Probable Minimal Fibrosis (F0-F1)	**Borderline Fibrosis (F2)**	**Probable Advanced Fibrosis (F3)**	**Probable Cirrhosis (F4)**
Hepatitis B[2]	**Normal ALT**	≤ 6.0	> 6.0 - 9.0	> 9.0 - 12.0	> 12.0
	Elevated ALT^	≤ 7.5	> 7.5 - 12.0	> 12.0 - 13.4	> 13.4
Hepatitis C[3]		≤ 7.0	> 7.0 - 9.5	> 9.5 - 12.5	> 12.5
NAFLD[4]		≤ 5.8	> 5.8 - 9.6	> 9.6 - 11.5	> 11.5
Alcoholic Liver Disease[5]		< 5.9	> 7.8	> 11	> 19.5

Interpretation of Controlled Attenuation Parameter (CAP) in dB/m[6]				
Liver Steatosis (% affected hepatocytes)	**No - Minimal (S0) (< 10%#)**	**Mild (S1) (10#-33%)**	**Moderate (S2) (34-66%)**	**Severe (S3) (>66%)**
CAP	< 248	248 - 267	268-279	≥ 280

* For liver stiffness result ≥ 6 kPa, possible fibrotic liver or other abnormal liver condition is suggested which requires further investigation.

^ **ALT** = Alanine aminotransferase. (Liver stiffness measurement likely overestimates the severity of liver fibrosis when ALT is higher than five times the upper limit of laboratory normal reference value)

Depending on the trial – some trials define S0 as <5% also.

References:

1. Fung J, Lee C, Chan M, Seto W, Wong DK, Lai C, & Yuen M. (2013). Defining normal liver stiffness range in a normal healthy Chinese population without liver disease. *Plos ONE*, *8*(12), doi:10.1371/journal.pone.0085067

2. Chan HLY, Wong GLH, Choi PCL, Chan AWH, Chim AML, Yiu KKL, Chan FKL, Sung JJY, & Wong VWS. (2009). Alanine aminotransferase-based algorithms of liver stiffness measurement by transient elastography (Fibroscan) for liver fibrosis in chronic hepatitis B. *Journal of Viral Hepatitis, 16,* pp. 36-44

3. Castera L, Vergniol J, Foucher J, Le Bail B, Chanteloup E, Haaser M, Darriet M, Couzigou P, & de Ledinghen V. (2005). Prospective comparison of transient elastography, Fibrotest, APRI, and liver biopsy for the assessment of fibrosis in chronic hepatitis C. *Gastroenterology, 128* (2), pp. 343-350.

4. Wong VWS, Vergniol J, Wong GLH, Foucher J, Chan HLY, Bail BL, Choi PCL, Kowo M, Chan AWH, Merrouche W, Sung JJY, & de Ledinghen V. (2010). Diagnosis of fibrosis and cirrhosis using liver stiffness measurement in nonalcoholic fatty liver disease. *Hepatology, 51*(2), pp. 454-462

5. Nguyen-Khac E, Chatelain D, Tramier B, Decrombecque C, Robert B, Joly J, Brevet M, Grignon P, Lion S, le Page L, & Dupas J. (2008). Assessment of asymptomatic liver fibrosis in alcoholic patients using fibroscan: prospective comparison with seven non-invasive laboratory tests. *Alimentary Pharmacology & Therapeutics*, *28*(10), 1188-1198. doi:10.1111/j.1365-2036.2008.03831.x

6. Karlas T, Petroff D, Sasso M, Fan JG, Mi YQ, de Ledinghen V, Kumar M, Lupsor-Platon M, Han KH, Cardoso AC, Ferraioli G, Chan WK, Wong VWS, Myers RP, Chayama K, Friedrich-Rust M, Beaugrand M, Shen F, Hiriart JB, Sarin SK, Badea R, Jung KS, Marcellin P, Filice C, Mahadeva S, Wong GLH, Crotty P, Masaki K, Bojunga J, Bedossa P, Keim V, Wiegand J. (2017). Individual patient data meta-analysis of controlled attenuation parameter (CAP) technology for assessing steatosis. *Journal of Hepatology*, *66* (5), pp. 1022-1030