



## Cue-alcohol associative learning in female rats

Roberto U. Cofresí<sup>a,c</sup>, Marie-H. Monfils<sup>a,b</sup>, Nadia Chaudhri<sup>d</sup>, Rueben A. Gonzales<sup>a,c</sup>, Hongjoo J. Lee<sup>a,b,\*</sup>

<sup>a</sup> The University of Texas at Austin, Institute for Neuroscience, Austin, TX, 78712, United States

<sup>b</sup> The University of Texas at Austin, College of Liberal Arts, Department of Psychology, Austin, TX, 78712, United States

<sup>c</sup> The University of Texas at Austin, College of Pharmacy, Division of Pharmacology & Toxicology, Psychology, Austin, TX, 78712, United States

<sup>d</sup> Concordia University, Department of Psychology, Center for Studies in Behavioral Neurobiology, Montreal, Quebec, H4B1R6, Canada

### ARTICLE INFO

#### Article history:

Received 21 January 2019

Received in revised form

14 March 2019

Accepted 19 March 2019

#### Keywords:

Oral alcohol

Low dose

Pavlovian conditioning

Female rat

Cue reactivity

### ABSTRACT

The ability of environmental cues to trigger alcohol-seeking behaviors is believed to facilitate problematic alcohol use. We previously showed that the development of this cue-evoked alcohol approach reflects cue-alcohol learning and memory in the adult male rat; however, we do not know whether the same is true for similarly aged female rats. Consequently, adult Long-Evans female rats were allowed to drink unsweetened alcohol in the home cage (Monday, Wednesday, Friday; 24-h two-bottle choice; 5 weeks) and were subsequently split into two experimental groups: Paired and Unpaired. Groups were matched for ingested doses and alcohol bottle preference across the pre-conditioning home cage period. Both groups were trained in conditioning chambers using a Pavlovian procedure. For the Paired group, the chamber houselight was illuminated to signal access to an alcohol sipper. Houselight onset was yoked for the Unpaired group, but access to the alcohol sipper was scheduled to occur only during the intervening periods (in the absence of light). We found that in the Paired, but not Unpaired group, an alcohol approach reaction was conditioned to houselight illumination, and the level of cue-conditioned reactivity predicted drinking behavior within trials. Groups experienced equivalently low but non-negligible blood alcohol concentrations over the course of conditioning sessions. We conclude that cue-triggered alcohol-seeking behavior in adult female rats reflects associative learning about the relationship between alcohol availability and houselight illumination.

© 2019 Elsevier Inc. All rights reserved.

### Introduction

Environmental stimuli that have been routinely paired with alcohol can acquire the ability to trigger alcohol-seeking behaviors and thereby contribute to problematic alcohol use. The implicit associative learning process that allows environmental stimuli paired with alcohol to acquire the ability to trigger alcohol-seeking behaviors, Pavlovian or classical cue conditioning, is believed to operate in a fundamentally similar way in males and females. However, males and females may differ in levels of susceptibility to specific ways in which cues can contribute to problematic alcohol use (Barker & Taylor, 2017). In the field of preclinical non-human animal models, there is a growing appreciation for qualitative and quantitative differences in the processes contributing to addiction-

like behavior and its expression (e.g., drug cue learning, drug cue reactivity) between male and female individuals (Becker & Koob, 2016). In light of this growing appreciation of biological sex differences, the burden of proof is on researchers to demonstrate that our models of addiction-like behavioral phenomena in non-human animals operate similarly in males and females of the model organism species, and if not, to document the differences. Despite this, many preclinical studies of alcohol-cue conditioning, especially those that use rats as the model organism and voluntary alcohol drinking paradigms, including our own (Cofresí et al., 2019; , 2017; Cofresí, Lee, Monfils, Chaudhri, & Gonzales, 2018; Knight et al., 2016; ; Krank, 2003; Krank, O'Neill, Squarey, & Jacob, 2008; Lamb, Ginsburg, Greig, & Schindler, 2019; Lamb, Ginsburg, & Schindler, 2016; LeCocq, Lahlou, Chahine, Padillo, & Chaudhri, 2018; Millan, Reese, Grossman, Chaudhri, & Janak, 2015; Sparks, Sciascia, Ayorech, & Chaudhri, 2014; Srey, Maddux, & Chaudhri, 2015; Tomie, Festa, Sparta, & Pohorecky, 2003; Tomie, Kuo, Apor, Salomon, & Pohorecky, 2004; Tomie, Miller, Dranoff, & Pohorecky,

\* Corresponding author. 108 E. Dean Keeton Stop, A8000, Austin, Texas, 78712, United States. Tel.: +1 512 232 8055.

E-mail address: [leejh@austin.utexas.edu](mailto:leejh@austin.utexas.edu) (H.J. Lee).

2006; Villaruel & Chaudhri, 2016), were conducted exclusively in male rats, and therefore, little is known about how Pavlovian alcohol-cue conditioning proceeds in female rats.

Here, we determined whether female rats were capable of associating an environmental stimulus with alcohol using the two-stage paradigm that we initially developed in male rats. In the first stage, we provided intermittent 24-h access to unsweetened alcohol in the rat's home cage alongside free access to food and water for 5 weeks. In the second stage, we tested the ability of time-limited unsweetened alcohol drinking opportunities to condition alcohol-seeking behavior to an antecedent visual cue in a physical environment different from the rat's home cage. In the latter test, a persistent change in the behavioral response to cue presentation could have been due to learning to associate the cue with alcohol access or non-associative learning driven by repeated exposure to the cue or to alcohol. To distinguish between these possibilities, we characterized behavior during cue presentation in female rats that were trained on two versions of the same conditioning paradigm. In one, alcohol access was explicitly paired with houselight illumination (a visual cue), whereas in the other, the two events were explicitly unpaired. Behavioral changes observed in the Paired group, but not observed in the Unpaired group, reflect alcohol-associative learning about the visual cue. To examine whether persistent behavioral changes during visual cue presentation were driven by differences in the ability of rats in the Paired and Unpaired groups to consume alcohol during the visual cue conditioning sessions, we characterized consummatory behavior (sipper licking latency and intensity). To verify that rats in both groups had similar blood alcohol concentrations during visual cue conditioning sessions, we took blood samples at the end of a conditioning session, determined the relationship between ingested dose and blood alcohol concentration, and retrospectively predicted blood alcohol concentration at the end of each conditioning session as a function of ingested dose. In doing so, we also evaluated the extent to which changes in behavior across visual cue conditioning might be driven by alcohol's post-ingestive pharmacology. Finally, we tested whether conditioned behavioral reactivity to the visual cue for alcohol in the paired group predicted alcohol consummatory behavior and ingested dose, two predictions derived from Tomie's model for how alcohol-cue reactivity promotes problematic alcohol use (Tomie, 1996; Tomie & Sharma, 2013).

## Methods and materials

### Subjects

Subjects were adult female Long-Evans rats (Envigo; Indianapolis, Indiana) weighing 200–225 g at arrival. Rats were singly housed in shoebox-style Plexiglas® home cages containing Sanichips® bedding and a Bio-Serv Gummy Bone (polyurethane; 5 cm × 2.5 cm). Metal wire cage tops were used. Standard chow pellets were loaded into a large cup inside the cage. Tap water was provided via gravity-fed sipper inserted at approximately 45° from the cage top. Chow and water were replenished daily. Bedding was replaced weekly. Cages were located in a temperature- and humidity-controlled room (22 ± 2 °C). All procedures took place 4–5 h into the light phase of a 12-h light/dark cycle unless otherwise indicated. Drinking solutions were prepared from 95% ethyl alcohol (ACS/USP grade, Pharmco-AAPER) and tap water every 3 days. These were kept and served at room temperature (20 °C). All procedures were approved by the Institutional Animal Care and Use Committee at the University of Texas at Austin, and conducted in accordance with NIH guidelines.

### Pre-conditioning ethanol drinking in the home cage

This procedure was described in detail elsewhere (Cofresí et al., 2017, 2018; Sparks et al., 2014). Briefly, rats were provided a bottle of unsweetened ethanol (15% ethanol in tap water; v/v; 15E) and a new bottle of water for 24 h every Monday, Wednesday, and Friday for 5 weeks. Bottle placement on the cage top alternated (ethanol on left v. right side) across sessions. Rats that failed to drink in week 1 were provided 5% and then 10% ethanol in tap water (v/v; 5E, 10E) to promote drinking. Any rats drinking <1 g/kg/24 h across week 5 were not retained for conditioning.

### Ethanol-reinforced classical conditioning

The conditioning chambers used were described in Cofresí et al. (2017). The conditioning procedures were previously described (Cofresí et al., 2019). Briefly, rats were assigned to “Paired” or “Unpaired” conditioning such that the resulting groups were matched for ingested doses across the 5 weeks of pre-conditioning drinking sessions. Rats in both groups were trained to use the retractable ethanol sipper in the conditioning chamber and habituated to chamber houselight illumination. Rats then underwent cue conditioning across 12 consecutive days. Each conditioning session consisted of eight trials. The inter-trial interval (ITI) was variable (mean: 280 s, minimum: 160 s, maximum: 360 s). After a 5-min wait period, the session started (with the first ITI). This was signaled to the rat by onset of the exhaust fan. The session ended when the final ITI (selected after trial 8) elapsed. This was signaled by offset of the exhaust fan inside each cubicle. During each trial in a session, the chamber houselight was illuminated for 20 s. In the Unpaired group, there was no consequent event. In the Paired group, the retractable bottle assembly was activated 10 s into the illumination to present a metal sipper such that ethanol access and houselight illumination co-terminated. Sipper presentations for the Unpaired group occurred mid-ITI, beginning in ITI 2 and ending in ITI 9. Houselight illumination onset, offset, and ITIs were yoked between groups. Sipper presentations were yoked within groups. Licking the sipper produced 10E or 15E, whichever the rat was drinking at the end of the pre-conditioning phase.

### Blood collection and ethanol analysis

After the 12th conditioning session, 1–2 additional sessions were given. At the end of one of these sessions, blood was sampled from the lateral saphenous vein while the rat was under isoflurane anesthesia. Ethanol concentration (mg/dL) in the blood sample was determined using gas chromatography with flame ionization detection as in Carrillo et al. (2008) and Cofresí et al. (2018, 2019).

### Behavior measurement

Trials from conditioning sessions 1–12 were sampled for behavior from digital video recordings. As in Cofresí et al. (2017, 2018, 2019) and Lee et al. (2005), instantaneous observations were made every 1.25 s starting 5 s before houselight illumination, such that there were four observations per 5-sec bin across the illumination period. At each observation, the rater noted the absence or presence of mutually exclusive behavioral states (sipper site approach: approaching or exploring the sipper insertion hole; orienting to light: both forepaws off the floor, supported by hind limbs; other: grooming, resting). Each of the 5-sec bins corresponded to a meaningful trial phase. Since the original paradigm (Paired group) was designed with houselight illumination as the

conditional stimulus (CS), the bins were labeled with reference to the CS. The pre-CS bin is the 5-sec period before CS presentation, and CS bins 1–4 are the 5-sec periods across CS presentation. Sipper licking was automatically recorded using a contact lickometer circuit. The latency(s) to first lick was also recorded on every trial. If no lick was registered within 10 s of sipper onset, then a maximum latency of 10 s was recorded. A second, modified contact lickometer circuit was used to automatically record forepaw contacts with the area of the chamber wall immediately around the sipper insertion hole independently of sipper licking, as described in Cofresí et al. (2019). The latency(s) to first forepaw contact after houselight onset was recorded. If no forepaw contact was registered within 30 s of houselight onset, then a maximum latency of 30 s was recorded. The latency(s) to first forepaw contact after sipper onset was also recorded. If no forepaw contact was registered within 20 s of sipper onset, then a maximum latency of 20 s was recorded. For both post-houselight and post-sipper onset forepaw contact latencies, if sustained forepaw contact was ongoing at the time of houselight/sipper onset (namely, if the rat was “holding on” to the area around the sipper insertion hole), then a negative latency was recorded because initiation of on-going contact was at an earlier time than onset of the houselight/sipper. Infrared photo beams were used to track general locomotion in the stimulus-rich (houselight fixture and sipper hole) vs. stimulus-poor (bare wall) zones of the conditioning chamber.

The dose of ethanol ingested by each rat was also monitored. For every home cage drinking session, bottles on an empty control cage were used to measure loss due to evaporation and spillage and correct drinking solution intake values across all subjects. For every conditioning session, a weigh boat underneath each bottle assembly collected spillage, and drinking solution intake values were corrected at the level of each individual subject. Drinking solution intake was measured as the corrected mass difference in bottle weight pre- and post-session. To obtain the ingested ethanol dose, the amount (g) of pure ethanol consumed was computed and expressed relative to body weight (kg) of each rat.

### Statistical analysis

Mixed factorial analysis of variance (ANOVA) was the primary statistical analysis technique used to analyze behavioral and drinking data. The threshold for statistical significance was  $p < 0.05$ . Significant results in the omnibus ANOVA were followed up as appropriate (e.g., ANOVA  $F$  tests were used to decompose interactions of two or more factors, and  $t$  tests were used to decompose the main effect of a factor). Bonferroni correction was applied at every follow-up stage to minimize false discovery. In a few instances, we used other statistical procedures. However, the threshold for statistical significance remained at  $p < 0.05$  for these other analyses. For example, we used Pearson's correlation test to evaluate the relationship between blood ethanol and ingested dose.

All analysis was done in R version 3.5.1 (R Core Team, 2018) using the car package (Fox & Weisberg, 2011). Data were plotted in R using the ggplot2 package (Wickham, 2009) and finalized in Inkscape version 0.92.2 (Inkscape Team, 2017).

### Results

Of 20 rats obtained for the study, 19 were conditioned based on *a priori* retention criterion: ingested dose  $\geq 1$  g/kg/24 h on average across the last week of the pre-conditioning phase. Pre-conditioning home cage ethanol drinking data are presented in Supplemental Figure 1.

Of the 19 rats that were conditioned, 17 were retained based on our *a priori* inclusion criterion: ingested dose  $\geq 0.30$  g/kg/session on average across the last three conditioning sessions. The two rats that failed to meet the latter criterion were both in the Unpaired group. One of those two had been conditioned with 10E, and the other had been conditioned with 15E. Of the 17 that met our *a priori* inclusion criterion, all seven in the Unpaired group and eight out of 10 in the Paired group had been conditioned with 15E. The remaining two out of 10 rats in the Paired group that met our *a priori* inclusion criterion had been conditioned with 10E.

During the waiting period before the first conditioning session, rats in the Paired group and Unpaired group alike were more active in the stimulus-rich half of the conditioning chamber (i.e., with the houselight fixture and sipper insertion hole) than the stimulus-poor half, and this did not change over the course of conditioning (Supplemental Fig. 2A). Rats in the Paired group and Unpaired group alike, however, did make increasingly more forepaw contacts with the area around the sipper insertion hole during the pre-session waiting period over the course of conditioning (Supplemental Fig 2B).

### Drinking in the conditioning chamber across ethanol-reinforced classical conditioning


Rats in the Paired and Unpaired groups drank similarly across the conditioning phase. Ingested doses increased significantly across conditioning sessions [session main effect:  $F(11,165) = 26.08$ ,  $p < 0.001$ ], and the pattern of increase was similar between groups (group main effect and group  $\times$  session interaction: NS; Figure 1A).

Seventeen rats met our *a priori* minimum drinking criteria across conditioning sessions 10–12. We wanted to monitor the blood ethanol concentrations achieved after the conditioning sessions, but we wanted to avoid the possible effects of the invasive blood sampling procedure on the behavior in subsequent sessions. Therefore, the rats were exposed to one or two additional conditioning sessions, and blood was sampled 8–11 min after the 8th sipper presentation. Blood ethanol concentration (BEC) at the end of the conditioning session was significantly related to ingested dose (Pearson's  $r = +0.76$ ,  $t_{15} = 4.60$ ,  $p < 0.001$ ; Figure 1B). Body weights ranged from 260 to 318 g. Ingested dose ranged from 0.35 to 1.2 g/kg, with a mean  $\pm$  S.E.M. of  $0.72 \pm 0.05$  g/kg. BECs ranged from 0 to 38.5 mg/dL, with a mean  $\pm$  S.E.M. of  $15.9 \pm 3.4$  mg/dL. The Paired and Unpaired groups did not differ in BEC, ingested dose, the relationship between dose and BEC, sampling time, or body weight on blood sampling day (Table 1). Thus, a single simple regression equation was used to predict BEC as a function of ingested dose across the conditioning sessions. Estimated end of session BEC across conditioning sessions did not differ between the Paired and Unpaired groups [session main effect:  $F(11,165) = 10.67$ ,  $p < 0.001$ ; group main effect and group  $\times$  session interaction: NS; Figure 1C]. Overall, end of session estimated BEC were low but non-zero after session 6.


### Acquisition of houselight cue-triggered ethanol seeking in the paired group, but not the unpaired group

The Paired and Unpaired groups differed in sipper site approach frequency during the trial phases across training [group  $\times$  session interaction:  $F(11,165) = 5.94$ ,  $p < 0.001$ ; group  $\times$  trial phase interaction:  $F(2,30) = 5.18$ ,  $p < 0.05$ ; Figure 2A]. For rats in the Paired group, sipper site approach frequency increased over sessions [simple session effect:  $F(11,99) = 12.03$ ,  $p < 0.001$ ] and as a function of houselight illumination period [simple trial phase effect:  $F(2,18) = 4.78$ ,  $p < 0.05$ ; pairwise  $t$  tests for pre-CS bin  $<$  CS bin 1 and CS bin 1  $<$  CS bin 2:  $t_9 > 5.25$ ,  $p < 0.0001$ ]. In contrast, sipper site


## A. Ingested Doses Across Conditioning



## B. Measured Post-Session Blood Ethanol



## C. Estimated Post-Session Blood Ethanol



**Fig. 1. Equivalent ethanol exposure across cue conditioning.** A: Ingested dose (g/kg) per session shown across conditioning sessions in all the animals. Horizontal line shows *a priori* inclusion criterion (dose  $\geq 0.30$  g/kg/session across sessions 10–12). B: Relationship between blood ethanol concentrations detected approximately 10 min after the 8th 10-sec drinking opportunity in a conditioning session and total ingested ethanol doses in the same session for adult, female Long-Evans rats. Black and white triangles represent Paired group ( $n = 10$ ) and Unpaired group ( $n = 7$ ), respectively. Regression line and 95% confidence limits shown by solid line and shaded area, respectively. C: Mean  $\pm$  S.E.M. estimated blood ethanol concentrations across conditioning sessions (approximately 10 min after the 8th drinking opportunity in the session) using ingested doses from A and the regression equation from B for the same 17 rats.

approach frequency remained at floor across trial phases and sessions for rats in the Unpaired group (simple session & trial phase effects: NS). The difference in sipper site approach level was

clearest in CS bin 2 [simple group effect:  $F(1,15) = 19.78$ ,  $p < 0.001$ ; Figure 2A, rightmost panel].

To confirm these findings, we also analyzed sipper site (face-plate) contact frequency, which was measured automatically using a modified lickometer circuit, and thus, free of rater bias. Results were similar to those presented above. The Paired and Unpaired groups differed in sipper site contact frequency across training [group main effect:  $F(1,15) = 5.78$ ,  $p < 0.03$ ; session main effect:  $F(11,166) = 4.18$ ,  $p < 0.001$ ; group  $\times$  session interaction:  $F(11,165) = 5.16$ ,  $p < 0.001$ ], but the group  $\times$  trial phase interaction effect was not statistically significant [ $F(2,30) = 1.81$ , NS]. However, it can be seen in Figure 2B that for rats in the Paired group, sipper site contact frequency during CS bin 1 and 2 increased over sessions, whereas contact during the pre-CS bin remained at the base level. In contrast, sipper site contact frequency remained at the base level across sessions in every bin for rats in the Unpaired group (Fig. 2B).

The frequency of houselight illumination-elicited orienting across sessions is presented in Supplemental Figure 3.

### Houselight cue-elicited ethanol-seeking reaction dynamics in session 12

Our previous studies in the Paired group male rats found that the ability of the houselight cue to elicit ethanol seeking appears to decrease across trials within sessions reliably by conditioning session 12 (Cofresí et al., 2018, 2019). In order to verify whether the same behavior pattern occurs in the Paired group female rats, we examined trial-by-trial behavior in conditioning session 12. Overall, female rats in the Paired group exhibited a robust sipper site approach reaction to houselight illumination in session 12, whereas those in the Unpaired group did not [group main effect:  $F(1,15) = 13.93$ ,  $p < 0.003$ ; Figure 3A]. Focusing on trial phase CS2, sipper site approach frequency was greater for the Paired group than for the Unpaired group in every trial ( $t_{15} \geq 2.10$ ,  $p \leq 0.05$ ) (Fig. 3A).

Similar results were obtained when we analyzed per-trial sipper site contacts (i.e., forepaw contact with the faceplate around the sipper insertion hole) in session 12. Overall, rats in the Paired group made many contacts after houselight onset, whereas those in the Unpaired group made few to no contacts [group main effect:  $F(1,15) = 6.23$ ,  $p < 0.025$ ; Figure 3B].

The per-trial frequency of houselight illumination-elicited orienting in session 12 is presented in Supplemental Figure 4.

### Acquisition of similar reactions to sipper presentation across ethanol-reinforced classical conditioning in the paired and unpaired groups


Equipment malfunction resulted in failure to record sipper licking during at least one session for one rat in the Unpaired group, reducing sample size to 6 for these analyses.

**Table 1**  
Blood ethanol concentrations, bodyweight, and drinking between groups on blood sampling day.

	Paired ( $n = 10$ )	Unpaired ( $n = 7$ )	
BEC (mg/dL)	$18.41 \pm 4.70$	$12.46 \pm 5.13$	$T_{15} = 0.839$ , NS
Time after 8th sipper presentation (min)	$9.09 \pm 0.25$	$9.33 \pm 0.33$	$T_{15} = -0.588$ , NS
Bodyweight (g)	$283.3 \pm 5.23$	$282.0 \pm 3.72$	$T_{15} = 0.185$ , NS
Ethanol (g)	$0.219 \pm 0.017$	$0.178 \pm 0.020$	$T_{15} = 1.529$ , NS
Dose (g/kg)	$0.777 \pm 0.0658$	$0.634 \pm 0.072$	$T_{15} = 1.438$ , NS
BEC-Dose Correlation Coefficient	$0.677 (0.08, 0.92)^*$	$0.892 (0.42, 0.98)^*$	$T_{13} = 0.552$ , NS

BEC stands for blood ethanol concentration. For rows 2–6, entries in columns 2–3 are mean  $\pm$  S.E.M. For row 7, entries in columns 2–3 are Pearson's product-moment correlation coefficients with lower and upper 95% confidence limits in parentheses. Asterisks indicate  $p < 0.05$  for the null hypothesis that the true correlation coefficient equals zero. For rows 2–6, entries in column 4 are Student's  $t$  test results for the null hypothesis that the true mean difference between groups equals zero. For row 7, the entry in column 4 represents the Student's  $t$  test result for the null hypothesis that group does not moderate the relationship between BEC and dose.





**Fig. 2. Conditioning of houselight-elicited anticipatory seeking.** Mean  $\pm$  S.E.M. level of sipper site approach (A) and faceplate contacts (B) paneled by trial phase (pre-CS bin: 5-sec bin before houselight onset; CS bins 1–2: 1st and 2nd 5-sec bin of illumination) shown across conditioning sessions (8 trials/session, 1 session/day, 12 consecutive days) for adult, female Long-Evans rats. Black and white triangles represent Paired group ( $n = 10$ ) and Unpaired group ( $n = 7$ ), respectively. Approach data (maximum response level was 4) were derived from offline manual videoscoring (see main text Methods: Behavior Measurement for videoscoring details). Contact data were collected online using a modified lickometer (see main text Methods: Behavior Measurement for details).


There was a decrease across sessions in average latency to start licking per trial [session main effect:  $F(11,154) = 12.58$ ,  $p < 0.001$ ; Figure 4A]. There was a concomitant increase across sessions in average licks per trial [session main effect:  $F(11,154) = 16.57$ ,  $p < 0.001$ ; Figure 4B]. Statistically significant group  $\times$  session interaction effects were also detected [in latency:  $F(11,154) = 2.05$ ,  $p < 0.05$ ; in licks:  $F(11,154) = 3.15$ ,  $p < 0.05$ ], but simple effects decomposition revealed that these were driven by trivial differences between groups early in conditioning (sessions 1, 2, and/or 3) that were not statistically significant after Bonferroni correction (Fig. 4A and B). Importantly, by the end of conditioning, there was no significant difference between the Paired and Unpaired groups in either the average latency to start licking or the average licks per trial (both group main effects, both session main effects, and both group  $\times$  session interactions over sessions 10–12: NS before and after Bonferroni correction; Figure 4A and B).

#### Correlation between cue-elicited ethanol seeking and ethanol drinking behavior in the paired group

For ease of comparison to Cofresí et al. (2018), each Paired group rat's asymptotic level of behavior was estimated as the average across conditioning sessions 10–12. The cue-elicited ethanol-seeking reaction was indexed by the level of approach during trial phase CS2 per trial because that is the within-trial period during which it was at its peak. Indices of ethanol drinking behavior included latency to start licking the sipper per trial, the number of licks per trial, and the total ingested dose of ethanol per session.

Cue-elicited ethanol seeking explained 61% of the total variance in the average latency to lick the sipper, 58% of the variance in the average numbers of licks, and 40% of the variance in the average total ingested dose in the Paired group. Specifically, we found that on average across trials at asymptote, higher levels of cue-elicited ethanol seeking were significantly associated with lower latency to start ethanol sipper licking [ $r = -0.78$ ,  $t_8 = -3.53$ ,  $p < 0.01$ ;


Figure 5A], more licking [ $r = +0.76$ ,  $t_8 = +3.15$ ,  $p < 0.02$ ; Figure 5B], and larger ingested doses [ $r = +0.63$ ,  $t_8 = +2.314$ ,  $p < 0.05$ ; Figure 5C].




**Fig. 3. Within-session dynamics of houselight-elicited anticipatory seeking.** Mean  $\pm$  S.E.M. level of anticipatory sipper site approach (A) and faceplate contacts (B) in the 5 s before light onset (bin -1) and over the 10-sec post-light onset but pre-sipper onset (CS bin 1 and 2, each 5 s) paneled by trial (1–8) within conditioning session 12 for adult, female Long-Evans rats. Black and white triangles represent Paired group ( $n = 10$ ) and Unpaired group ( $n = 7$ ), respectively. Approach data (maximum response level was 4) were derived from offline manual videoscoring.

## Drinking Per Trial Across Conditioning

### A. Average Licking Latency Per Trial



### B. Average Total Licks Per Trial



**Fig. 4. Equivalent drinking behavior across cue conditioning.** Mean  $\pm$  S.E.M. (A) latency (sec) to start licking per trial and (B) number of licks per trial shown across conditioning sessions (8 trials/session, 1 session/day, 12 consecutive days) for adult, female Long-Evans rats. Black and white triangles represent Paired group ( $n = 10$ ) and Unpaired group ( $n = 6$  out of 7 due to equipment malfunction), respectively.

## Discussion


In the present study, we had the following goals: 1) determine whether cue-triggered alcohol-seeking behaviors in female rats resulted from repeated exposure to alcohol, the cue, or associative learning, and 2) test whether the covariation between alcohol-cue reactivity and drinking behavior existed within episodes as predicted by a major theoretical framework for understanding the role of Pavlovian alcohol cues in alcohol use behavior.

### Pre-conditioning free-choice alcohol drinking and preference

Adult female rats drank just as much alcohol at the start as at the end of the 5-week pre-conditioning home cage drinking period in the present study (Supplemental Fig. 1A), which replicates previous findings (Butler, Carter, & Weiner, 2014; Morales, McGinnis, & McCool, 2015). Unlike in those studies, however, our female rats appeared to lose their initial aversion to the taste of unsweetened alcohol (Supplemental Fig. 1B). This could be accounted for by the rats learning to associate the taste of alcohol with its post-ingestive reinforcing effects (metabolic or pharmacological or both). This minor discrepancy between our study and the studies of Morales et al. (2015) and Butler et al. (2014) is most likely attributable to our use of a lower alcohol concentration in the drinking solution (10–15% alcohol v/v in tap water in our study compared to 20% alcohol v/v in their studies).

### Acquisition of alcohol-cue reactivity

After acquisition of voluntary drinking, we tested for cue-alcohol associative learning. The only difference between the two



**Fig. 5. Conditioned cue reactivity predicts drinking latency, drinking intensity, and ingested dose in Paired group.** Relationships of latency to start licking per trial (A), total licks per trial (B), and ingested dose per session (C) to houselight-elicited sipper site approach level per trial (maximum = 4) on average across conditioning sessions 10–12. Data were from 10 adult, female Long-Evans rats. Solid lines in each panel represent the regression line. Dashed lines represent the upper and lower 95% confidence limits around the regression line.

groups (paired and unpaired) was the presence of a positive contingency between houselight illumination (CS) and alcohol access in the Paired group. In both groups, rats learned to react to sipper presentation with rapid initiation of vigorous consummatory licking (Fig. 4A and B), and learned to react to initial oral alcohol receipt with an increase in the rate of consummatory licking (Supplemental Fig. 6A). Rats in both groups ingested similar doses of alcohol across conditioning (Fig. 1A). Similar levels of alcohol were detected in blood approximately 10 min after the 8th drinking opportunity in a conditioning session (Fig. 1B), and similar levels were predicted to be experienced over the course of conditioning (Fig. 1C). Although total ingested doses by these female rats were numerically larger than those ingested by male rats in the same paradigm, blood alcohol levels in the female rats were similar to those of male rats in this (Cofresí et al., 2018, 2019) and similar paradigms (LeCocq et al., 2018). During the 5-min pre-session “waiting” periods, both groups moved around more in the stimulus-rich than stimulus-poor side of the conditioning chamber (Supplemental Fig. 2A) and made a similar number of sipper site (faceplate) contacts (Supplemental Fig. 2B). However, only rats in

the Paired group acquired houselight illumination-elicited anticipatory sipper site approach and contact behavior (Fig. 2A and B). This is strong behavioral evidence that cue-triggered alcohol-seeking behaviors arise from associative learning and are not merely due to repeated exposure to alcohol or the cue within the same context. Additionally, it confirms that associative learning about antecedent conditional stimuli for alcohol access in this (Cofresí et al., 2019) and similar paradigms (Srey et al., 2015) is not restricted to male rats.

Despite equivalent alcohol exposure, rats in the Unpaired group did not develop cue-triggered alcohol-directed behavior. However, we did observe persistence of the overt attentional orienting reaction to houselight illumination – specifically, orienting during the second half of light illumination (Supplemental Figs. 3–4) – in the unpaired female rats. We have also observed the same form of persistent orienting in male rats that went through a similar habituation and conditioning paradigm, with the houselight being explicitly unpaired with alcohol access (Cofresí et al., 2019). Our present findings in female rats suggest that in both sexes, the persistent overt attentional response in the Unpaired group may be a conditioned attentional response (Delamater & Holland, 2008; Holland, 1980) that reflects associative learning about houselight offset as a predictor of alcohol access.

#### *Trial-by-trial dynamics of alcohol-cue reactivity*

In the present study, in conditioning session 12, the Paired group female rats exhibited no within-session trial-by-trial decay in the level of houselight illumination-elicited sipper site approach and contact (Fig. 3A and B). Female rats in the Paired and Unpaired groups alike exhibited no trial-by-trial change in the latency to approach the sipper area upon sipper presentation (Supplemental Fig. 5A), but did exhibit a small trial-by-trial increase in the latency to start drinking (Supplemental Fig. 5B), and a small decrease in the overall intensity of drinking from trials 1–4 to 5–8 (Supplemental Fig. 6B). In contrast, in our previous study, equivalently experienced, paired-group male rats exhibited trial-by-trial decreases in the vigor of both houselight illumination-elicited sipper site approach and drinking behavior, whereas male rats in the unpaired group did not (Cofresí et al., 2018, 2019).

We explained our previous findings by positing that rats may experience progressive within-session specific satiety for alcohol, and consequently, progressive devaluation of the alcohol reinforcer (Samson, Czachowski, & Slawecki, 2000; Samson, Slawecki, Sharpe, & Chappell, 1998). Cue-elicited goal-directed behavior is known to be sensitive to between-session reinforcer devaluation (e.g., specific satiety, pairing with illness) in food and sugar cue conditioning paradigms (Holland & Rescorla, 1975; Morrison, Bamkole, & Nicola, 2015). Based on that literature, we argued that if within-session specific satiety for alcohol and consequent devaluation of the alcohol reinforcer were taking place, then we would expect trial-by-trial decay in the level of houselight cue-elicited alcohol seeking. The present findings suggest that while cue-elicited alcohol seeking may be sensitive to progressive within-session specific satiety for alcohol and consequent devaluation of the alcohol reinforcer in male rats, it may not be similarly sensitive in female rats.

We also previously argued that if the vigor of alcohol drinking behavior had come to be in part controlled by the conditioned alcohol cue, then it too would be sensitive to progressive within-session satiety for alcohol and consequent devaluation of the alcohol reinforcer. If so, then we would expect a trial-by-trial decrease in the vigor of alcohol drinking behavior specifically among the Paired, but not Unpaired, group female rats. Given that both Paired and Unpaired groups in the present study exhibited

trial-by-trial decreases in the vigor of alcohol drinking behavior (Supplemental Fig. 5B & 6B), we cannot argue that the conditioned alcohol cue in Paired-group female rats exerted any direct control over the vigor of their alcohol drinking behavior. However, our finding that both Paired and Unpaired group female rats exhibited trial-by-trial decreases in alcohol drinking behavior agrees with the idea that progressive within-session specific satiety took place.

Thus, it is tempting to interpret the insensitivity of cue-elicited alcohol seeking to within-session specific satiety for alcohol in the female rat as an indication that despite conditioning the alcohol cue at a similar rate and reacting to that cue with what looks like the same response, male rats encoded the alcohol cue in a stimulus-outcome memory, whereas female rats encoded the alcohol cue in a stimulus-response memory. A more parsimonious, and more easily tested, alternative explanation is that male and female rats may simply be differentially sensitive to different types of reinforcer devaluation in general or specifically, different types of devaluation applied to an alcohol reinforcer. Either explanation has implications for the sensitivity of alcohol-cue reactivity to different behavioral interventions between men and women.

#### *Cue-triggered alcohol-directed reactivity promotes alcohol intake*

According to a model for alcohol abuse proposed by Tomie and colleagues (Tomie, 1996; Tomie & Sharma, 2013), alcohol-cue reactivity should co-vary with alcohol drinking. One of our previous studies confirmed this prediction in male rats (Cofresí et al., 2018). In the present study, we extend this finding to female rats. Specifically, we found that greater levels of houselight illumination-elicited alcohol seeking predicted faster initiation of drinking, more drinking, and the ingestion of larger alcohol doses (Fig. 5A–C). These relationships could be due to a causal response chain or between-subject differences in biopsychological factors that influence conditioning rates, reactivity levels, and drinking.

#### *The present study in context*

Our present findings are not surprising given that female rats have been shown to condition behavioral reactions to cues predicting: 1) appetitive stimuli such as food or sugar pellets (Anderson & Petrovich, 2015; Pitchers et al., 2015), 2) aversive stimuli such as mild foot shock (Kosten, Lee, & Kim, 2006; Milad, Igoe, Lebron-Milad, & Novales, 2009; Pryce, Lehmann, & Feldon, 1999), and 3) drugs of abuse such as cocaine (Feltenstein, Henderson, & See, 2011; Kippin et al., 2005), especially in involuntary drug exposure paradigms (Bobzean, Dennis, & Perrotti, 2014; Campbell, Wood, & Spear, 2000; Russo, Festa, et al., 2003; Russo, Jenab, et al., 2003). In fact, there is evidence for appetitive and aversive conditioning to cues predicting involuntary alcohol exposure in female rats (Nentwig, Myers, & Grisel, 2017; Sherrill, Berthold, Koss, Juraska, & Gulley, 2011; Torres, Walker, Beas, & O'Dell, 2014). Additionally, there is indirect evidence for female rats conditioning to appetitive cues for voluntary alcohol consumption from studies of cue-induced reinstatement of extinguished alcohol self-administration behaviors (Bertholomey, Nagarajan, & Torregrossa, 2016; Randall, Stewart, & Besheer, 2017). The main contribution of the present study to our field is as an empirical demonstration that appetitive Pavlovian conditioning to voluntary alcohol consumption progresses similarly in female as well as in male rats. Our unequivocal verification of this basic phenomenon in female rats is important for other pre-clinical laboratories conducting behavioral or neurobiological studies of alcohol-cue reactivity using the rat as a model organism.



## On the role of the estrous cycle

The present study was not designed to evaluate estrous cycle effects on alcohol learning and memory in the freely cycling female rat. In fact, we chose not to monitor the estrous cycle in our study for two reasons. First and foremost, we wanted to minimize procedural differences between the present study and our previous studies in male rats. Second, we were concerned that daily vaginal lavage could be capable of altering the conditioning properties of alcohol because it is an invasive, stressful procedure (Sharp, Zammit, Azar, & Lawson, 2003). Others have applied daily lavage and not observed detrimental effects on home cage alcohol drinking and operant self-administration (Priddy et al., 2017). (However, it should be noted that, to our knowledge, no proper experiment evaluating the potential effects of daily vaginal lavage [as a stressor] on the alcohol intake of female rats could be found in the literature.) Importantly, Priddy et al. (2017) also reported null effects of estrous cycle phase, in agreement with earlier studies in freely cycling female rats (Roberts, Smith, Weiss, Rivier, & Koob, 1998). Another recent study, in which vaginal lavage was done only once after the final operant self-administration session, also failed to find an effect of estrous cycle phase in freely cycling female rats (Bertholomey et al., 2016). Despite these null effects of the estrous cycle on alcohol intake in female rats, a recent meta-analysis across human and non-human animal models indicated that gonadal hormones do exert modulatory effects on alcohol intake (Erol, Ho, Winham, & Karpyak, 2017). Moreover, failure of the estrous cycle to modulate overall voluntary alcohol consumption does not preclude the estrous cycle from modulating cue reactivity phenomena. Indeed, extinction of fear and cocaine cues, and especially, post-extinction relapse-like return of reactivity to those cues, have been shown to be modulated by estrous cycle phase in the female rat (Feltenstein et al., 2011; Kippin et al., 2005; Milad et al., 2009). Additionally, studies of conditioned place preference to involuntary alcohol exposure in female rats (Torres et al., 2014) and female mice (Hilderbrand & Lasek, 2018) alike strongly implicate gonadal hormone variation over the estrous cycle in modulating the appetitive conditioning properties of alcohol. Consequently, future studies should characterize the role of the female rat estrous cycle, if any, in appetitive Pavlovian conditioning to voluntary alcohol consumption, its extinction, and post-extinction relapse-like response return.

## Conclusion

We found that an alcohol access-related cue acquired the ability to elicit an alcohol approach response in female rats only if that cue positively predicted alcohol access. In doing so, we confirmed that associative learning about antecedent conditional stimuli for alcohol access in our paradigm, by extension in similar paradigms, is not restricted to male rats. Within-session patterns of cue-elicited alcohol seeking and drinking by female rats exhibited subtle differences from what we have previously observed in male rats. Overall, our findings underscore the importance of Pavlovian conditioning processes in alcohol self-administration across the sexes as well as the need for increased study of the female sex in preclinical animal models of alcohol-cue reactivity.

## Acknowledgments

Funding was provided by NIH NIAAA R37AA11852 (RAG), NIH NIAAA T32AA007471 (RUC), and CIHR MOP-137030 (NC).

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.alcohol.2019.03.003>.

## References

- Anderson, L. C., & Petrovich, G. D. (2015). Renewal of conditioned responding to food cues in rats: Sex differences and relevance of estradiol. *Physiology & Behavior*, 151, 338–344. <https://doi.org/10.1016/j.physbeh.2015.07.035>.
- Barker, J. M., & Taylor, J. R. (2017). Sex differences in incentive motivation and the relationship to the development and maintenance of alcohol use disorders. *Physiology & Behavior*, 203, 91–99. <https://doi.org/10.1016/j.physbeh.2017.09.027>.
- Becker, J. B., & Koob, G. F. (2016). Sex differences in animal models: Focus on addiction. *Pharmacological Reviews*, 68, 242–263. <https://doi.org/10.1124/pr.115.011163>.
- Bertholomey, M. L., Nagarajan, V., & Torregrossa, M. M. (2016). Sex differences in reinstatement of alcohol seeking in response to cues and yohimbine in rats with and without a history of adolescent corticosterone exposure. *Psychopharmacology*, 233, 2277–2287. <https://doi.org/10.1007/s00213-016-4278-x>.
- Bobzean, S. A., Dennis, T. S., & Perrotti, L. I. (2014). Acute estradiol treatment affects the expression of cocaine-induced conditioned place preference in ovariectomized female rats. *Brain Research Bulletin*, 103, 49–53. <https://doi.org/10.1016/j.brainresbull.2014.02.002>.
- Butler, T. R., Carter, E., & Weiner, J. L. (2014). Adolescent social isolation does not lead to persistent increases in anxiety-like behavior or ethanol intake in female long-evans rats. *Alcoholism: Clinical and Experimental Research*, 38, 2199–2207. <https://doi.org/10.1111/acer.12476>.
- Campbell, J. O., Wood, R. D., & Spear, L. P. (2000). Cocaine and morphine-induced place conditioning in adolescent and adult rats. *Physiology & Behavior*, 68, 487–493.
- Carrillo, J., Howard, E. C., Moten, M., Houck, B. D., Czachowski, C. L., & Gonzales, R. A. (2008). A 3-day exposure to 10% ethanol with 10% sucrose successfully initiates ethanol self-administration. *Alcohol*, 42, 171–178. <https://doi.org/10.1016/j.alcohol.2008.01.005>.
- Cofresí, R. U., Grote, D. J., Le, E. V. T., Monfils, M.-H., Chaudhri, N., Gonzales, R. A., et al. (2019). Alcohol-associated antecedent stimuli elicit alcohol seeking in non-dependent rats and may activate the insula. *Alcohol*, 76, 91–102. <https://doi.org/10.1016/j.alcohol.2018.08.004>.
- Cofresí, R. U., Lee, H. J., Monfils, M.-H., Chaudhri, N., & Gonzales, R. A. (2018). Characterizing conditioned reactivity to sequential alcohol-predictive cues in well-trained rats. *Alcohol*, 69, 41–49. <https://doi.org/10.1016/j.alcohol.2017.11.034>.
- Cofresí, R. U., Lewis, S. M., Chaudhri, N., Lee, H. J., Monfils, M.-H., & Gonzales, R. A. (2017). Postretrieval extinction attenuates alcohol cue reactivity in rats. *Alcoholism: Clinical and Experimental Research*, 41, 608–617. <https://doi.org/10.1111/acer.13323>.
- Delamater, A. R., & Holland, P. C. (2008). The influence of CS-US interval on several different indices of learning in appetitive conditioning. *Journal of Experimental Psychology: Animal Behavior Processes*, 34, 202–222. <https://doi.org/10.1037/0097-7403.34.2.202>.
- Erol, A., Ho, A. M.-C., Winham, S. J., & Karpyak, V. M. (2017). Sex hormones in alcohol consumption: A systematic review of evidence. *Addiction Biology*, 157–169. <https://doi.org/10.1111/adb.12589>.
- Feltenstein, M. W., Henderson, A. R., & See, R. E. (2011). Enhancement of cue-induced reinstatement of cocaine-seeking in rats by yohimbine: Sex differences and the role of the estrous cycle. *Psychopharmacology*, 216, 53–62. <https://doi.org/10.1007/s00213-011-2187-6>.
- Fox, J., & Weisberg, S. (2011). *An [R] companion to applied regression* (2nd ed.). Thousand Oaks, CA: Sage Publications, Inc.
- Hilderbrand, E. R., & Lasek, A. W. (2018). Estradiol enhances ethanol reward in female mice through activation of ER $\alpha$  and ER $\beta$ . *Hormones and Behavior*, 98, 159–164. <https://doi.org/10.1016/j.yhbeh.2018.01.001>.
- Holland, P. C. (1980). Influence of visual conditioned stimulus characteristics on the form of Pavlovian appetitive conditioned responding in rats. *Journal of Experimental Psychology: Animal Behavior Processes*, 6, 81–97.
- Holland, P. C., & Rescorla, R. A. (1975). The effect of two ways of devaluing the unconditioned stimulus after first- and second-order appetitive conditioning. *Journal of Experimental Psychology: Animal Behavior Processes*, 1, 355–363.
- Inkscape Team. (2017). Inkscape: Free, open-source SVG editor. URL: <https://www.inkscape.org/>.
- Kippin, T. E., Fuchs, R. A., Mehta, R. H., Case, J. M., Parker, M. P., Bimonte-Nelson, H. A., et al. (2005). Potentiation of cocaine-primed reinstatement of drug seeking in female rats during estrus. *Psychopharmacology*, 182, 245–252. <https://doi.org/10.1007/s00213-005-0071-y>.
- Knight, C. P., Hauser, S. R., Deehan, G. A., Jr., Toalston, J. E., McBride, W. J., & Rodd, Z. A. (2016). Oral conditioned cues can enhance or inhibit ethanol (EtOH)-Seeking and EtOH-relapse drinking by alcohol-preferring (P) rats. *Alcoholism*:



- Clinical and Experimental Research*, 40, 906–915. <https://doi.org/10.1111/acer.13027>.
- Kosten, T. A., Lee, H. J., & Kim, J. J. (2006). Early life stress impairs fear conditioning in adult male and female rats. *Brain Research*, 1087, 142–150. <https://doi.org/10.1016/j.brainres.2006.03.009>.
- Krank, M. D. (2003). Pavlovian conditioning with ethanol: Sign-tracking (autoshaping), conditioned incentive, and ethanol self-administration. *Alcoholism: Clinical and Experimental Research*, 27, 1592–1598. <https://doi.org/10.1097/01.ALC.0000092060.09228.DE>.
- Krank, M. D., O'Neill, S., Square, K., & Jacob, J. (2008). Goal- and signal-directed incentive: Conditioned approach, seeking, and consumption established with unsweetened alcohol in rats. *Psychopharmacology*, 196, 397–405. <https://doi.org/10.1007/s00213-007-0971-0>.
- Lamb, R. J., Ginsburg, B. C., Greig, A., & Schindler, C. W. (2019). Effects of rat strain and method of inducing ethanol drinking on Pavlovian-Instrumental-Transfer with ethanol-paired conditioned stimuli. *Alcohol*, 79, 47–57. <https://doi.org/10.1016/j.alcohol.2019.01.003>.
- Lamb, R. J., Ginsburg, B. C., & Schindler, C. W. (2016). Effects of an ethanol-paired CS on responding for ethanol and food: Comparisons with a stimulus in a Truly-Random-Control group and to a food-paired CS on responding for food. *Alcohol*, 57, 15–27. <https://doi.org/10.1016/j.alcohol.2016.10.009>.
- LeCocq, M. R., Lahlou, S., Chahine, M., Padillo, L. N., & Chaudhri, N. (2018). Modeling relapse to pavlovian alcohol-seeking in rats using reinstatement and spontaneous recovery paradigms. *Alcoholism: Clinical and Experimental Research*, 42, 1795–1806. <https://doi.org/10.1111/acer.13825>.
- Lee, H. J., Groshek, F., Petrovich, G. D., Cantalini, J. P., Gallagher, M., & Holland, P. C. (2005). Role of amygdalo-nigral circuitry in conditioning of a visual stimulus paired with food. *The Journal of Neuroscience*, 25, 3881–3888. <https://doi.org/10.1523/JNEUROSCI.0416-05.2005>.
- Milad, M. R., Igoe, S. A., Lebron-Milad, K., & Novales, J. E. (2009). Estrous cycle phase and gonadal hormones influence conditioned fear extinction. *Neuroscience*, 164, 887–895. <https://doi.org/10.1016/j.neuroscience.2009.09.011>.
- Millan, E. Z., Reese, R. M., Grossman, C. D., Chaudhri, N., & Janak, P. H. (2015). Nucleus accumbens and posterior amygdala mediate cue-triggered alcohol seeking and suppress behavior during the omission of alcohol-predictive cues. *Neuropsychopharmacology*, 40, 2555–2565. <https://doi.org/10.1038/npp.2015.102>.
- Morales, M., McGinnis, M. M., & McCool, B. A. (2015). Chronic ethanol exposure increases voluntary home cage intake in adult male, but not female, Long-Evans rats. *Pharmacology Biochemistry and Behavior*, 139, 67–76. <https://doi.org/10.1016/j.pbb.2015.10.016>.
- Morrison, S. E., Bamkole, M. A., & Nicola, S. M. (2015). Sign tracking, but not goal tracking, is resistant to outcome devaluation. *Frontiers in Neuroscience*, 9, 1–12. <https://doi.org/10.3389/fnins.2015.00468>.
- Nentwig, T. B., Myers, K. P., & Grisel, J. E. (2017). Initial subjective reward to alcohol in Sprague-Dawley rats. *Alcohol*, 58, 19–22. <https://doi.org/10.1016/j.alcohol.2016.11.005>.
- Pitchers, K. K., Flagel, S. B., O'Donnell, E. G., Solberg Woods, L. C., Sarter, M., & Robinson, T. E. (2015). Individual variation in the propensity to attribute incentive salience to a food cue: Influence of sex. *Behavioural Brain Research*, 278, 462–469. <https://doi.org/10.1016/j.bbr.2014.10.036>.
- Priddy, B. M., Carmack, S. A., Thomas, L. C., Vendruscolo, J. C. M., Koob, G. F., & Vendruscolo, L. F. (2017). Sex, strain, and estrous cycle influences on alcohol drinking in rats. *Pharmacology Biochemistry and Behavior*, 152, 61–67. <https://doi.org/10.1016/j.pbb.2016.08.001>.
- Pryce, C. R., Lehmann, J., & Feldon, J. (1999). Effect of sex on fear conditioning is similar for context and discrete CS in Wistar, Lewis and Fischer rat strains. *Pharmacology Biochemistry and Behavior*, 64, 753–759.
- R Core Team. (2018). *R: A Language and Environment for Statistical Computing*. Vienna, Austria.
- Randall, P. A., Stewart, R. T., & Besheer, J. (2017). Sex differences in alcohol self-administration and relapse-like behavior in Long-Evans rats. *Pharmacology Biochemistry and Behavior*, 156, 1–9. <https://doi.org/10.1016/j.pbb.2017.03.005>.
- Roberts, A. J., Smith, A. D., Weiss, F., Rivier, C., & Koob, G. F. (1998). Estrous cycle effects on operant responding for ethanol in female rats. *Alcoholism: Clinical and Experimental Research*, 22, 1564–1569.
- Russo, S. J., Festa, E. D., Fabian, S. J., Gazi, F. M., Kraish, M., Jenab, S., et al. (2003). Gonadal hormones differentially modulate cocaine-induced conditioned place preference in male and female rats. *Neuroscience*, 120, 523–533.
- Russo, S. J., Jenab, S., Fabian, S. J., Festa, E. D., Kemen, L. M., & Quinones-Jenab, V. (2003). Sex differences in the conditioned rewarding effects of cocaine. *Brain Research*, 970, 214–220.
- Samson, H. H., Czachowski, C. L., & Slawecki, C. J. (2000). A new assessment of the ability of oral ethanol to function as a reinforcing stimulus. *Alcoholism: Clinical and Experimental Research*, 24, 766–773.
- Samson, H. H., Slawecki, C. J., Sharpe, A. L., & Chappell, A. (1998). Appetitive and consummatory behaviors in the control of ethanol consumption: A measure of ethanol seeking behavior. *Alcoholism: Clinical and Experimental Research*, 22, 1783–1787.
- Sharp, J., Zammit, T., Azar, T., & Lawson, D. (2003). Stress-like responses to common procedures in individually and group-housed female rats. *Contemporary Topics in Laboratory Animal Science*, 42, 9–18.
- Sherrill, L. K., Berthold, C., Koss, W. A., Juraska, J. M., & Gulley, J. M. (2011). Sex differences in the effects of ethanol pre-exposure during adolescence on ethanol-induced conditioned taste aversion in adult rats. *Behavioural Brain Research*, 225, 104–109. <https://doi.org/10.1016/j.bbr.2011.07.003>.
- Sparks, L. M., Sciascia, J. M., Ayorech, Z., & Chaudhri, N. (2014). Vendor differences in alcohol consumption and the contribution of dopamine receptors to Pavlovian-conditioned alcohol-seeking in Long-Evans rats. *Psychopharmacology*, 231, 753–764. <https://doi.org/10.1007/s00213-013-3292-5>.
- Srey, C. S., Maddux, J.-M. N., & Chaudhri, N. (2015). The attribution of incentive salience to pavlovian alcohol cues: A shift from goal-tracking to sign-tracking. *Frontiers in Behavioral Neuroscience*, 9, 54. <https://doi.org/10.3389/fnbeh.2015.00054>.
- Tomie, A. (1996). Locating reward cue at response manipulandum (CAM) induces symptoms of drug abuse. *Neuroscience & Biobehavioral Reviews*, 20, 505–535.
- Tomie, A., Festa, E. D., Sparta, D. R., & Pohorecky, L. A. (2003). Lever conditioned stimulus-directed autoshaping induced by saccharin-ethanol unconditioned stimulus solution: Effects of ethanol concentration and trial spacing. *Alcohol*, 30, 35–44.
- Tomie, A., Kuo, T., Apor, K. R., Salomon, K. E., & Pohorecky, L. A. (2004). Autoshaping induces ethanol drinking in nondeprived rats: Evidence of long-term retention but no induction of ethanol preference. *Pharmacology Biochemistry and Behavior*, 77, 797–804. <https://doi.org/10.1016/j.pbb.2004.02.005>.
- Tomie, A., Miller, W. C., Dranoff, E., & Pohorecky, L. A. (2006). Intermittent presentations of ethanol sipper tube induce ethanol drinking in rats. *Alcohol and Alcoholism*, 41, 225–230. <https://doi.org/10.1093/alcalc/agl002>.
- Tomie, A., & Sharma, N. (2013). Pavlovian sign-tracking model of alcohol abuse. *Current Drug Abuse Reviews*, 6, 201–219.
- Torres, O. V., Walker, E. M., Beas, B. S., & O'Dell, L. E. (2014). Female rats display enhanced rewarding effects of ethanol that are hormone dependent. *Alcoholism: Clinical and Experimental Research*, 38, 108–115. <https://doi.org/10.1111/acer.12213>.
- Villaruel, F. R., & Chaudhri, N. (2016). Individual differences in the attribution of incentive salience to a pavlovian alcohol cue. *Frontiers in Behavioral Neuroscience*, 10, 238. <https://doi.org/10.3389/fnbeh.2016.00238>.
- Wickham, H. (2009). *ggplot2: elegant graphics for data analysis*. New York, New York: Springer-Verlag.