

Very Brief Notes

Professor David Barrett

Transcribed by Thomas Cohn

9/28/18

Defn: $f \in C^\infty \Leftrightarrow f \in C^r \forall r \in \mathbb{N} \cup \{0\}$.

$$f \in C^1(A^{\text{osso}V}, W)$$

$$f \in C^2 \Leftrightarrow Df \in C^1$$

$$f \in C^r \Leftrightarrow Df \in C^{r-1} \text{ (for } r = 2, 3, 4, \dots)$$

$$f \in C^r \Leftrightarrow D_{\vec{u}_1} D_{\vec{u}_2} \cdots D_{\vec{u}_k} f \text{ exists and is continuous } \forall \vec{u}_1, \dots, \vec{u}_k \in V \text{ with } k \leq r.$$

From Wednesday, $f \in C^2(A^{\text{osso}V}, \mathbb{R}^n) \rightarrow D_{\vec{u}_1} D_{\vec{u}_2} f = D_{\vec{u}_2} D_{\vec{u}_1} f$.

This also works for $f \in C^2(A, W)$ with $\dim W < +\infty$ (see study exercise 7).

Suppose $f \in C^r$. Consider $D_{\vec{u}_1} D_{\vec{u}_2} \cdots D_{\vec{u}_r} f$. We can interchange \vec{u}_j with \vec{u}_{j+1} , so we can arbitrarily permute \vec{u}_j .

Ex: Suppose $f \in C^r(\mathbb{R}^n, \mathbb{R})$. How many distinct $D_{j_1} \cdots D_{j_r} f$ are there?