

# Math 591 Lecture 33

Professor Alejandro Uribe-Ahumada

*Transcribed by Thomas Cohn*

11/18/20

## Global formulae for $d$

First, we want to consider the pairing of forms and vector fields. If  $\alpha \in \Omega^k$  and  $X_1, \dots, X_k \in \mathfrak{X}(M)$ , then  $\alpha(X_1, \dots, X_k) : M \rightarrow \mathbb{R}$  is a function on  $M$ .

**Prop:** (1) If  $\alpha \in \Omega^1(M)$ ,  $Y_0, Y_1 \in \mathfrak{X}(M)$ , then

$$(d\alpha)(Y_0, Y_1) = Y_0(\alpha(Y_1)) - Y_1(\alpha(Y_0)) - \alpha([Y_0, Y_1])$$

Proof: WOLOG assume  $\alpha = f dg$ . Then  $d\alpha = df \wedge dg$ . Thus,

$$(d\alpha)(Y_0, Y_1) = df(Y_0)dg(Y_1) - df(Y_1)dg(Y_0) = Y_0(f)Y_1(g) - Y_1(f)Y_0(g)$$

Well, the RHS of this is

$$Y_0(fY_1(g)) - Y_1(fY_0(g)) - f[Y_0, Y_1]g = (Y_0f)(Y_1g) + \cancel{fY_0Y_1g} - (Y_1fY_0g + \cancel{fY_1Y_0g}) - f(Y_0Y_1g - Y_1Y_0g)$$

□

**Prop:** (2) If  $\alpha \in \Omega^2(M)$ ,  $Y_0, Y_1, Y_2 \in \mathfrak{X}(M)$ , then

$$(d\alpha)(Y_0, Y_1, Y_2) = Y_0\alpha(Y_1, Y_2) - Y_1\alpha(Y_0, Y_2) + Y_2\alpha(Y_0, Y_1) - \alpha([Y_0, Y_1], Y_2) + \alpha([Y_0, Y_2], Y_1) - \alpha([Y_1, Y_2], Y_0)$$

In general, if  $\alpha \in \Omega^k(M)$ ,  $Y_0, \dots, Y_k \in \mathfrak{X}(M)$ , then

$$(d\alpha)(Y_0, \dots, Y_k) = \sum_{j=0}^k (-1)^j Y_j \alpha(Y_0, \dots, \widehat{Y_j}, \dots, Y_k) + \sum_{0 \leq i < j \leq k} (-1)^{i+j} \alpha([Y_i, Y_j], Y_0, \dots, \widehat{Y_i}, \dots, \widehat{Y_j}, \dots, Y_k)$$

where  $\widehat{\phantom{x}}$  indicates that we omit that term.

Refer to Proposition 20.14 in Tu/Proposition 14.32 in Lee.

Another way to think of this is, given  $\alpha \in \Omega^k(M)$ ,  $X_1, \dots, X_k \in \mathfrak{X}(M)$ , and  $p \in M$ , we have  $(X_i)_p \in T_p M$ , so

$$\alpha_p : \underbrace{T_p M \times \dots \times T_p M}_k = (T_p M)^k \rightarrow \mathbb{R} \\ p \mapsto \alpha_p((X_1)_p, \dots, (X_k)_p) \in \mathbb{R}$$

## Lie Derivatives of Forms

**Defn:** Let  $X \in \mathfrak{X}(M)$ ,  $\alpha \in \Omega^k(M)$ , and  $p \in M$ . Let  $\varphi$  be the local flow of  $X$ . Form the curve  $(-\varepsilon, \varepsilon) \ni t \mapsto (\varphi_t^* \alpha)_p \in \bigwedge^k(T_p^* M)$ . (Note that  $\bigwedge^k(T_p^* M)$  is independent of  $t$ .) Then we defined

$$(\mathcal{L}_X \alpha)_p = \left. \frac{d}{dt} (\varphi_t^* \alpha)_p \right|_{t=0}$$

Observe: If  $k = 0$ , then  $\alpha$  is a function on  $M$ , so

$$(\mathcal{L}_X \alpha)_p = \left. \frac{d}{dt} (\alpha \circ \varphi_t)_p \right|_{t=0} = \left. \frac{d}{dt} (\alpha(\varphi_t(p))) \right|_{t=0} = \dot{\varphi}_t \Big|_{t=0} (\alpha)$$

Thus,  $(\mathcal{L}_X \alpha)_p = X(\alpha)(p)$  if  $\alpha$  is a function.

## First Properties

**Prop:**

- a) The expression  $\mathcal{L}_X(\alpha)$  is  $\mathbb{R}$ -linear w.r.t  $X$  and  $\alpha$ .
- b)  $\mathcal{L}_X(\alpha \wedge \beta) = \mathcal{L}_X(\alpha) \wedge \beta + \alpha \wedge \mathcal{L}_X(\beta) - \mathcal{L}_X$  is a derivation.
- c)  $\mathcal{L}_X(d\alpha) = d\mathcal{L}_X(\alpha)$ .
- d) If  $F : M \rightarrow M$  is a diffeomorphism, then  $\mathcal{L}_X(F^*\alpha) = F^*(\mathcal{L}_{F_*(X)}\alpha)$ .

Proof: (b) holds because  $\varphi_t^*(\alpha \wedge \beta) = (\varphi_t^*\alpha) \wedge (\varphi_t^*\beta)$ . We just need to show the right hand side satisfies the usual product rule with respect to  $\frac{d}{dt}$ .

(c) holds because  $\varphi_t^*(d\alpha) = d(\varphi_t^*\alpha)$ ,  $\forall t$ . This uses the fact that  $d$  and  $\frac{d}{dt}$  commute, i.e.,  $\frac{\partial^2}{\partial t \partial x^j} = \frac{\partial^2}{\partial x^j \partial t}$ .

(d) holds because for  $p \in M$ ,  $v_1, \dots, v_k \in T_p M$ ,

$$\mathcal{L}_X(F^*\alpha)_p(v_1, \dots, v_k) = \frac{d}{dt} \underbrace{\varphi_t^*(F^*\alpha)_p(v_1, \dots, v_k)}_{(\varphi_t^* \circ F^*)(\alpha) = (F \circ \varphi_t)^*(\alpha)} \Big|_{t=0} = \frac{d}{dt} \alpha((F \circ \varphi_t)_{*,p}(v_1), \dots, (F \circ \varphi_t)_{*,p}(v_k)) \Big|_{t=0}$$

Observe:  $t \mapsto F \circ \varphi_t$  is the integral curve of  $F_*(X)$  at  $F(p)$ . So

$$\mathcal{L}_X(F^*\alpha)_p(v_1, \dots, v_k) = \dots = (\mathcal{L}_{F_*(X)}\alpha)(F_{*,p}(v_1), \dots, F_{*,p}(v_k)) = F^*(\mathcal{L}_{F_*(X)}\alpha)(v_1, \dots, v_k)$$

□

**Cor:** If  $\mathcal{L}_X\alpha = 0$  (everywhere), then  $\varphi_t^*\alpha = \alpha$ ,  $\forall t$ .

Proof: We will show  $\forall p \in M$ , the curve  $t \mapsto (\varphi_t^*\alpha)_p \in \bigwedge^k T_p^*M$  is constant.

Proof:  $\forall s$ ,

$$\frac{d}{dt} \varphi_t^*\alpha \Big|_{t=s} = \frac{d}{dt} (\varphi_{t+s}^*\alpha) \Big|_{t=0} = \frac{d}{dt} \varphi_t^*(\varphi_s^*\alpha) \Big|_{t=0} = \mathcal{L}_X(\varphi_s^*\alpha) = \varphi_s^* \underbrace{\mathcal{L}_{(\varphi_s)_*(X)}\alpha}_{=X} = \varphi_s^* \underbrace{\mathcal{L}_X\alpha}_{=0} = 0$$

□

## Towards Cartan's Formula

We need interior multiplication/contraction.

**Defn:** Given  $\alpha \in \Omega^k$ ,  $X \in \mathfrak{X}(M)$ , we define  $\iota_X\alpha \in \Omega^{k-1}$  by,  $\forall p \in M$  and  $v_1, \dots, v_{k-1} \in T_p M$ ,

$$(\iota_X\alpha)_p(v_1, \dots, v_{k-1}) = \alpha_p(X_p, v_1, \dots, v_{k-1})$$

**Thm:** (Cartan's Magic Formula)  $\mathcal{L}_X = d \circ \iota_X + \iota_X \circ d : \Omega^k \rightarrow \Omega^k$ .

Picture:

$$\begin{array}{ccccccc} \dots & \xrightarrow{d} & \Omega^{k-1} & \xrightarrow{d} & \Omega^k & \xrightarrow{d} & \Omega^{k+1} \xrightarrow{d} \dots \\ & & & \swarrow \iota_X & \Downarrow \mathcal{L}_X & \searrow \iota_X & \\ \dots & \xrightarrow{d} & \Omega^{k-1} & \xrightarrow{d} & \Omega^k & \xrightarrow{d} & \Omega^{k+1} \xrightarrow{d} \dots \end{array}$$

where  $\mathcal{L}_X$  is the “sum of two paths”.

On Friday, we'll show that when  $\alpha = dx \wedge dy$  and  $M = \mathbb{R}^2$ , then  $\mathcal{L}_X(\alpha) = (\text{div } X)\alpha$ , in the calc III sense.