

Math 591 Lecture 26

Thomas Cohn

11/2/20

Lie Derivatives

The general notion of a Lie derivative with respect to a vector field X is “pull back by θ_t , the time t map of the flow of X , and then differentiate with respect to t at $t = 0$ ”. We call this \mathcal{L}_X .

Ex: We can pull back functions, so we can have Lie derivatives of functions.

Given X , θ_t , and $f \in C^\infty(M)$, $\phi_t^*(f)(p) = f(\phi_t(p))$ (pullback). Now differentiate:

$$\left. \frac{d}{dt} \phi_t^*(f)(p) \right|_{t=0} = \left. \frac{d}{dt} f(\phi_t(p)) \right|_{t=0} = X(f)(p)$$

So $(\mathcal{L}_X f)(p) = X(f)(p) = df(X_p)$. So $\mathcal{L}_X f$ is again a function on M .

Recall: A 1-form α is an assignment to each $p \in M$, an element $\alpha_p \in T_p^*M$. I.e. α is a section of the cotangent bundle. In local coordinates, we can write $\alpha = \sum \alpha_i dx^i$, for $\alpha_i \in C^\infty(U)$.

Ex: 1-forms can be pulled back by any F , by $F^*(\alpha)(v) = \alpha(F_*, \cdot(v))$. So

$$(\mathcal{L}_X \alpha)(p) = \left. \frac{d}{dt} \underbrace{\phi_t^* \alpha_{\phi_t(p)}}_{\substack{\in T_p^*M \\ \text{for each } t}} \right|_{t=0}$$

Thus, $\mathcal{L}_X \alpha$ is again a 1-form on M .

Special case: $\alpha = df$ for some $f \in C^\infty(M)$. Then by the chain rule (1)

$$F^*(df)(v) = df(F_*, \cdot(v)) \stackrel{(1)}{=} d(f \circ F)(v) = d(F^*f)(v)$$

so $F^*(df) = d(F^*f)$.

So, in general,

$$\mathcal{L}_X(df) = \left. \frac{d}{dt} \phi_t^*(df) \right|_{t=0} = \left. \frac{d}{dt} d\phi_t^*(f) \right|_{t=0} \stackrel{(1)}{=} d \left(\left. \frac{d}{dt} \phi_t^* f \right|_{t=0} \right) = d(df(X)) = d(X(f))$$

Where (1) is true because $\frac{d^2}{\partial t \partial x^i} = \frac{\partial^2}{\partial^i \partial t}$, so the differentiations commute.

In fact, in general, $\mathcal{L}_X(\alpha) = d(\alpha(X)) + \dots$ (we'll fill in the other term later).

Now, back to vector fields...

Note: Vector fields can be pulled back by diffeomorphisms. This means if $X, Y \in \mathfrak{X}(M)$, $\mathcal{L}_X Y$ is defined by, $\forall p \in M$,

$$\mathcal{L}_X(Y)(p) = \left. \underbrace{\frac{d}{dt} [(\phi_t)_*, p]^{-1} (Y_{\phi_t(p)})}_{=V_t \in T_p M} \right|_{t=0}$$

Prop: $\mathcal{L}_X Y = [X, Y]$.

Proof: Compute V_t in coordinates $(U, (x^1, \dots, x^n))$. Restrict to $V \subseteq U$ such that $\phi_t(V) \subseteq U$. Write $\phi_t = (\phi_t^1, \dots, \phi_t^n)$. Then

$$J_t = \left(\frac{\partial \phi_t^i}{\partial x^j}(p) \right)_{(i,j)} \quad Y = \sum_{i=1}^n g_i \frac{\partial}{\partial x^i} \quad X = \sum_{i=1}^n f_i \frac{\partial}{\partial x^i}$$

For a fixed p , let

$$\tilde{Y}_t = Y_{\phi_t(p)} = \begin{pmatrix} g_1(\phi_t(p)) \\ \vdots \\ g_n(\phi_t(p)) \end{pmatrix}$$

Then, let \tilde{V}_t be the column vector with components of V_t . We have

$$\tilde{V}_t = J_t^{-1} \tilde{Y}_t$$

Differentiating with $\frac{d}{dt}$ on both sides, we get

$$\dot{\tilde{V}}_t = J_t^{-1} \dot{\tilde{Y}}_t - J_t^{-1} \dot{J}_t J_t^{-1} \tilde{Y}_t$$

which at $t = 0$, yields

$$\underbrace{\dot{\tilde{V}}_t \Big|_{t=0}}_{\substack{\text{components} \\ \text{of } \mathcal{L}_X Y}} = \begin{pmatrix} X(g_1)(p) \\ \vdots \\ X(g_n)(p) \end{pmatrix} - ?$$

Well,

$$\dot{J}_t = \left(\frac{\partial^2 \phi_t^i}{\partial t \partial x^j}(p) \right)_{(i,j)} = \left(\frac{\partial^2 \phi_t^i}{\partial x^j \partial t}(p) \right)_{(i,j)} = \left(\frac{\partial f_i}{\partial x^j}(p) \right)_{(i,j)}$$

since $\frac{\partial \phi_t^i}{\partial t}(p) = f_i$. Thus, we have

$$\dot{\tilde{V}}_t \Big|_{t=0} = \begin{pmatrix} X(g_1)(p) \\ \vdots \\ X(g_n)(p) \end{pmatrix} - \left(\frac{\partial f_i}{\partial x^j}(p) \right)_{(i,j)} \begin{pmatrix} g_1(p) \\ \vdots \\ g_n(p) \end{pmatrix}$$

Therefore,

$$\begin{aligned} \mathcal{L}_X Y &= \sum_{i=1}^n X(g_i) \frac{\partial}{\partial x^i} - \sum_{i,j=1}^n \frac{\partial f_i}{\partial x^j} g_j \frac{\partial}{\partial x^i} \\ &= \sum_{i=1}^n X(g_i) \frac{\partial}{\partial x^i} - \sum_{i=1}^n Y(f_i) \frac{\partial}{\partial x^i} \\ &= \sum_{i=1}^n (X(g_i) - Y(f_i)) \frac{\partial}{\partial x^i} \\ &= [X, Y] \end{aligned}$$

□

Thm: Using the same notation as last time, $\forall t \in (-\varepsilon, \varepsilon)$, $V_t = \frac{d}{dt}(\phi_{t,*})^{-1}(Y_{\phi_t}(p)) = (\phi_{t,*})^{-1}[X, Y]_{\phi_t(p)}$.

Proof: We already have it for $t = 0$. Now, consider V_{t+s} , and use the group law/translation lemma. □

Cor: If $[X, Y] = 0$, then the flows of X and Y commute.