

Math 635 Lecture 7

Thomas Cohn

2/3/21

Review: Connections on Vector Bundles

Given $\mathcal{E} \rightarrow M$ a vector bundle, a connection is an operator

$$\begin{aligned} \nabla : \mathfrak{X}(M) \times \Gamma(\mathcal{E}) &\rightarrow \Gamma(\mathcal{E}) \\ (X, s) &\mapsto \nabla_X s \end{aligned}$$

with universal quantifiers $\forall X, Y \in \mathfrak{X}(M)$, $s, t \in \Gamma(\mathcal{E})$, and $f \in C^\infty(M)$:

- (1) $\nabla_{X+Y}s = \nabla_X s + \nabla_Y s$
- (2) $\nabla_{fX}s = f\nabla_X s$
- (3) $\nabla_X(s+t) = \nabla_X s + \nabla_X t$
- (4) $\nabla_X(fs) = f\nabla_X s + X(f)s$

Properties (1) and (2) together are written as ∇ is linear in X over $C^\infty(M)$. Note that property (4) implies that $\forall c \in \mathbb{R}$, $\nabla_X(cs) = c\nabla_X s$.

Last time, we saw that $(\nabla_X s)(p)$ only depends on $X_p \in T_p M$ and $s|_U$ for any (arbitrarily small) neighborhood U of p . In other words, if we fix a section $s \in \Gamma(\mathcal{E})$, $\nabla_X s$ is a 1-form that takes values in sections. And if we fix $X \in \mathfrak{X}(M)$, then $\nabla_X \cdot : \Gamma(\mathcal{E}) \rightarrow \Gamma(\mathcal{E})$ is a derivation.

Ex: Say $\mathcal{E} = M \times \mathbb{R}^r \rightarrow M$. (This is locally the general case.) Then $\Gamma(M \times \mathbb{R}^r) \ni s \leftrightarrow \vec{f} : M \rightarrow \mathbb{R}^r$ by $\forall p \in M$, $s(p) = (p, \vec{f}(p))$. This is the same as having a global moving frame $\Gamma(M \times \mathbb{R}^r) \ni E_i \leftrightarrow \vec{f}_i(p) = (0, \dots, 1, \dots, 0)$ (with a 1 in the i th entry), for $i = 1, \dots, r$, because $s = \sum_{i=1}^r f^i E_i$ for $f^i \in C^\infty(M, \mathbb{R})$, where $\vec{f} = (f^1, \dots, f^r)$.

Suppose we have a connection ∇ . Last time, we defined $\vartheta = (\theta_i^j)$, the connection matrix associated with (E_1, \dots, E_r) , by $\forall X, \forall j$,

$$\nabla_X E_j = \sum_{i=1}^r \theta_i^j(X) E_i$$

$\forall i, j, \theta_i^j \in \Omega^1(M)$ (a C^∞ one-form). Then:

$$\nabla_X s = \sum_{i=1}^r \nabla_X (f^i E_i) = \sum_{i=1}^r X(f^i) E_i + \sum_{i=1}^r f^i \nabla_X E_i = \sum_{i=1}^r X(f^i) E_i + \sum_{i,j=1}^r f^i \theta_i^j(X) E_j$$

The corresponding vector-valued function on M is

$$\nabla_X \vec{f} = \left(X(f^1) + \sum_{i=1}^r \theta_i^1(X) f^i, \dots, X(f^r) + \sum_{i=1}^r \theta_i^r(X) f^i \right) = (X(f^1), \dots, X(f^r)) + \left(\sum_{i=1}^r \theta_i^1(X) f^i, \dots, \sum_{i=1}^r \theta_i^r(X) f^i \right)$$

So in vector/matrix notation, with \vec{f} as a column vector, we write

$$\nabla_X \vec{f} = d\vec{f}(X) + \vartheta(X) \vec{f}$$

where $\vartheta(X) = (\theta_i^j(X))$ with lower index i being the columns, and upper index j being the rows.

Conversely, here, ϑ can be any $r \times r$ matrix of one-forms, and this can be used to define a connection on the trivial bundle!

Observe: The previous calculation is valid locally, given some moving frame (E_1, \dots, E_r) of $\mathcal{E} \rightarrow M$ on $U \subseteq M$. Suppose (F_1, \dots, F_r) is another moving frame on U . Then $\forall i, F_i = \sum_j a_i^j E_j$, where the matrix $A = (a_i^j)$ is invertible at each $p \in U$,

and $\forall i, j, a_i^j \in C^\infty(U)$. Given ∇ , we get ϑ , the connection matrix corresponding to the E_j 's, and $\tilde{\vartheta}$, the connection matrix corresponding to the F_j 's.

Exer: Check that $\tilde{\vartheta} = A^{-1}dA + A^{-1}\vartheta A$.

A special feature of the case where $\mathcal{E} = TM$ ($r = n = \dim M$). Consider again a moving frame (E_1, \dots, E_n) . In this case, we can write $X = \sum_k a^k E_k$. In the generic case,

$$\nabla_X E_i = \sum_j \theta_i^j(X) E_j = \sum_{j,k} a^k \underbrace{\theta_i^j(E_k)}_{\text{Christoffel Symbols}} E_j$$

Defn: $\forall i, j, k, \Gamma_{ki}^j \stackrel{\text{def}}{=} \theta_i^j(E_k) \in C^\infty(U)$ are the Christoffel symbols.

Note that $\forall i, k$, we get $\nabla_{E_k} E_i = \sum_j \Gamma_{ki}^j E_j$. So Γ_{ki}^j determines ϑ , and therefore ∇ on U .

Now, back to the general case: vector bundle $\mathcal{E} \rightarrow M$ with connection ∇ . We want to look at “parallelism” and “parallel transport”.

Defn: Let $\gamma : [a, b] \rightarrow M, s \in \Gamma(\mathcal{E})$. Then s is said to be covariant constant, or parallel, iff $\forall t \in [a, b], (\nabla_{\dot{\gamma}(t)} s)(\gamma(t)) = 0$.

Let's analyze this equation... It will turn out to be a system of ordinary differential equations!

Let (E_1, \dots, E_r) be a moving frame on U , a neighborhood of $\gamma(t)$. For the time being, just assume $\text{Im}(\gamma) \subseteq U$. We can write $s = \sum_i f^i E_i$, for some $f^i \in C^\infty(U)$. Let ϑ be the connection matrix w.r.t. the E_i 's.

Introduce $\vec{f} = (f^1, \dots, f^r) : U \rightarrow \mathbb{R}^r$. We saw that $\nabla_{\dot{\gamma}} \vec{f} = d\vec{f}(\dot{\gamma}) + \vartheta(\dot{\gamma})\vec{f}$. Define $f^i(t) = f^i(\gamma(t))$, so here, $\vec{f}(t) = \vec{f}(\gamma(t))$. Then

$$d\vec{f}(\dot{\gamma}) = \frac{d}{dt} \vec{f} \quad \text{and} \quad \frac{d}{dt} \vec{f} + \vartheta(\dot{\gamma})\vec{f} = \vec{0}$$

(taking \vec{f} to be the column vector of the $f^i(t)$'s, and $\vartheta(\dot{\gamma})$ being a t -dependent matrix).