

Seminar 1: Counting Techniques

Carlos Buitrago

Table of contents

1	Basics of Combinatorics	1
2	Fundamental Counting Principles	2
2.1	Sum Rule	2
2.1.1	Example	2
2.2	Product Rule	2
2.2.1	Example	2
2.3	Pigeonhole Principle	2
2.3.1	Example	2
2.4	Counting Formulas	3
2.5	Binomial Theorem	3
3	Fundamental Binomial Identities	3
3.0.1	(1) Symmetry Identity	3
3.0.2	(2) Pascal Identity	3
3.0.3	(3) Combinations with Repetition	3
3.0.4	(4) Sum of Binomial Coefficients	3
3.0.5	(5) Sum of Squares	3
3.0.6	(6) Hockey-Stick Identity	4
3.1	Useful Consequences	4
4	Inclusion–Exclusion Principle	4

1 Basics of Combinatorics

Seminar 1

Probability Theory

by Carlos Buitrago · Fall 2026

2 Fundamental Counting Principles

2.1 Sum Rule

Suppose that a task can be performed in n_1 ways or in n_2 ways, and that the two possibilities are mutually exclusive. Then the total number of possible outcomes equals $n_1 + n_2$.

2.1.1 Example

A class consists of 15 boys and 12 girls. If we wish to choose a class representative, then the number of possible choices is

$$15 + 12 = 27.$$

2.2 Product Rule

Suppose that a procedure consists of k consecutive steps. If Step i can be performed in n_i ways for every $i = 1, \dots, k$, then the total number of possible outcomes equals

$$n_1 n_2 \cdots n_k.$$

2.2.1 Example

A password consists of three letters followed by two digits. Since each letter can be chosen in 26 ways and each digit in 10 ways, the total number of possible passwords equals

$$26^3 \cdot 10^2$$

2.3 Pigeonhole Principle

If $n + 1$ objects are placed into n boxes, then at least one box must contain at least two objects.

2.3.1 Example

Among 13 people there are always at least two born in the same month. Indeed, the objects are the people and the boxes are the months of the year. Since $13 > 12$, the conclusion follows immediately.

2.4 Counting Formulas

	With Repetition	Without Repetition
Combinations	$\binom{n+k-1}{k}$	$\binom{n}{k}$
Arrangements	n^k	$\frac{n!}{(n-k)!}$

2.5 Binomial Theorem

For every $n \in \mathbb{Z}_+$,

$$(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}.$$

3 Fundamental Binomial Identities

3.0.1 (1) Symmetry Identity

$$\binom{n}{k} = \binom{n}{n-k}.$$

3.0.2 (2) Pascal Identity

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}.$$

3.0.3 (3) Combinations with Repetition

$$\binom{n+k-1}{k}.$$

3.0.4 (4) Sum of Binomial Coefficients

$$\sum_{k=0}^n \binom{n}{k} = 2^n.$$

3.0.5 (5) Sum of Squares

$$\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n}.$$

3.0.6 (6) Hockey-Stick Identity

$$\binom{n+m}{n} = \sum_{i=0}^m \binom{n+i-1}{n-1}.$$

3.1 Useful Consequences

From the Hockey-Stick Identity we obtain

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

and

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}.$$

4 Inclusion–Exclusion Principle

Suppose that $a_1, \dots, a_N$ are objects and $d_1, \dots, d_n$ are properties.

Let $N(d_i)$ denote the number of objects satisfying property d_i , and let $N(d_{i_1}, \dots, d_{i_k})$ denote the number of objects satisfying all properties $d_{i_1}, \dots, d_{i_k}$ simultaneously.

Then the number of objects satisfying none of the properties is given by

$$N - \sum_i N(d_i) + \sum_{i < j} N(d_i, d_j) - \sum_{i < j < k} N(d_i, d_j, d_k) + \dots + (-1)^n N(d_1, \dots, d_n).$$