

# Seminar 1: Basics of Combinatorics

Carlos Buitrago

## Table of contents

|          |                                                 |          |
|----------|-------------------------------------------------|----------|
| <b>1</b> | <b>Fundamental Counting Principles</b>          | <b>1</b> |
| 1.1      | Sum Rule . . . . .                              | 1        |
| 1.1.1    | Example . . . . .                               | 2        |
| 1.2      | Product Rule . . . . .                          | 2        |
| 1.2.1    | Example . . . . .                               | 2        |
| 1.3      | Pigeonhole Principle . . . . .                  | 2        |
| 1.3.1    | Example . . . . .                               | 2        |
| 1.4      | Counting Formulas . . . . .                     | 2        |
| 1.5      | Binomial Theorem . . . . .                      | 3        |
| <b>2</b> | <b>Fundamental Binomial Identities</b>          | <b>3</b> |
| 2.0.1    | Symmetry Identity . . . . .                     | 3        |
| 2.0.2    | Pascal Identity . . . . .                       | 3        |
| 2.0.3    | Combinations with Repetition . . . . .          | 3        |
| 2.0.4    | Sum of Binomial Coefficients . . . . .          | 3        |
| 2.0.5    | Sum of Squares . . . . .                        | 3        |
| 2.0.6    | Hockey-Stick Identity . . . . .                 | 4        |
| 2.1      | Useful Consequences . . . . .                   | 4        |
| <b>3</b> | <b>Inclusion–Exclusion Principle</b>            | <b>4</b> |
| <b>4</b> | <b>Why Combinatorics Matters in Probability</b> | <b>4</b> |

## 1 Fundamental Counting Principles

### 1.1 Sum Rule

Suppose that a task can be performed in  $n_1$  ways or in  $n_2$  ways, and that the two possibilities are mutually exclusive. Then the total number of possible outcomes equals  $n_1 + n_2$ .

### 1.1.1 Example

A class consists of 15 boys and 12 girls. If we wish to choose a class representative, then the number of possible choices is

$$15 + 12 = 27.$$

## 1.2 Product Rule

Suppose that a procedure consists of  $k$  consecutive steps. If Step  $i$  can be performed in  $n_i$  ways for every  $i = 1, \dots, k$ , then the total number of possible outcomes equals

$$n_1 n_2 \cdots n_k.$$

### 1.2.1 Example

A password consists of three letters followed by two digits. Since each letter can be chosen in 26 ways and each digit in 10 ways, the total number of possible passwords equals

$$26^3 \cdot 10^2$$

## 1.3 Pigeonhole Principle

If  $n + 1$  objects are placed into  $n$  boxes, then at least one box must contain at least two objects.

### 1.3.1 Example

Among 13 people there are always at least two born in the same month. Indeed, the objects are the people and the boxes are the months of the year. Since  $13 > 12$ , the conclusion follows immediately.

## 1.4 Counting Formulas

|              | With Repetition    | Without Repetition  |
|--------------|--------------------|---------------------|
| Combinations | $\binom{n+k-1}{k}$ | $\binom{n}{k}$      |
| Arrangements | $n^k$              | $\frac{n!}{(n-k)!}$ |

## 1.5 Binomial Theorem

For every  $n \in \mathbb{Z}_+$ ,

$$(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}.$$

## 2 Fundamental Binomial Identities

### 2.0.1 Symmetry Identity

$$\binom{n}{k} = \binom{n}{n-k}.$$

### 2.0.2 Pascal Identity

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}.$$

### 2.0.3 Combinations with Repetition

$$\binom{n+k-1}{k}.$$

### 2.0.4 Sum of Binomial Coefficients

$$\sum_{k=0}^n \binom{n}{k} = 2^n.$$

### 2.0.5 Sum of Squares

$$\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n}.$$

### 2.0.6 Hockey-Stick Identity

$$\binom{n+m}{n} = \sum_{i=0}^m \binom{n+i-1}{n-1}.$$

## 2.1 Useful Consequences

From the Hockey-Stick Identity we obtain

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

and

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}.$$

## 3 Inclusion–Exclusion Principle

Suppose that  $a_1, \dots, a_N$  are objects and  $d_1, \dots, d_n$  are properties.

Let  $N(d_i)$  denote the number of objects satisfying property  $d_i$ , and let  $N(d_{i_1}, \dots, d_{i_k})$  denote the number of objects satisfying all properties  $d_{i_1}, \dots, d_{i_k}$  simultaneously.

Then the number of objects satisfying none of the properties is given by

$$N - \sum_i N(d_i) + \sum_{i < j} N(d_i, d_j) - \sum_{i < j < k} N(d_i, d_j, d_k) + \dots + (-1)^n N(d_1, \dots, d_n).$$

## 4 Why Combinatorics Matters in Probability

In many elementary probability problems, all outcomes are assumed to be equally likely. In that case, the probability of an event  $A$  is

$$\mathbb{P}(A) = \frac{|A|}{|\Omega|}.$$

Here  $\Omega$  is the sample space,  $|\Omega|$  is the total number of possible outcomes, and  $|A|$  is the number of favorable outcomes.


Therefore, probability often begins with counting.

For example, if we toss a fair coin 10 times, then there are  $2^{10}$  possible outcomes. The probability of getting exactly 4 heads is

$$\mathbb{P}(\text{exactly 4 heads}) = \frac{\binom{10}{4}}{2^{10}}.$$

Combinatorics tells us how many outcomes are possible. Probability tells us how likely they are.

Binomial Distribution: Simulation vs Theory



Birthday Problem: Simulation vs Theory

