

A Quick Introduction to Differential Forms

And Generalized Stokes's Theorem

Cola

The Chinese University of Hong Kong, Shenzhen

December 15, 2025

Unification of Calculus

The primary goal of differential forms is to provide a **unified framework** for the various integration theorems of vector calculus.

- Green's Theorem (2D)
- Stokes's Theorem (3D Curl)
- Divergence Theorem (3D)

All these are specific instances of the **Generalized Stokes's Formula**:

$$\int_{\partial D} \omega = \int_D d\omega$$

Classical Examples: 2D

Green's Formula (Line Integrals)

$$\int_{\partial D} Pdx + Qdy = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dxdy \quad (1)$$

2D Flux

$$\int_{\partial D} -Qdx + Pdy = \iint_D \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} \right) dxdy \quad (2)$$

Classical Examples: 3D

Divergence Theorem

$$\iint_{\partial D} Pdydz + Qdxdz + Rdxdy = \iiint_D \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dxdydz \quad (3)$$

In this theory, we work in \mathbb{R}^n with coordinates x^1, x^2, \dots, x^n .

0-Forms and 1-Forms

0-Forms

Smooth functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$, i.e., $f \in C^\infty(\mathbb{R}^n)$.
Example: $f(\mathbf{x}) = \sum_{i=1}^n (x^i)^2$.

1-Forms

Vector space spanned by $\{dx^1, \dots, dx^n\}$. *Example:*
 $\omega = (x^1)^2 dx^1 + (x^2) dx^2$.

2-Forms and the Wedge Product

A $\binom{n}{2}$ dimensional space spanned by $\{dx^i \wedge dx^j\}$.

Properties of \wedge

- ① **Anti-symmetry:** $dx^i \wedge dx^j = -dx^j \wedge dx^i$
- ② **Nilpotency:** $dx^i \wedge dx^i = 0$

Example: $\omega = f dx^1 \wedge dx^2 + g dx^2 \wedge dx^3$.

k -Forms ($1 \leq k \leq n$)

A space spanned by $\{dx^{i_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_k}\}$ with dimension $\binom{n}{k}$.

- **Anti-symmetry:** Swapping any two indices flips the sign.
- **Vanishing:** If $k > n$, the form is identically zero because at least one dx^i must repeat.

The Exterior Derivative d

The map $d : \Omega^k(\mathbb{R}^n) \rightarrow \Omega^{k+1}(\mathbb{R}^n)$ is defined as:

- **On 0-forms:** $df = \sum_i \frac{\partial f}{\partial x^i} dx^i$.
- **On k -forms:** If $\omega = \sum a_I dx^I$, then:

$$d\omega = \sum da_I \wedge dx^I$$

Examples of Exterior Differentiation

Example (n=2, 1-form \rightarrow 2-form)

$$\omega = Pdx + Qdy \implies d\omega = \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx \wedge dy$$

Example (n=3, 2-form \rightarrow 3-form)

$$\begin{aligned} \omega &= Pdy \wedge dz + Qdx \wedge dz + Rdx \wedge dy \\ d\omega &= \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dx \wedge dy \wedge dz \end{aligned}$$

Integration of k -Forms

Suppose $D \subseteq \mathbb{R}^n$ is a k -dimensional region and $\omega = \sum a_I dx^I$ is a k -form.

$$\int_D \omega := \sum \int_D a_{i_1 \dots i_k} dx^{i_1} \dots dx^{i_k} \quad (4)$$

Example ($n=3$): Surface integral of a 2-form:

$$\int_S \omega = \int_S P dydz + \int_S Q dx dz + \int_S R dx dy$$

The Generalized Stokes's Theorem

Theorem

Let D be a $(k + 1)$ -dimensional region with boundary ∂D . If ω is a k -form, then:

$$\int_D d\omega = \int_{\partial D} \omega \quad (5)$$

- **Dimension** $n = 3$: Recovers the Divergence and Curl theorems.
- **Dimension** $n = 2$: Recovers Green's Theorem.
- **Dimension** $n = 1$: Recovers the Fundamental Theorem of Calculus.

Case $n = 1$: Fundamental Theorem of Calculus

Let $D = [a, b]$ be a 1-dimensional region in \mathbb{R}^1 . The boundary is $\partial D = \{b\} - \{a\}$.

Derivation

Let $\omega = f(x)$ be a 0-form. Then $d\omega = f'(x)dx$. Applying Generalized Stokes:

$$\int_D d\omega = \int_{[a,b]} f'(x)dx$$

$$\int_{\partial D} \omega = f(b) - f(a)$$

This yields the standard Fundamental Theorem of Calculus:

$$\int_a^b f'(x)dx = f(b) - f(a)$$

Case $n = 2$: Green's Theorem

Let $D \subset \mathbb{R}^2$ be a region with boundary curve ∂D .

Derivation

Let $\omega = Pdx + Qdy$ be a 1-form. We calculate the 2-form $d\omega$:

$$\begin{aligned} d\omega &= dP \wedge dx + dQ \wedge dy \\ &= \left(\frac{\partial P}{\partial x} dx + \frac{\partial P}{\partial y} dy \right) \wedge dx + \left(\frac{\partial Q}{\partial x} dx + \frac{\partial Q}{\partial y} dy \right) \wedge dy \\ &= \frac{\partial P}{\partial y} dy \wedge dx + \frac{\partial Q}{\partial x} dx \wedge dy \\ &= \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx \wedge dy \end{aligned}$$

Generalized Stokes $\int_D d\omega = \int_{\partial D} \omega$ recovers Green's Theorem.

Case $n = 3$: Classical Stokes's Theorem

Let $S \subset \mathbb{R}^3$ be a 2D surface with boundary curve $\partial S = C$.

Derivation

Let $\omega = F_1 dx + F_2 dy + F_3 dz$ (a 1-form representing vector field \mathbf{F}). The exterior derivative $d\omega$ is a 2-form:

$$d\omega = \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z} \right) dy \wedge dz + \left(\frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x} \right) dz \wedge dx + \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) dx \wedge dy$$

Integrating $d\omega$ over S is equivalent to the flux of $\nabla \times \mathbf{F}$, yielding:

$$\iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S} = \oint_C \mathbf{F} \cdot d\mathbf{r}$$

Case $n = 3$: Divergence Theorem

Let $V \subset \mathbb{R}^3$ be a 3D solid with boundary surface $\partial V = S$.

Derivation

Let $\omega = Pdy \wedge dz + Qdz \wedge dx + Rdx \wedge dy$ be a 2-form.

$$\begin{aligned} d\omega &= dP \wedge dy \wedge dz + dQ \wedge dz \wedge dx + dR \wedge dx \wedge dy \\ &= \frac{\partial P}{\partial x} dx \wedge dy \wedge dz + \frac{\partial Q}{\partial y} dy \wedge dz \wedge dx + \frac{\partial R}{\partial z} dz \wedge dx \wedge dy \\ &= \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dx \wedge dy \wedge dz \end{aligned}$$

Generalized Stokes $\int_V d\omega = \int_S \omega$ recovers the Divergence Theorem:

$$\iiint_V (\nabla \cdot \mathbf{F}) dV = \iint_S \mathbf{F} \cdot d\mathbf{S}$$