


Jet collimation and acceleration

Oliver Porth
Anton Pannekoek Institute,
Amsterdam




Event Horizon Telescope



The radio jet in M87 (Virgo A)

Elliptical galaxy in center of Virgo cluster 50 Million lightyears away


There is evidence for a central dark mass of $3\text{-}6 \times 10^9 M_{\text{sun}}$



The radio jet in M87 (Virgo A)

Elliptical galaxy in center of Virgo cluster 50 Million lightyears away

There is evidence for a central dark mass of $3\text{-}6 \times 10^9 M_{\text{sun}}$




LOFAR

The radio jet in M87 (Virgo A)

Elliptical galaxy in center of Virgo cluster 50 Million lightyears away

There is evidence for a central dark mass of $3\text{-}6 \times 10^9 M_{\text{sun}}$




Composite: H. Falcke (RU Nijmegen)

The radio jet in M87 (Virgo A)

Elliptical galaxy in center of Virgo cluster 50 Million lightyears away


There is evidence for a central dark mass of $3\text{-}6 \times 10^9 M_{\text{sun}}$



The radio jet in M87 (Virgo A)

Elliptical galaxy in center of Virgo cluster 50 Million lightyears away


There is evidence for a central dark mass of $3\text{-}6 \times 10^9 M_{\text{sun}}$



The radio jet in M87 (Virgo A)

Elliptical galaxy in center of Virgo cluster 50 Million lightyears away


There is evidence for a central dark mass of $3\text{-}6 \times 10^9 M_{\text{sun}}$



The radio jet in M87 (Virgo A)

Elliptical galaxy in center of Virgo cluster 50 Million lightyears away


There is evidence for a central dark mass of $3\text{-}6 \times 10^9 M_{\text{sun}}$




The radio jet in M87 (Virgo A)

Elliptical galaxy in center of Virgo cluster 50 Million lightyears away

There is evidence for a central dark mass of $3\text{-}6 \times 10^9 M_{\text{sun}}$



Global Structure of M87




Asada & Nakamura
(2012),
Hada et al. (2013)

¹Reynolds et al. (1996), Li et al. (2009), de Gasperin et al. (2012), Broderick et al. (2015), Prieto et al. (2016)

Global Structure of M87

- Parabolic ($z \propto r^{1.7}$) over $10^5 r_s$
- Above bondi sale: **conical** streamlines

$$z \propto r$$



Asada & Nakamura
(2012),
Hada et al. (2013)

¹Reynolds et al. (1996), Li et al. (2009), de Gasperin et al. (2012), Broderick et al. (2015), Prieto et al. (2016)

Global Structure of M87

- **Parabolic ($z \propto r^{1.7}$) over $10^5 r_s$**
- Above bondi sale: **conical** streamlines

$$z \propto r$$

Jet power¹:

10^{42} erg/s - 10^{45} erg/s


BH-mass:

$3.45 \cdot 10^9 M_{\odot}$ (Walsh et al., 2013)

$6.14 \cdot 10^9 M_{\odot}$ (Gebhard et al. 2011)

Distance:

16.8 Mpc (Bird+, 2010, Blakeslee+ 2009, Cantiello 2018)




Asada & Nakamura (2012),
Hada et al. (2013)

¹Reynolds et al. (1996), Li et al. (2009), de Gasperin et al. (2012), Broderick et al. (2015), Prieto et al. (2016)



some theory...

...ok, a lot of theory



How do Jets Accelerate: The magnetic paradigm



How do Jets Accelerate: The magnetic paradigm



How do Jets Accelerate: The magnetic paradigm



$$r_{LC}\Omega = c$$




Example: light cylinder for sun

$$\Omega = \frac{2\pi}{24d} = 3 \times 10^{-6} \text{ s}^{-1}$$


$$c = 3 \times 10^{10} \text{ cm/s}$$

$$\Rightarrow r_{LC} = c/\Omega = 10^{16} \text{ cm} = 660 \text{ AU}$$

How do Jets Accelerate: The magnetic paradigm




How do Jets Accelerate: The magnetic paradigm




Adopted from Alexander (Sasha) Tchekhovskoy

How do Jets Accelerate: The magnetic paradigm



Adopted from Alexander (Sasha) Tchekhovskoy


How do Jets Accelerate: The magnetic paradigm



Adopted from Alexander (Sasha) Tchekhovskoy

force-free


How do[✓] Jets Accelerate: The magnetic paradigm



Assume the jets
are **massless (force-free)**
for simplicity

force-free

How do jets Accelerate: The magnetic paradigm




$$B_\varphi = -\frac{v_\varphi}{c} B_r = -\frac{\Omega R}{c} B_r$$

Assume the jets
are **massless (force-free)**
for simplicity

force-free

How do jets Accelerate: The magnetic paradigm




$$E = \left| -\frac{\vec{v}}{c} \times \vec{B} \right| = +\frac{\Omega R}{c} B_r$$

$$B_\varphi = -\frac{v_\varphi}{c} B_r = -\frac{\Omega R}{c} B_r$$

Assume the jets
are **massless (force-free)**
for simplicity

force-free

How do jets Accelerate: The magnetic paradigm



$$\frac{v}{c} = \left| \frac{\vec{E} \times \vec{B}}{B^2} \right| = \frac{E}{B}$$


$$E = \left| -\frac{\vec{v}}{c} \times \vec{B} \right| = +\frac{\Omega R}{c} B_r$$

$$B_\varphi = -\frac{v_\varphi}{c} B_r = -\frac{\Omega R}{c} B_r$$

Assume the jets
are **massless (force-free)**
for simplicity

force-free

How do jets Accelerate: The magnetic paradigm



$$\frac{v}{c} = \left| \frac{\vec{E} \times \vec{B}}{B^2} \right| = \frac{E}{B} \Rightarrow 1 - \frac{v^2}{c^2} = \frac{B^2 - E^2}{B^2}$$

$$E = \left| -\frac{\vec{v}}{c} \times \vec{B} \right| = +\frac{\Omega R}{c} B_r$$



$$B_\varphi = -\frac{v_\varphi}{c} B_r = -\frac{\Omega R}{c} B_r$$

Assume the jets
are **massless (force-free)**
for simplicity

force-free

How do jets Accelerate: The magnetic paradigm

$$\gamma^2 = \frac{1}{1 - v^2/c^2} = \frac{B_r^2 + B_\varphi^2}{B_r^2 + B_\varphi^2 - E^2} = 1 + \frac{B_\varphi^2}{B_r^2} = 1 + (\Omega R/c)^2$$



$$\frac{v}{c} = \left| \frac{\vec{E} \times \vec{B}}{B^2} \right| = \frac{E}{B} \Rightarrow 1 - \frac{v^2}{c^2} = \frac{B^2 - E^2}{B^2}$$

$$E = \left| -\frac{\vec{v}}{c} \times \vec{B} \right| = +\frac{\Omega R}{c} B_r$$



$$B_\varphi = -\frac{v_\varphi}{c} B_r = -\frac{\Omega R}{c} B_r$$

Assume the jets
are **massless (force-free)**
for simplicity

force-free

How do jets Accelerate: The magnetic paradigm


$$\gamma^2 = \frac{1}{1 - v^2/c^2} = \frac{B_r^2 + B_\varphi^2}{B_r^2 + B_\varphi^2 - E^2} = 1 + \frac{B_\varphi^2}{B_r^2} = 1 + (\Omega R/c)^2$$



$$-\frac{v^2}{c^2} = \frac{B^2 - E^2}{B^2}$$

Assume the jets
are **massless (force-free)**
for simplicity

mass loaded - Poynting dominated flows



mass loaded - Poynting dominated flows


Fluxes between streamlines:

$$F_M = \Gamma \rho v_p$$

$$F_E = \Gamma F_M + F_{Poynting}$$

$$F_B = B_p$$

$$F_L = -r B_p B_\phi + r \Gamma v_\phi \Gamma \rho v_p$$



mass loaded - Poynting dominated flows

Fluxes between streamlines:


$$F_M = \Gamma \rho v_p$$

$$F_E = \Gamma F_M + F_{Poynting}$$

$$F_B = B_p$$

$$F_L = -r B_p B_\phi + r \Gamma v_\phi \Gamma \rho v_p$$

Ratios of fluxes: conserved along the stream line Ψ !



Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{\text{kin}} + F_{\text{Poynting}}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$

$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r \Gamma v_\phi$$

mass loaded - Poynting dominated flows

Fluxes between streamlines:


$$F_M = \Gamma \rho v_p$$

$$F_E = \Gamma F_M + F_{Poynting}$$

$$F_B = B_p$$

$$F_L = -r B_p B_\phi + r \Gamma v_\phi \Gamma \rho v_p$$

Ratios of fluxes: conserved along the stream line Ψ !



$$\mu = \frac{F_E}{F_M} = \Gamma + \frac{|\mathbf{E} \times \mathbf{B}|}{\rho v_p \Gamma} = \Gamma \left(1 + \frac{F_{Poynting}}{F_{Kinetic}} \right)$$

Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{kin} + F_{Poynting}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$

$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r \Gamma v_\phi$$

mass loaded - Poynting dominated flows


Fluxes between streamlines:

$$F_M = \Gamma \rho v_p$$

$$F_E = \Gamma F_M + F_{Poynting}$$

$$F_B = B_p$$

$$F_L = -r B_p B_\phi + r \Gamma v_\phi \Gamma \rho v_p$$



Ratios of fluxes: conserved along the stream line Ψ !

$$\mu = \frac{F_E}{F_M} = \Gamma + \frac{|\mathbf{E} \times \mathbf{B}|}{\rho v_p \Gamma} = \Gamma \left(1 + \frac{F_{Poynting}}{F_{Kinetic}} \right)$$

$$\sigma \parallel$$

Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{kin} + F_{Poynting}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$

$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r \Gamma v_\phi$$

mass loaded - Poynting dominated flows


Fluxes between streamlines:

$$F_M = \Gamma \rho v_p$$

$$F_E = \Gamma F_M + F_{Poynting}$$

$$F_B = B_p$$

$$F_L = -r B_p B_\phi + r \Gamma v_\phi \Gamma \rho v_p$$



Ratios of fluxes: conserved along the stream line Ψ !

$$\mu = \frac{F_E}{F_M} = \Gamma + \frac{|\mathbf{E} \times \mathbf{B}|}{\rho v_p \Gamma} = \Gamma \left(1 + \frac{F_{Poynting}}{F_{Kinetic}} \right)$$

$$\sigma \parallel$$

$$\mu = \Gamma(1 + \sigma); \quad \Gamma \rightarrow \mu(\sigma \rightarrow 0)$$

Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{kin} + F_{Poynting}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$

$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r \Gamma v_\phi$$

mass loaded - Poynting dominated flows


Fluxes between streamlines:

$$F_M = \Gamma \rho v_p$$

$$F_E = \Gamma F_M + F_{Poynting}$$

$$F_B = B_p$$

$$F_L = -r B_p B_\phi + r \Gamma v_\phi \Gamma \rho v_p$$



Ratios of fluxes: conserved along the stream line Ψ !

$$\mu = \frac{F_E}{F_M} = \Gamma + \frac{|\mathbf{E} \times \mathbf{B}|}{\rho v_p \Gamma} = \Gamma \left(1 + \frac{F_{Poynting}}{F_{Kinetic}} \right)$$

$$\mu = \Gamma(1 + \sigma); \quad \Gamma \rightarrow \mu(\sigma \rightarrow 0)$$

Maximum achievable Lorentz-factor: μ

Jet accelerates by decreasing its magnetization σ

Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{kin} + F_{Poynting}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$


$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r \Gamma v_\phi$$

Force-free breaks at the fast point!

speed of fast waves?


$$v_{FMS}^2 = \frac{B'^2}{\rho + B'^2} = \frac{\sigma}{1 + \sigma}$$



Force-free breaks at the fast point!

speed of fast waves?

$$v_{FMS}^2 = \frac{B'^2}{\rho + B'^2} = \frac{\sigma}{1 + \sigma} \quad 1 - v_{FMS}^2 = \frac{1}{1 + \sigma}$$




Force-free breaks at the fast point!

speed of fast waves?

$$v_{FMS}^2 = \frac{B'^2}{\rho + B'^2} = \frac{\sigma}{1 + \sigma} \quad 1 - v_{FMS}^2 = \frac{1}{1 + \sigma}$$

$$\Gamma_{FMS}^2 = 1 + \sigma \Rightarrow \Gamma_{FMS} \approx \sigma^{1/2}$$




Force-free breaks at the fast point!

speed of fast waves?

$$v_{FMS}^2 = \frac{B'^2}{\rho + B'^2} = \frac{\sigma}{1 + \sigma} \quad 1 - v_{FMS}^2 = \frac{1}{1 + \sigma}$$

$$\Gamma_{FMS}^2 = 1 + \sigma \Rightarrow \Gamma_{FMS} \approx \sigma^{1/2}$$

$$\Gamma_{FMS}^3 = \Gamma(1 + \sigma) \Rightarrow \Gamma_{FMS} = \mu^{1/3}$$



Force-free breaks at the fast point!


force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves?

$$v_{FMS}^2 = \frac{B'^2}{\rho + B'^2} = \frac{\sigma}{1 + \sigma} \quad 1 - v_{FMS}^2 = \frac{1}{1 + \sigma}$$

$$\Gamma_{FMS}^2 = 1 + \sigma \Rightarrow \Gamma_{FMS} \approx \sigma^{1/2}$$

$$\Gamma_{FMS}^3 = \Gamma(1 + \sigma) \Rightarrow \Gamma_{FMS} = \mu^{1/3}$$



Force-free breaks at the fast point!

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

$$\Gamma = \mu^{1/3}$$

Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

$$\Gamma = \mu^{1/3} \ll \mu$$

Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

$$\Gamma = \mu^{1/3} \ll \mu = 10^3$$

Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

$$10 = \Gamma = \mu^{1/3} \ll \mu = 10^3$$


Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

$$10 = \Gamma = \mu^{1/3} \ll \mu = 10^3$$




Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

$$10 = \Gamma = \mu^{1/3} \ll \mu = 10^3$$



Adopted from Sasha


Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

$$10 = \Gamma = \mu^{1/3} \ll \mu = 10^3$$




Force-free breaks at the fast point!

force-free: $\sigma = \infty$ fast waves travel at c

speed of fast waves: $\Gamma_{\text{FMS}} = \sigma^{1/2}$

force-free breaks down when $\Gamma = \Gamma_{\text{FMS}} = \sigma^{1/2} = \mu^{1/3}$

$$10 = \Gamma = \mu^{1/3} \ll \mu = 10^3$$



Adopted from Sasha

(unconfined) relativistic jets accelerate
inefficiently
=> σ -Problem

Poynting dominated flows - in axisymmetry

Beyond the light cylinder ($r\Omega \equiv c$) :

$$v_\phi \ll r\Omega; \quad v_p \rightarrow 1 : \quad E = |B_\phi|; \quad |B_\phi| = r\Omega B_p$$


Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{\text{kin}} + F_{\text{Poynting}}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$

$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r\Gamma v_\phi$$



Poynting dominated flows - in axisymmetry

Beyond the light cylinder ($r\Omega \equiv c$) :

$$v_\phi \ll r\Omega; \quad v_p \rightarrow 1 : \quad E = |B_\phi|; \quad |B_\phi| = r\Omega B_p$$

$$\mu = \Gamma + \frac{r\Omega B_p |B_\phi|}{\rho v_p \Gamma}$$

$$\phi = r^2 B_p$$

$$\Rightarrow \mu = \Gamma + \frac{\Omega(\Psi)^2 r^2 B_p}{k(\Psi)}$$

Michel (1969)


Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{\text{kin}} + F_{\text{Poynting}}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$

$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r\Gamma v_\phi$$



Poynting dominated flows - in axisymmetry

Beyond the light cylinder ($r\Omega \equiv c$) :


$$v_\phi \ll r\Omega; \quad v_p \rightarrow 1 : \quad E = |B_\phi|; \quad |B_\phi| = r\Omega B_p$$

$$\mu = \Gamma + \frac{r\Omega B_p |B_\phi|}{\rho v_p \Gamma}$$

$$\phi = r^2 B_p$$

$$\Rightarrow \mu = \Gamma + \frac{\Omega(\Psi)^2 r^2 B_p}{k(\Psi)}$$

Michel (1969)



Acceleration proceeds via differential „bunching“ of field lines $\phi \rightarrow 0$
 \Rightarrow No acceleration without collimation!

E.g. straight (monopolar) field lines:

$$B_p(r) \propto r^{-2} \Rightarrow \phi = \text{const.}$$

Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{\text{kin}} + F_{\text{Poynting}}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$

$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r\Gamma v_\phi$$

Poynting dominated flows - in axisymmetry

Beyond the light cylinder ($r\Omega \equiv c$) :


$$v_\phi \ll r\Omega; \quad v_p \rightarrow 1 : \quad E = |B_\phi|; \quad |B_\phi| = r\Omega B_p$$

$$\mu = \Gamma + \frac{r\Omega B_p |B_\phi|}{\rho v_p \Gamma}$$

$$\phi = r^2 B_p$$

$$\Rightarrow \mu = \Gamma + \frac{\Omega(\Psi)^2 r^2 B_p}{k(\Psi)}$$

Michel (1969)



Acceleration proceeds via differential „bunching“ of field lines $\phi \rightarrow 0$
 \Rightarrow No acceleration without collimation!

E.g. straight (monopolar) field lines:

$$B_p(r) \propto r^{-2} \Rightarrow \phi = \text{const.}$$

Equipartition scale for collimating ($z \propto r^{1.5}$) jet :

$$z_{\text{eq}} = 2.7 \times 10^4 r_S \left(\frac{\mu}{30} \right)^3 = 2 \left(\frac{\mu}{30} \right)^3 \frac{M_\bullet}{10^9 M_\odot} \text{pc}$$

Conserved quantities:

$$\mu(\Psi) \equiv \frac{F_{\text{kin}} + F_{\text{Poynting}}}{\rho \Gamma v_p}$$

$$r\Omega(\Psi) \equiv v_\phi - v_p \frac{B_\phi}{B_p}$$


$$k(\Psi) \equiv \frac{\rho \Gamma v_p}{B_p}$$

$$l(\Psi) \equiv -\frac{I}{2\pi k} + r\Gamma v_\phi$$

Barkov & Komissarov (2008)
 Beskin & Nokhrina (2009)
 Porth et al. (2011)

Conditions for relativistic jet formation


- Rotating magnetosphere with open magnetic fields (need Blandford)
- High magnetisation: Need to extract energy
- Collimating agent: disk wind, stellar remnant, shocked gas




Wei et al (2012)

Conditions for relativistic jet formation

- Rotating magnetosphere with open magnetic fields (need Blandford)
- High magnetisation: Need to extract energy
- Collimating agent: disk wind, stellar remnant, shocked gas




Wei et al (2012)





Conditions for relativistic jet formation

- Rotating magnetosphere with open magnetic fields (need Blandford)
- High magnetisation: Need to extract energy
- Collimating agent: disk wind, stellar remnant, shocked gas





Wei et al (2012)



Conditions for relativistic jet formation

- Rotating magnetosphere with open magnetic fields (need Blandford)
- High magnetisation: Need to extract energy
- Collimating agent: disk wind, stellar remnant, shocked gas




2D numerical simulations

- 5 orders of magnitude: can be directly compared to observations
- Jets gradually accelerate
 - as a power-law in distance
 - to relativistic Lorentz factors
- Jets efficiently convert magnetic to kinetic energy (to equipartition)
- Pinching and shear instabilities!
 - Mass-loading and jet deceleration



Koushik
Chatterjee




2D numerical simulations

- 5 orders of magnitude: can be directly compared to observations
- Jets gradually accelerate
 - as a power-law in distance
 - to relativistic Lorentz factors
- Jets efficiently convert magnetic to kinetic energy (to equipartition)
- Pinching and shear instabilities!
 - Mass-loading and jet deceleration




Koushik
Chatterjee




Jet instabilities

- (AGN) Jets can cover 10^5 - 10^7 of initial radii without disruption
- Optical synchrotron needs continuous particle re-acceleration thus dissipation (e.g. Meisenheimer 2003)
- How can we avoid disrupting fluid instabilities?
- What is the deciding factor between the FR1 and FR2 division?
- How can we have dissipation without disruption?





Credit: A. Bridle



Many thanks to Manel Peruchó for the diagram (in press)!


Dangerous jet instabilities






Kelvin Helmholtz type instabilities (KHI):
(Turland & Scheuer (1976); Ferrari+ (1978);
Hardee (2004); Perucho et al (2004,07);
Bodo et al (2006); Rossi et al (2008); ...)

Can be stabilized with:
Relativistic bulk motion,
Thick shear layer, poloidal fields,
jet expansion


Can be stabilized with:
Relativistic bulk motion, shear, being force-free,
electric fields, poloidal fields, **jet expansion**



Dangerous jet instabilities





Current driven instabilities (CDI): Pinch, Kink and higher orders
(Bateman (1978), Begelman (1998), Appl et al. (2000), Baty (2005), Lery et al. (2000); Narayan et al. (2009); Moll et al (2011); Mizuno et al (2011,12,14); O'Neill et al (2012); Mignone et al. (2013); Anjiri+ (2014); ...)




Can be stabilized with:
Relativistic bulk motion, shear, being force-free,
electric fields, poloidal fields, jet expansion

Dangerous jet instabilities




MHD jet in powerlaw atmosphere




External gas:
 $p_{\text{ext}} = p_0 z^{-\kappa}$



Sufficient criterion for jet stability:
Causally disconnected flows are globally stable!



MHD jet in powerlaw atmosphere




Sufficient criterion for jet stability:
Causally disconnected flows are globally stable!




External gas:
 $p_{\text{ext}} = p_0 z^{-\kappa}$



MHD jet in powerlaw atmosphere



- Waves traveling back and forth the jet giving rise to oscillations of jet boundary
- For $\kappa \geq 2$ flow is conical and no waves traverse across the jet
- Jet boundary becomes disconnected for $\kappa \geq 2$





looking inside an expanding jet




- letting jets expand increases their stability
- Formation of sub-structure inside the jet
- shocks and magnetic reconnection
=> particle acceleration

looking inside an expanding jet




- letting jets expand increases their stability
- Formation of sub-structure inside the jet
- shocks and magnetic reconnection
=> particle acceleration

(a)



6.2 Myr

(b)



6.2 Myr

(c)



5.7 Myr

“communication”: important for jet stability, FR1/FR2 division
 (OP +2015, Tchekhovskoy+ 2016)

- Stabilization due to inhibited causal contact across the jet
 - Higher jet power: narrower Mach cones => more stable
 - Less collimated jet: Mach cone points away from axis => more stable
- Steep ambient pressure gradient enhances jet stability: necessarily globally stable for $\kappa \geq 2$
- Formation of unstable jet core:
 - can seed jet internal turbulence
 - dissipates energy and thus aids in jet acceleration

quick summary

- Rotating magnetospheres: jets!
- Magnetic acceleration: gradual, depends on (external) collimation
- Plethora of instabilities threatening jets, some good (dissipation), some bad (disruption)
- Communication and causality is key in jet acceleration and stability