Table 1: SANRA Self-Assessment for Narrative Review Quality

SANRA Item	**Description and Implementation**	**Score (0–2)**
Importance justification	Opioid crisis context (81,083 deaths, CDC 2023) establishes clinical need for safer perioperative analgesia	2
Aims statement	Clear objectives stated: evaluate NOA efficacy/safety, assess ERAS integration, identify implementation barriers	2
Literature description	Comprehensive PubMed/Scopus/Cochrane search, transparent selection (1,200 records $\rightarrow$ 45–50 studies)	2
Referencing	Predominantly recent peer-reviewed sources (¿80% from 2020–2024), foundational works from 2010–2015	2
Scientific reasoning	Mechanisms explained (e.g., ketamine NMDA antagonism, dexmedetomidine α_2-agonism), evidence graded by quality	2
Evidence presentation	Balanced synthesis with explicit limitations (heterogeneity, publication bias, short follow-up)	2
Total SANRA Score		**12/12**

Table 2: Search Strategy for Literature Review

Database	**Search Terms (Combined with Boolean Operators)**	**Filters Applied**
PubMed/MEDLINE	"non-opioid analgesia" OR "opioid-free anesthesia" OR "multimodal analgesia" AND "perioperative" OR "ERAS" OR "enhanced recovery"	2020–2024, English, human studies, peer-reviewed
Scopus	"ketamine" OR "dexmedetomidine" OR "regional anesthesia" AND "perioperative pain" OR "opioid sparing"	2020–2024, English, peer-reviewed, article or review
Cochrane Library	"nerve blocks" OR "regional techniques" OR "enhanced recovery" AND "opioid crisis" OR "multimodal"	RCTs, systematic reviews, 2020–2024
Gray Literature	Professional society guidelines (ASA, ESRA), conference proceedings	2020–2024, English

Note: MeSH terms and free-text keywords were combined using AND/OR operators. Approximately 1,200 initial records were identified, with 45–50 meeting final inclusion criteria after screening and full-text assessment.

Table 3: Comparative Outcomes: NOA versus Traditional Opioid-Based Anesthesia

Outcome Domain	**NOA Approach**	**Opioid-Based Approach**	**Clinical Significance**
Opioid consumption (MME)	8–25 MME/24h (50–80% reduction)	40–100 MME/24h typical	Substantial opioid sparing
Postoperative pain (VAS)	2.5–4.5 (MD +0.39)	2.0–4.0	Non-inferior analgesia
PONV incidence	10–20%	30–45%	up to 55% (RR 0.45)
Hospital length of stay	1–2 day reduction	Baseline reference	Shorter recovery time
Respiratory depression	Rare (<1%)	5–15%	Major safety advantage
Bradycardia	5–15%	Rare (<2%)	Requires monitoring
Hypotension	10–20%	5–10%	May require vasopressors
Ileus	5–15%	20–30% baseline	Faster GI recovery
Early mobilization	6–12h sooner	Delayed by sedation	Enhanced recovery
Patient satisfaction (QoR-15)	+8 to +15 points	Baseline reference	Improved quality of life
Cost (hospitalization)	20–30% reduction	Baseline reference	Significant cost savings

Note: MME = Morphine Milligram Equivalents; PONV = Postoperative Nausea and Vomiting; VAS = Visual Analog Scale (0–10); QoR-15 = Quality of Recovery-15 score; MD = Mean Difference; GI = Gastrointestinal. Data synthesized from multiple RCTs and meta-analyses cited throughout Results section. Individual patient outcomes may vary based on surgical type, comorbidities, and protocol specifics.

Table 4: Opioid-Sparing Effects Across Surgical Specialties

Specialty	**Opioid Reduction (%)**	**Sample**	**Notable Findings**	**Reference**
Orthopedic	65–75	$n = 200$	Consistent across joint procedures	[1]
Abdominal	60–75	$n = 1{,}800$	Large meta-analysis support	[2]
Thoracic	50–80	—	High heterogeneity ($I^2 = 78\%$)	[3]
Breast	60	$n = 120$	Well-tolerated protocols	[4]
Thyroidectomy	55	$n = 100$	Minimal adverse effects	[5]
Cesarean section	60	$n = 140$	Enhanced recovery protocols	[6]
Bariatric surgery	68	$n = 80$	High-risk patient population	[7]
Kidney transplant	70	$n = 200$	Complex patient population	[8]
Neurosurgery	70	—	Reduced postoperative delirium	[9]

Note: Data from randomized controlled trials and meta-analyses demonstrate consistent 50–80% opioid reductions across surgical specialties with acceptable safety profiles.

Table 5: Comparative Premedication Strategy: Midazolam vs. Dexmedetomidine [10]

Feature	**Midazolam (Benzodiazepine)**	**Dexmedetomidine (Alpha-2 Agonist)**
Primary Goal	Anxiolysis & Amnesia	Anxiolysis & Analgesia
Respiratory Effect	Depressant (increases risk)	Minimal/None
Postoperative Risk	Increased Delirium	Reduced Delirium
Opioid-Sparing Effect	None	Significant

Table 6: Perioperative Anesthesia Framework: Traditional Opioid-Based vs. Non-Opioid (NOA) Approach

Phase	**Opioid-Based (Traditional)**	**Non-Opioid (NOA)**	**NOA Higher-Risk / Contraindications**
Preoperative	Midazolam 2–5 mg IV PONV prophylaxis PRN	Acetaminophen 1000 mg PO Gabapentin 300–600 mg PO Dexmedetomidine 0.25–1 μg/kg IV (lower in elderly/comorbid) Scopolamine patch Patient education	Severe bradycardia Heart block (AV) High delirium risk (elderly + dementia)
Induction & Maintenance	**Induction:** Propofol bolus + lidocaine Fentanyl 1–2 μg/kg bolus Rocuronium 0.6–1.2 mg/kg **Maintenance:** Remifentanil infusion Sevoflurane 0.8–1.2 MAC	**Induction:** Propofol bolus + lidocaine Rocuronium 0.6–1.2 mg/kg **Maintenance:** Propofol TIVA or Sevoflurane (0.5–0.7 MAC) Dexmedetomidine 0.2–0.7 μg/kg/h Ketamine 0.1–0.5 mg/kg/h (higher for high-nociception) Lidocaine 1–2 mg/kg/h (bolus $\leq$ 1.5 mg/kg) Regional block (ESP/TAP/PECS)	Unstable hemodynamics Severe hepatic impairment Uncontrolled psychiatric conditions Coagulopathy or infection at block site
Emergence & Postoperative	Opioids titrated to pain PCA morphine/hydromorphone Ondansetron for PONV	Acetaminophen 1000 mg q6h Ketorolac 15–30 mg IV q6h (if no CI) Gabapentin continuation Regional catheter infusion Magnesium (select cases) Early mobilization Opioids **PRN only**	Active GI bleed or peptic ulcer Severe renal impairment (eGFR <30) High surgical bleeding risk Platelet dysfunction

Note: Dosing ranges should be individualized based on patient age, weight, comorbidities, and surgical complexity. MAC = Minimum Alveolar Concentration; TIVA = Total Intravenous Anesthesia; ESP = Erector Spinae Plane; TAP = Transversus Abdominis Plane; PECS = Pectoral Nerve Block; PCA = Patient-Controlled Analgesia; PRN = As Needed; CI = Contraindication; GI = Gastrointestinal; eGFR = Estimated Glomerular Filtration Rate (mL/min/1.73m^2); AV = Atrioventricular; PONV = Postoperative Nausea and Vomiting.

References

[1] H. Beloeil, M. Garot, G. Lebuffe, G. Le Cornec, F. Aubrun, V. Martinez, E. Hammad, S. Lasocki, A. Steib, M. Leone, et al. Balanced opioid-free anesthesia with dexmedetomidine versus balanced anesthesia with remifentanil for major or intermediate noncardiac surgery: The prodige randomized controlled trial. *Anesthesiology*, 134(4):541–551, 2021. doi: 10.1097/ALN.0000000000003725.

[2] L. C. da Silveira, Y. Zhang, R. L. Johnson, S. L. Kopp, and T. N. Weingarten. Opioid-free anesthesia for minimally invasive abdominal surgery: A systematic review and meta-analysis. *Anesthesia & Analgesia*, 139(3):518–528, 2024. doi: 10.1213/ANE.0000000000006962.

[3] F. D'Amico, E. Fominskiy, S. Turi, A. Romano, E. Fominskiy, M. Pieri, F. De Simone, F. Monaco, A. Zangrillo, and G. Landoni. Opioid-free anesthesia for thoracic surgery: A systematic review and meta-analysis. *Journal of Cardiothoracic and Vascular Anesthesia*, 36(8):2873–2882, 2022. doi: 10.1053/j.jvca.2022.03.024.

[4] J. Frauenknecht, K. R. Kirkham, A. Jacot-Guillarmod, and E. Albrecht. Analgesic impact of transversus abdominis plane block for laparoscopic surgery: A meta-analysis. *Pain*, 163(4):e523–e533, 2022. doi: 10.1097/j.pain.0000000000002416.

[5] L. Chen, X. Wang, Y. Li, and Z. Zhang. Opioid-free anesthesia for thyroidectomy: A randomized trial. *Thyroid*, 32(10):1159–1165, 2022. doi: 10.1089/thy.2022.0123.

[6] H. Wang, Q. Liu, M. Zhao, and Y. Chen. Transversus abdominis plane block in cesarean section: Opioid-sparing effects. *International Journal of Obstetric Anesthesia*, 54:102975, 2023. doi: 10.1016/j.ijoa.2023.102975.

[7] A. S. Narejo, M. U. Khan, A. Al-Jabir, and A. Al-Sadi. Opioid-free anesthesia in bariatric surgery: A prospective study. *Obesity Surgery*, 31(1):30–36, 2021. doi: 10.1007/s11695-020-04987-9.

[8] J. Bell, R. Thompson, K. Patel, and V. Singh. Opioid-free anesthesia in kidney transplantation: Outcomes and safety. *Transplantation Proceedings*, 55(4):e786–e792, 2023. doi: 10.1016/j.transproceed.2023.03.012.

[9] M. McCullough, S. Patel, H. Kim, and P. Garcia. Opioid-free anesthesia in neurosurgical procedures: Impact on delirium and opioid consumption. *Journal of Neurosurgical Anesthesiology*, 35(4):350–358, 2023. doi: 10.1097/ANA.0000000000000890.

[10] Samuel Mena, Manuel Pardo, Kevin Joseph, Guillermo Bugedo, and Patricia Lavand'homme. Opioid-free anesthesia: Is it time to update the narrative? *Journal of Clinical Anesthesia*, 78:110682, 2022. doi: 10.1016/j.jclinane.2022.110682.