

# Variational Mixture of Normalizing Flows

Guilherme Grijó Pen Freitas Pires

November 15, 2019

Thesis to obtain the Master of Science degree in  
**Electrical and Computer Engineering**



# Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions

# Introduction and Motivation

- Deep generative models have been an active area of research, with promising results.

# Introduction and Motivation

- Deep generative models have been an active area of research, with promising results.
  - **Implicit distributions:** Generative adversarial networks [Goodfellow et al., 2014], Variational Autoencoder [Kingma and Welling, 2014]
    - Don't allow explicit access to the density function

# Introduction and Motivation

- Deep generative models have been an active area of research, with promising results.
  - **Implicit distributions:** Generative adversarial networks [Goodfellow et al., 2014], Variational Autoencoder [Kingma and Welling, 2014]
    - Don't allow explicit access to the density function
  - **Explicit distributions:** Normalizing flows [Rezende and Mohamed, 2015]
    - Allow explicit access to the density function
    - Lack an approach to introduce discrete structure (multi-modality) in the modelled distribution.

# Introduction and Motivation: Goal

- The goal of this work was the development of a mixture of flexible distributions.

# Introduction and Motivation: Goal

- The goal of this work was the development of a mixture of flexible distributions.
- This requires answering two questions:

# Introduction and Motivation: Goal

- The goal of this work was the development of a mixture of flexible distributions.
- This requires answering two questions:
  - What should be the “family” of the mixture components?



# Introduction and Motivation: Goal

- The goal of this work was the development of a mixture of flexible distributions.
- This requires answering two questions:
  - What should be the “family” of the mixture components?
  - How should the mixture components’ parameters be estimated?

- Mixture Models

# Outline

- Mixture Models
- Variational Inference

# Outline

- Mixture Models
- Variational Inference
  - The chosen framework for estimating the parameters of the proposed model

# Outline

- Mixture Models
- Variational Inference
  - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows

- Mixture Models
- Variational Inference
  - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows
  - The chosen family for the mixture model components

- Mixture Models
- Variational Inference
  - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows
  - The chosen family for the mixture model components
- Variational Mixture of Normalizing Flows

- Mixture Models
- Variational Inference
  - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows
  - The chosen family for the mixture model components
- Variational Mixture of Normalizing Flows
- Experiments and results



- Mixture Models
- Variational Inference
  - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows
  - The chosen family for the mixture model components
- Variational Mixture of Normalizing Flows
- Experiments and results
- Conclusions and future work

# Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models**
- 3 Normalizing Flows
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions

# Mixture Models: Definition

- A mixture model is used to model data that is assumed to contain subgroups.

# Mixture Models: Definition

- A mixture model is used to model data that is assumed to contain subgroups.
- Typically, it is assumed that the “subgroup-conditional” distributions belong to the same family, but have different parameters.


# Mixture Models: Definition

- A mixture model is used to model data that is assumed to contain subgroups.
- Typically, it is assumed that the “subgroup-conditional” distributions belong to the same family, but have different parameters.
- Formally, a mixture model’s joint distribution (for a single instance  $x$ ) is given by:


$$p(\boldsymbol{x}, c) = p(\boldsymbol{x}|c)p(c),$$

where  $c$  is the latent variable that indexes the subgroup to which  $x$  belongs

# Mixture Models: Plate diagram



# Mixture Models: Mixture of Gaussians



# Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows**
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions



# Normalizing Flows: Change of Variables

Given

$$\begin{cases} z \sim p(z) \\ x = g(z; \theta) \end{cases}$$

# Normalizing Flows: Change of Variables

Given

$$\begin{cases} z \sim p(z) \\ x = g(z; \theta) \end{cases}$$

then:

$$\begin{aligned} f_X(\mathbf{x}) &= f_Z(g^{-1}(\mathbf{x}; \theta)) \left| \det \left( \frac{d}{d\mathbf{x}} g^{-1}(\mathbf{x}; \theta) \right) \right| \\ &= f_Z(g^{-1}(\mathbf{x}; \theta)) \left| \det \left( \frac{d}{d\mathbf{z}} g(\mathbf{z}; \theta) \right) \Big|_{\mathbf{z}=g^{-1}(\mathbf{x}; \theta)} \right|^{-1} \end{aligned}$$

# Normalizing Flows: Change of Variables

Given

$$\begin{cases} z \sim p(z) \\ x = g(z; \theta) \end{cases}$$

then:

$$\begin{aligned} f_X(x) &= f_Z(g^{-1}(x; \theta)) \left| \det \left( \frac{d}{dx} g^{-1}(x; \theta) \right) \right| \\ &= f_Z(g^{-1}(x; \theta)) \left| \det \left( \frac{d}{dz} g(z; \theta) \right) \Big|_{z=g^{-1}(x; \theta)} \right|^{-1} \end{aligned}$$

This can be optimized w.r.t.  $\theta$ , so as to approximate an arbitrary distribution

# Normalizing Flows: Change of Variables

The described in the previous slide can be useful if

# Normalizing Flows: Change of Variables

The described in the previous slide can be useful if

- The base density has a closed form expression and is easy to sample from

# Normalizing Flows: Change of Variables

The described in the previous slide can be useful if

- The base density has a closed form expression and is easy to sample from
- The determinant of the Jacobian of  $g$  is computationally cheap - not the case, in general

# Normalizing Flows: Change of Variables

The described in the previous slide can be useful if

- The base density has a closed form expression and is easy to sample from
- The determinant of the Jacobian of  $g$  is computationally cheap - not the case, in general
- The gradient of the determinant of the Jacobian of  $g$  w.r.t  $\theta$  is computationally cheap

# Normalizing Flows: Change of Variables

The framework of Normalizing Flows consists of composing several transformations that fulfill the three listed conditions.



# Normalizing Flows: Affine Coupling Layer

An example of such a transformation is the Affine Coupling Layer [Dinh, Sohl-Dickstein, and Bengio, 2017].

Splitting  $z$  into  $(z_1, z_2)$ ,

# Normalizing Flows: Affine Coupling Layer

An example of such a transformation is the Affine Coupling Layer [Dinh, Sohl-Dickstein, and Bengio, 2017].

Splitting  $z$  into  $(z_1, z_2)$ ,

$$\begin{cases} x_1 &= z_1 \odot \exp(s(z_2)) + t(z_2) \\ x_2 &= z_2. \end{cases}$$

# Normalizing Flows: Affine Coupling Layer

An example of such a transformation is the Affine Coupling Layer [Dinh, Sohl-Dickstein, and Bengio, 2017].

Splitting  $z$  into  $(z_1, z_2)$ ,

$$\begin{cases} x_1 &= z_1 \odot \exp(s(z_2)) + t(z_2) \\ x_2 &= z_2. \end{cases}$$

This transformation has the following Jacobian matrix:

$$J_{f(z)} = \begin{bmatrix} \frac{\partial x_1}{\partial z_1} & \frac{\partial x_1}{\partial z_2} \\ \frac{\partial x_2}{\partial z_1} & \frac{\partial x_2}{\partial z_2} \end{bmatrix} = \begin{bmatrix} \text{diag}(\exp(s(z_2))) & \frac{\partial x_1}{\partial z_2} \\ \mathbf{0} & I \end{bmatrix}$$

# Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows
- 4 Variational Inference**
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions

# Variational Inference: Preamble

Consider a joint probability distribution  $p(\boldsymbol{x}, \boldsymbol{z})$ .

# Variational Inference: Preamble

Consider a joint probability distribution  $p(x, z)$ . Suppose  $x$  is observed and  $z$  is latent.

# Variational Inference: Preamble

Consider a joint probability distribution  $p(\mathbf{x}, \mathbf{z})$ . Suppose  $\mathbf{x}$  is observed and  $\mathbf{z}$  is latent. If we want to infer the most probable values of  $\mathbf{z}$ , given  $\mathbf{x}$ , by Bayes' Law:

$$\begin{aligned} p(\mathbf{z}|\mathbf{x}) &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{p(\mathbf{x})} \\ &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{\int p(\mathbf{x}|\mathbf{z}')p(\mathbf{z}')d\mathbf{z}'} \end{aligned}$$

# Variational Inference: Preamble

Consider a joint probability distribution  $p(\mathbf{x}, \mathbf{z})$ . Suppose  $\mathbf{x}$  is observed and  $\mathbf{z}$  is latent. If we want to infer the most probable values of  $\mathbf{z}$ , given  $\mathbf{x}$ , by Bayes' Law:

$$\begin{aligned} p(\mathbf{z}|\mathbf{x}) &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{p(\mathbf{x})} \\ &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{\int p(\mathbf{x}|\mathbf{z}')p(\mathbf{z}')d\mathbf{z}'} \end{aligned}$$

**Problem:** The integral in the denominator is intractable for most interesting models.



# Variational Inference: Preamble

Consider a joint probability distribution  $p(\mathbf{x}, \mathbf{z})$ . Suppose  $\mathbf{x}$  is observed and  $\mathbf{z}$  is latent. If we want to infer the most probable values of  $\mathbf{z}$ , given  $\mathbf{x}$ , by Bayes' Law:

$$\begin{aligned} p(\mathbf{z}|\mathbf{x}) &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{p(\mathbf{x})} \\ &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{\int p(\mathbf{x}|\mathbf{z}')p(\mathbf{z}')d\mathbf{z}'} \end{aligned}$$

**Problem:** The integral in the denominator is intractable for most interesting models.

- Variational inference is an approximate inference framework, that can be used to overcome this intractability.

# Variational Inference: Goal

Given a *variational* family  $q(\mathbf{z}; \boldsymbol{\lambda})$ , find the parameters  $\boldsymbol{\lambda}$  that minimize the Kullback-Leibler divergence between  $q(\mathbf{z}; \boldsymbol{\lambda})$  and  $p(\mathbf{z}|\mathbf{x})$

# Variational Inference: ELBO

$$KL(q||p) = \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z}$$

# Variational Inference: ELBO

$$\begin{aligned}KL(q||p) &= \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} \\ &= \int q(\mathbf{z}) (\log q(\mathbf{z}) - \log p(\mathbf{z}|\mathbf{x})) d\mathbf{z}\end{aligned}$$

# Variational Inference: ELBO

$$\begin{aligned}KL(q||p) &= \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - \log p(\mathbf{z}|\mathbf{x})) d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - (\log p(\mathbf{x}, \mathbf{z}) - \log p(\mathbf{x}))) d\mathbf{z}\end{aligned}$$

# Variational Inference: ELBO

$$\begin{aligned}KL(q||p) &= \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - \log p(\mathbf{z}|\mathbf{x})) d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - (\log p(\mathbf{x}, \mathbf{z}) - \log p(\mathbf{x}))) d\mathbf{z} \\&= \mathbb{E}_q[\log q(\mathbf{z})] - \mathbb{E}_q[\log p(\mathbf{x}, \mathbf{z})] + \log p(\mathbf{x})\end{aligned}$$

# Variational Inference: ELBO

$$\begin{aligned}KL(q||p) &= \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - \log p(\mathbf{z}|\mathbf{x})) d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - (\log p(\mathbf{x}, \mathbf{z}) - \log p(\mathbf{x}))) d\mathbf{z} \\&= \mathbb{E}_q[\log q(\mathbf{z})] - \mathbb{E}_q[\log p(\mathbf{x}, \mathbf{z})] + \log p(\mathbf{x})\end{aligned}$$

Which yields the lower bound (ELBO):

$$\begin{aligned}ELBO(q) &= \mathbb{E}_q[\log p(\mathbf{x}, \mathbf{z})] - \mathbb{E}_q[\log q(\mathbf{z})] \\&= \mathbb{E}_q[\log p(\mathbf{x}|\mathbf{z})] + \mathbb{E}_q[\log p(\mathbf{z})] - \mathbb{E}_q[\log q(\mathbf{z})]\end{aligned}$$

# Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows**
- 6 Conclusions



How can we leverage the flexibility of normalizing flows, and endow it with multimodal, discrete structure, like in a mixture model?

How can we leverage the flexibility of normalizing flows, and endow it with multimodal, discrete structure, like in a mixture model?

Mixture of normalizing flows.

How can we leverage the flexibility of normalizing flows, and endow it with multimodal, discrete structure, like in a mixture model?

Mixture of normalizing flows.  $\rightarrow$  Approximate inference is required.

Recall the ELBO:


$$ELBO(q) = \mathbb{E}_q[\log p(\boldsymbol{x}|\boldsymbol{z})] + \mathbb{E}_q[\log p(\boldsymbol{z})] - \mathbb{E}_q[\log q(\boldsymbol{z})]$$

Recall the ELBO:



$$ELBO(q) = \mathbb{E}_q[\log p(\mathbf{x}|\mathbf{z})] + \mathbb{E}_q[\log p(\mathbf{z})] - \mathbb{E}_q[\log q(\mathbf{z})]$$

Let the variational posterior  $q(\mathbf{z}|\mathbf{x})$  be parameterized by a neural network. We jointly optimize this objective, hence we learn the variational posterior and the generative components simultaneously.

# VMoNF: Overview





# VMoNF: Experiments - Pinwheel (5 wings)





## Training Animation



# VMoNF: Experiments - 2 Circles




# VMoNF: Experiments - 2 Circles (semi supervised)



Note: 32 labeled points, 1024 unlabeled points

# VMoNF: Experiments - MNIST



# Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions**

# Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]

# Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology

# Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology
  - Investigate the effect of a consistency loss regularization term

# Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology
  - Investigate the effect of a consistency loss regularization term
- Weight-sharing between components



# Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology
  - Investigate the effect of a consistency loss regularization term
- Weight-sharing between components
- Balance between complexities

# Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology
  - Investigate the effect of a consistency loss regularization term
- Weight-sharing between components
- Balance between complexities
- (Controlled) component annihilation

Thank you!