

Variational Mixture of Normalizing Flows

Guilherme Grijó Pen Freitas Pires

November 15, 2019

Thesis to obtain the Master of Science degree in
Electrical and Computer Engineering



Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions

Introduction and Motivation

- Deep generative models have been an active area of research, with promising results.

Introduction and Motivation

- Deep generative models have been an active area of research, with promising results.
 - **Implicit distributions:** Generative adversarial networks [Goodfellow et al., 2014], Variational Autoencoder [Kingma and Welling, 2014]
 - Don't allow explicit access to the density function

Introduction and Motivation

- Deep generative models have been an active area of research, with promising results.
 - **Implicit distributions:** Generative adversarial networks [Goodfellow et al., 2014], Variational Autoencoder [Kingma and Welling, 2014]
 - Don't allow explicit access to the density function
 - **Explicit distributions:** Normalizing flows [Rezende and Mohamed, 2015]
 - Allow explicit access to the density function
 - Lack an approach to introduce discrete structure (multi-modality) in the modelled distribution.

Introduction and Motivation: Goal

- The goal of this work was the development of a mixture of flexible distributions.

Introduction and Motivation: Goal

- The goal of this work was the development of a mixture of flexible distributions.
- This requires answering two questions:

Introduction and Motivation: Goal

- The goal of this work was the development of a mixture of flexible distributions.
- This requires answering two questions:
 - What should be the “family” of the mixture components?

Introduction and Motivation: Goal

- The goal of this work was the development of a mixture of flexible distributions.
- This requires answering two questions:
 - What should be the “family” of the mixture components?
 - How should the mixture components’ parameters be estimated?

- Mixture Models

Outline

- Mixture Models
- Variational Inference

Outline

- Mixture Models
- Variational Inference
 - The chosen framework for estimating the parameters of the proposed model

Outline

- Mixture Models
- Variational Inference
 - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows

- Mixture Models
- Variational Inference
 - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows
 - The chosen family for the mixture model components

- Mixture Models
- Variational Inference
 - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows
 - The chosen family for the mixture model components
- Variational Mixture of Normalizing Flows

- Mixture Models
- Variational Inference
 - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows
 - The chosen family for the mixture model components
- Variational Mixture of Normalizing Flows
- Experiments and results

- Mixture Models
- Variational Inference
 - The chosen framework for estimating the parameters of the proposed model
- Normalizing Flows
 - The chosen family for the mixture model components
- Variational Mixture of Normalizing Flows
- Experiments and results
- Conclusions and future work

Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models**
- 3 Normalizing Flows
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions

Mixture Models: Definition

- A mixture model is used to model data that is assumed to contain subgroups.

Mixture Models: Definition

- A mixture model is used to model data that is assumed to contain subgroups.
- Typically, it is assumed that the “subgroup-conditional” distributions belong to the same family, but have different parameters.


Mixture Models: Definition

- A mixture model is used to model data that is assumed to contain subgroups.
- Typically, it is assumed that the “subgroup-conditional” distributions belong to the same family, but have different parameters.
- Formally, a mixture model’s joint distribution (for a single instance x) is given by:

$$p(\mathbf{x}, c) = p(\mathbf{x}|c)p(c),$$

where c is the latent variable that indexes the subgroup to which x belongs

Mixture Models: Plate diagram



Mixture Models: Mixture of Gaussians




Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows**
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions

Normalizing Flows: Change of Variables

Given

$$\begin{cases} z \sim p(z) \\ x = g(z; \theta) \end{cases}$$

Normalizing Flows: Change of Variables

Given

$$\begin{cases} z \sim p(z) \\ x = g(z; \theta) \end{cases}$$

then:

$$\begin{aligned} f_X(\mathbf{x}) &= f_Z(g^{-1}(\mathbf{x}; \theta)) \left| \det \left(\frac{d}{d\mathbf{x}} g^{-1}(\mathbf{x}; \theta) \right) \right| \\ &= f_Z(g^{-1}(\mathbf{x}; \theta)) \left| \det \left(\frac{d}{d\mathbf{z}} g(\mathbf{z}; \theta) \right) \Big|_{\mathbf{z}=g^{-1}(\mathbf{x}; \theta)} \right|^{-1} \end{aligned}$$

Normalizing Flows: Change of Variables

Given

$$\begin{cases} z \sim p(z) \\ x = g(z; \theta) \end{cases}$$

then:

$$\begin{aligned} f_X(x) &= f_Z(g^{-1}(x; \theta)) \left| \det \left(\frac{d}{dx} g^{-1}(x; \theta) \right) \right| \\ &= f_Z(g^{-1}(x; \theta)) \left| \det \left(\frac{d}{dz} g(z; \theta) \right) \Big|_{z=g^{-1}(x; \theta)} \right|^{-1} \end{aligned}$$

This can be optimized w.r.t. θ , so as to approximate an arbitrary distribution

Normalizing Flows: Change of Variables

The described in the previous slide can be useful if

Normalizing Flows: Change of Variables

The described in the previous slide can be useful if

- The base density has a closed form expression and is easy to sample from

Normalizing Flows: Change of Variables

The described in the previous slide can be useful if

- The base density has a closed form expression and is easy to sample from
- The determinant of the Jacobian of g is computationally cheap - not the case, in general

Normalizing Flows: Change of Variables

The described in the previous slide can be useful if

- The base density has a closed form expression and is easy to sample from
- The determinant of the Jacobian of g is computationally cheap - not the case, in general
- The gradient of the determinant of the Jacobian of g w.r.t θ is computationally cheap

Normalizing Flows: Change of Variables

The framework of Normalizing Flows consists of composing several transformations that fulfill the three listed conditions.

Normalizing Flows: Change of Variables

The framework of Normalizing Flows consists of composing several transformations that fulfill the three listed conditions.

I.e., the function g is a composition of L functions h_ℓ ,
 $\ell = 0, 1, \dots, L - 1$.

Normalizing Flows: Change of Variables

The framework of Normalizing Flows consists of composing several transformations that fulfill the three listed conditions.

I.e., the function g is a composition of L functions h_ℓ , $\ell = 0, 1, \dots, L - 1$. Applying the formula to g , and taking the logarithm, yields:

$$\log f_X(\mathbf{x}) = \log f_Z(g^{-1}(\mathbf{x})) - \sum_{\ell=0}^{L-1} \log \left| \det \left(\frac{d}{d\mathbf{x}_\ell} h_\ell(\mathbf{x}_\ell) \right) \right|.$$

Normalizing Flows: Affine Coupling Layer

An example of such a transformation is the Affine Coupling Layer {[Dinh, Sohl-Dickstein, and Bengio, 2017]}.

Splitting z into (z_1, z_2) ,

Normalizing Flows: Affine Coupling Layer

An example of such a transformation is the Affine Coupling Layer {[Dinh, Sohl-Dickstein, and Bengio, 2017]}.

Splitting z into (z_1, z_2) ,

$$\begin{cases} x_1 &= z_1 \odot \exp(s(z_2)) + t(z_2) \\ x_2 &= z_2. \end{cases}$$

Normalizing Flows: Affine Coupling Layer

An example of such a transformation is the Affine Coupling Layer {[Dinh, Sohl-Dickstein, and Bengio, 2017]}.

Splitting z into (z_1, z_2) ,

$$\begin{cases} x_1 &= z_1 \odot \exp(s(z_2)) + t(z_2) \\ x_2 &= z_2. \end{cases}$$

This transformation has the following Jacobian matrix:

$$J_{f(z)} = \begin{bmatrix} \frac{\partial x_1}{\partial z_1} & \frac{\partial x_1}{\partial z_2} \\ \frac{\partial x_2}{\partial z_1} & \frac{\partial x_2}{\partial z_2} \end{bmatrix} = \begin{bmatrix} \text{diag}(\exp(s(z_2))) & \frac{\partial x_1}{\partial z_2} \\ \mathbf{0} & I \end{bmatrix}$$

Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows
- 4 Variational Inference**
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions

Variational Inference: Preamble

Consider a joint probability distribution $p(\boldsymbol{x}, \boldsymbol{z})$.

Variational Inference: Preamble

Consider a joint probability distribution $p(x, z)$. Suppose x is observed and z is latent.

Variational Inference: Preamble

Consider a joint probability distribution $p(\mathbf{x}, \mathbf{z})$. Suppose \mathbf{x} is observed and \mathbf{z} is latent. If we want to infer the most probable values of \mathbf{z} , given \mathbf{x} , by Bayes' Law:

$$\begin{aligned} p(\mathbf{z}|\mathbf{x}) &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{p(\mathbf{x})} \\ &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{\int p(\mathbf{x}|\mathbf{z}')p(\mathbf{z}')d\mathbf{z}'} \end{aligned}$$

Variational Inference: Preamble

Consider a joint probability distribution $p(\mathbf{x}, \mathbf{z})$. Suppose \mathbf{x} is observed and \mathbf{z} is latent. If we want to infer the most probable values of \mathbf{z} , given \mathbf{x} , by Bayes' Law:

$$\begin{aligned} p(\mathbf{z}|\mathbf{x}) &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{p(\mathbf{x})} \\ &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{\int p(\mathbf{x}|\mathbf{z}')p(\mathbf{z}')d\mathbf{z}'} \end{aligned}$$

Problem: The integral in the denominator is intractable for most interesting models.

Variational Inference: Preamble

Consider a joint probability distribution $p(\mathbf{x}, \mathbf{z})$. Suppose \mathbf{x} is observed and \mathbf{z} is latent. If we want to infer the most probable values of \mathbf{z} , given \mathbf{x} , by Bayes' Law:

$$\begin{aligned} p(\mathbf{z}|\mathbf{x}) &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{p(\mathbf{x})} \\ &= \frac{p(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{\int p(\mathbf{x}|\mathbf{z}')p(\mathbf{z}')d\mathbf{z}'} \end{aligned}$$

Problem: The integral in the denominator is intractable for most interesting models.

- Variational inference is an approximate inference framework, that can be used to overcome this intractability.

Variational Inference: Goal

Given a parametric family of distributions $q(z; \lambda)$, find the parameters λ that minimize the Kullback-Leibler divergence between $q(z; \lambda)$ and $p(z|x)$

Variational Inference: ELBO

$$KL(q||p) = \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z}$$

Variational Inference: ELBO

$$\begin{aligned}KL(q||p) &= \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} \\ &= \int q(\mathbf{z}) (\log q(\mathbf{z}) - \log p(\mathbf{z}|\mathbf{x})) d\mathbf{z}\end{aligned}$$

Variational Inference: ELBO

$$\begin{aligned}KL(q||p) &= \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - \log p(\mathbf{z}|\mathbf{x})) d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - (\log p(\mathbf{x}, \mathbf{z}) - \log p(\mathbf{x}))) d\mathbf{z}\end{aligned}$$

Variational Inference: ELBO

$$\begin{aligned}KL(q||p) &= \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - \log p(\mathbf{z}|\mathbf{x})) d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - (\log p(\mathbf{x}, \mathbf{z}) - \log p(\mathbf{x}))) d\mathbf{z} \\&= \mathbb{E}_q[\log q(\mathbf{z})] - \mathbb{E}_q[\log p(\mathbf{x}, \mathbf{z})] + \log p(\mathbf{x})\end{aligned}$$

Variational Inference: ELBO

$$\begin{aligned}KL(q||p) &= \int q(\mathbf{z}) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - \log p(\mathbf{z}|\mathbf{x})) d\mathbf{z} \\&= \int q(\mathbf{z}) (\log q(\mathbf{z}) - (\log p(\mathbf{x}, \mathbf{z}) - \log p(\mathbf{x}))) d\mathbf{z} \\&= \mathbb{E}_q[\log q(\mathbf{z})] - \mathbb{E}_q[\log p(\mathbf{x}, \mathbf{z})] + \log p(\mathbf{x})\end{aligned}$$

Which yields the lower bound (ELBO):

$$\begin{aligned}ELBO(q) &= \mathbb{E}_q[\log p(\mathbf{x}, \mathbf{z})] - \mathbb{E}_q[\log q(\mathbf{z})] \\&= \mathbb{E}_q[\log p(\mathbf{x}|\mathbf{z})] + \mathbb{E}_q[\log p(\mathbf{z})] - \mathbb{E}_q[\log q(\mathbf{z})]\end{aligned}$$

Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows**
- 6 Conclusions

Is it possible to combine the ideas from the previous sections, to obtain a mixture of flexible models?

Recall the ELBO:

$$ELBO(q) = \mathbb{E}_q[\log p(\mathbf{x}|\mathbf{z})] + \mathbb{E}_q[\log p(\mathbf{z})] - \mathbb{E}_q[\log q(\mathbf{z})]$$

VMoNF: Definition

Recall the ELBO:

$$ELBO(q) = \mathbb{E}_q[\log p(\mathbf{x}|\mathbf{z})] + \mathbb{E}_q[\log p(\mathbf{z})] - \mathbb{E}_q[\log q(\mathbf{z})]$$

Let the variational posterior $q(\mathbf{z}|\mathbf{x})$ be parameterized by a neural network.

VMoNF: Definition


Recall the ELBO:

$$ELBO(q) = \mathbb{E}_q[\log p(\mathbf{x}|\mathbf{z})] + \mathbb{E}_q[\log p(\mathbf{z})] - \mathbb{E}_q[\log q(\mathbf{z})]$$


Let the variational posterior $q(\mathbf{z}|\mathbf{x})$ be parameterized by a neural network.

We optimize this objective, by **jointly** learning the variational posterior and the generative components.



VMoNF: Overview



VMoNF: Experiments - Pinwheel (5 wings)




VMoNF: Experiments - Pinwheel (5 wings)





VMoNF: Experiments - Pinwheel (3 wings)

Training Animation


VMoNF: Experiments - 2 Circles





VMoNF: Experiments - 2 Circles



VMoNF: Experiments - 2 Circles (semi supervised)



VMoNF: Experiments - 2 Circles (semi supervised)



Note: 32 labeled points, 1024 unlabeled points

VMoNF: Experiments - MNIST




Table of Contents

- 1 Introduction and Motivation
- 2 Mixture Models
- 3 Normalizing Flows
- 4 Variational Inference
- 5 Variational Mixture of Normalizing Flows
- 6 Conclusions**

Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]

Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles \rightarrow Topology

Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology
 - Investigate the effect of a consistency loss regularization term

Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology
 - Investigate the effect of a consistency loss regularization term
- Weight-sharing between components

Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology
 - Investigate the effect of a consistency loss regularization term
- Weight-sharing between components
- Balance between complexities

Conclusions and Future Work

- Similar work is being pursued and published in prominent venues: [Dinh, Sohl-Dickstein, et al., 2019; Izmailov et al., 2019]
- Formally describe the reasons why the model fails in cases like the 2 circles → Topology
 - Investigate the effect of a consistency loss regularization term
- Weight-sharing between components
- Balance between complexities
- (Controlled) component annihilation

Thank you!