
Amik St-Cyr
**On Optimized Schwarz Preconditioning for High-Order
Spectral Element Methods**

National Center for Atmospheric Research
1850 Table Mesa Drive
Boulder
CO 80305.
`amik@ucar.edu`
Martin. J. Gander
Stephen J. Thomas

On Optimized Schwarz Preconditioning for High-Order Spectral Element Methods

A. St-Cyr¹, M.J. Gander² and S. J. Thomas³

February 2, 2004

¹**amik@ucar.edu**, NCAR 1850 Table Mesa Drive Boulder, CO 80305, USA.

²**mgander@math.mcgill.ca**, McGill 805 Sherbrooke W. Montreal QC, Canada
H3A2K6

³**thomas@ucar.edu**, NCAR 1850 Table Mesa Drive Boulder, CO 80305, USA.

Abstract

Optimized Schwarz preconditioning is applied to a spectral element method for the modified Helmholtz equation and pseudo-Laplacian arising in incompressible flow solvers. The preconditioning is performed on an element-by-element basis. The method enables one to use non-overlapping elements, yielding an effective algorithm in terms of communication between elements and implementation. Two approaches are tested. The first consists of constructing a P_1 finite element problem on each overlapping element. In the second, the preconditioner is applied directly on a non-overlapping spectral element. Numerical results demonstrate an improvement in the iteration count over the classical Schwarz algorithm.

Introduction

The classical Schwarz algorithm uses Dirichlet transmission conditions between sub-domains. By introducing a more general Robin boundary condition, it is possible to optimize the convergence characteristics of the original algorithm [1, 2, 5, 4]. In this work, a study of the model equations $u - \Delta u = f$ and pseudo-Laplacian arising in incompressible flow solvers is performed. As suggested by the work of [3], the preconditioning is either implemented via a P_1 finite element formulation of the original problem build on the spectral element grid, or directly by solving a smaller spectral element problem without overlap on each spectral element. Although traditional Schwarz preconditioning combined with a coarse grid solver is quite efficient, the need for even more powerful preconditioning techniques stems from atmospheric modeling. Recently (see [9, 7]), a semi-implicit SEM was combined with OIFS time stepping, enabling time steps on the order of 20 times the advective CFL condition [10]. This directly reflects as a significant increase in the number of conjugate gradient iterations required to perform the semi-implicit step.

Bibliography

- [1] P. CHARTON, F. NATAF AND F. ROGIER (1991), *Méthode de décomposition de domaines pour l'équation d'advection-diffusion*, C. R. Acad. Sci., Vol. 313, No. 9, pp. 623-626.
- [2] P. CHEVALIER AND F. NATAF (1998), *Symmetrized method with optimized second-order conditions for the Helmholtz equation*, In Domain decomposition methods, 10 (Boulder, CO, 1997), pp. 400-407, Amer. Math. Soc., Providence, RI.
- [3] P.F. FISCHER, N.I. MILLER AND H.M. TUFO (2000), *An overlapping Schwarz method for spectral element simulation of three-dimensional incompressible flows*, in Parallel Solution of Partial Differential Equations, P. Bjorstad and M. Lusk, eds., Springer-Verlag, pp.159-180.
- [4] M. J. GANDER, *Optimized Schwarz Methods (2003)*, Research Report, No. 2003-01, Dept. of Mathematics and Statistics, McGill University, 33 pages, submitted.
- [5] M.J. GANDER, F. MAGOULES AND F. NATAF (2002), *Optimized Schwarz Methods without Overlap for the Helmholtz Equation*, SIAM Journal on Scientific Computing, Vol. 24, No 1, pp. 38-60.
- [6] J.W. LOTTES AND P.F. FISCHER (2003), *Hybrid Multigrid/Schwarz Algorithms for the Spectral Element Method*, submitted.
- [7] A. ST-CYR AND S.J. THOMAS (2004), *Non-linear operator integration factor splitting for the shallow water equations*, In preparation for J. Sci. Comp.
- [8] S. J. THOMAS, J. M. DENNIS, H. M. TUFO, AND P. F. FISCHER (2003), *A Schwarz Preconditioner for the Cubed-Sphere*, SIAM J. Sci. Comp., Vol. 25, No. 2, pp. 442-453.
- [9] S.J. THOMAS AND R.D. LOFT (2002), *Semi-implicit spectral element atmospheric model*. Journal of Scientific Computing, vol 17, 339-350.
- [10] D. XIU AND G.E. KARNIADAKIS (2001), *A semi-Lagrangian high-order method for Navier-Stokes equations*, J.C.P., No. 172, pp. 658-684.