
DU Lei
**A block IDR(s) method for nonsymmetric linear
equations with multiple right-hand sides**

Department of Computational Science and Engineering
Graduate School of Engineering
Nagoya University
Furo-cho
Chikusa-ku
Nagoya 464-8603
Japan
lei-du@na.cse.nagoya-u.ac.jp
SOGABE Tomohiro
YU Bo
YAMAMOTO Yusaku
ZHANG Shao-Liang

The problem for solving several linear systems of equations with the same coefficient matrix and different right-hand sides, which can be written as

$$Ax_i = b_i, A \in R^{n \times n}, b_i \in R^n, i = 1, \dots, m \quad (1)$$

appears in many applications, such as in electromagnetic scattering and structural mechanics.

Many methods have been proposed to solve this kind of problem. One main class of these methods is the block solvers, including the block CG, block QMR, block GMRES, block Bi-CGSTAB, etc. Recently, Based on the induced dimension reduction theorem, the IDR(s) was developed by Sonneveld and Gijzen[1]. This new family of algorithms has some distinguishing features, for example, IDR(s) may converge the solution using at most $n+n/s$ matrix-vector products, IDR(1) is equivalent to Bi-CGSTAB at the even residuals and IDR(s) with $s \geq 1$ is competitive with most Bi-CG based methods. For these reasons, we proposed the block IDR(s): a block version of IDR(s) for solving linear systems of equation with multiple right-hand sides. Numerical experiments show the effectiveness of our proposed method.

Reference: [1]P. Sonneveld and M. van Gijzen, IDR(s): a family of simple and fast algorithms for solving large nonsymmetric systems of linear equations, SIAM J. Sci Comput., 31(2008), pp. 1035-1062.