

Kinematic Coherence and Rational Harmonic Phase-Locking in the Globular Cluster NGC 6205

Colt Camron Lien

University of Arkansas at Monticello, Monticello, Arkansas 71656, USA

July 27, 2026

Abstract

Standard models of globular cluster dynamics predict chaotic kinematic relaxation and random thermal variance among stellar populations. This study investigates isolated subsets of macro-vortices within NGC 6205 (M13) exhibiting anomalous structural rigidity and suppressed kinetic drag. Using astrometric and thermodynamic data from Gaia DR3, the central parsecs of M13 were filtered for stars demonstrating radial velocity variance of < 2.0 km/s and constant photometric output. K-Means clustering applied to the proper motion vectors (μ_α , μ_δ) reveals a highly coherent 40-node network partitioned into three distinct kinematic clusters. Centroid measurements indicate the transverse spatial distances between these clusters scale to rational integer resonances (4 : 3 and 16 : 15). The uniform astrometric stability (RUWE ≈ 1.0) and rational spatial distribution strongly suggest this subset operates not as independent gravitational bodies, but as a phase-locked, macro-cymatic standing wave within a continuous fluid plenum.

1 Introduction

Current gravitational relaxation models struggle to fully account for localized, rigid kinematic coupling within dense globular clusters without invoking hypothetical intermediate-mass black holes (IMBHs) [8]. In standard astrophysics, massive celestial bodies in dense clusters should experience chaotic thermal variance and kinetic relaxation [11].

However, when viewed through a continuous fluid-dynamic framework, stellar masses operate as localized pressure wells experiencing kinetic drag [1, 7]. Extrapolating from macroscopic hydrodynamic quantum analog experiments [4, 2], if subsets of stars achieve perfect laminar flow, establishing an infinite bulk modulus ($\lim_{\rho \rightarrow \rho_{max}} K = \infty$), they should vibrate as a unified macroscopic acoustic network. This paper proposes that a subset of stars within NGC 6205 has achieved this exact state of kinematic phase-locking.

2 Data and Methodology

Data was extracted from the European Space Agency’s Gaia Data Release 3 (DR3) [5]. The initial query targeted the spatial coordinates of M13 (RA = 250.421° , Dec = 36.460°) within a 0.1° radius.

2.1 Kinetic Filtration Pipeline

To isolate nodes operating within a localized standing wave, the background telemetry was filtered for extreme kinetic stability and continuous thermodynamic output:

- Kinetic Drag Exclusion:** Nodes with a radial velocity error > 2.0 km/s were removed to isolate laminar flow.
- Dynamic Pressure Continuity:** Nodes were filtered strictly for `phot_variable_flag = 'CONSTANT'` or `'NOT_AVAILABLE'` to ensure unshifting kinetic pressure waves.
- Outlier Rejection:** Stars experiencing kinetic shear (proper motion deviations exceeding two standard deviations from the mean) were dropped, isolating bound kinematic structures [6], yielding a final rigid network of 40 nodes.

3 Transverse Kinematic Clustering

A 3D spatial-kinetic mapping of the 40-node network reveals the stars rest on a remarkably flat uniform radial velocity plane along the Z-axis, indicating shared localized dynamic pressure.




Figure 1: 3D spatial-kinetic mapping of the M13 network (edge-on perspective). The isolated nodes demonstrate extreme alignment along the Z-axis (Kinetic Velocity), indicating a shared macroscopic fluid pressure plane.

The proper motion density contour (Figure 2) displays a distinct tri-harmonic distribution. Crucially, the central nodes exhibit uniform Renormalised Unit Weight Error ($\text{RUWE} \approx 1.0$) [6]. In standard astrometry, this value indicates a distinct absence of chaotic, unresolved perturbations. This uniform kinematic stability across spatially distributed stars contradicts expectations of random thermal noise, pointing instead to a highly coherent macroscopic structure.

4 Rational Harmonic Scaling

To map the acoustic anchors of the network, a K-Means clustering algorithm ($n = 3$) [9] was applied to the proper motion vectors.

4.1 Centroid Measurements

The spatial coordinates (μ_α, μ_δ) for the three distinct macroscopic harmonic clusters were isolated, and the Euclidean distances between them were calculated:

- Transverse Distance A: 0.5048 mas/yr
- Transverse Distance B: 0.4728 mas/yr
- Transverse Distance C: 0.3544 mas/yr

4.2 Acoustic Integer Resonance

Analyzing the mathematical ratios of these transverse distances reveals a shift from irrational KAM stability metrics [3] to distinct rational integer fractions:

$$\text{Ratio A/B} = \frac{0.5048}{0.4728} = 1.0677 \quad (1)$$




Figure 2: KDE Plot of Transverse Kinetic Coupling in the M13 Acoustic Network. Note the three distinct harmonic clusters and the cyan coherence ($\text{RUWE} \approx 1.0$).

This mathematically aligns with a 16 : 15 (≈ 1.066) higher-order kinetic boundary state.

$$\text{Ratio B/C} = \frac{0.4728}{0.3544} = 1.3342 \quad (2)$$

This aligns with a 4 : 3 (≈ 1.333) first-order harmonic resonance. In fluid dynamics, wave interference dictates that nodes must entrain to low-integer ratios to achieve continuous constructive interference, minimizing localized fluid drag [7, 10]. The alignment with these specific rational geometric ratios is consistent with the preservation of a highly stable, localized macro-cymatic standing wave.

5 Discussion

The primary conduit node anchoring Cluster 1 exhibits a K-type thermodynamic output of 4284 K. This temperature provides the highly stable, continuous dynamic pressure gradient required to sustain the localized network without the violent flaring common in M-type stars.

The slight deviation from the irrational Golden Ratio ($\phi \approx 1.618$) expected in pure KAM stability [3] suggests the network is in a transient thermodynamic state. The shift to rational integer fractions, specifically the highly stable 4:3 first-order resonance, may represent a structural recalibration of tension designed to minimize fluid shear against the external localized pressure of the dense cluster environment.

6 Conclusion

Empirical astrometric data from Gaia DR3 highlights the existence of a highly coherent, 40-star

kinematic network embedded within NGC 6205. The uniform kinematic coupling and precise rational spacing present a significant challenge to standard gravitational relaxation models. Future observational cosmology parameters must shift from localized point-source emission searches to measuring instantaneous phase-shifts and geometric rigidity across macroscopic stellar subsets.

Code and Data Availability

All Python extraction scripts, K-Means clustering algorithms, and the filtered Gaia DR3 datasets utilized in this research are available in a public GitHub repository to allow for complete methodological transparency and independent peer verification.

AI Disclosure

The author acknowledges the use of an AI language model to assist with the structural organization, academic formatting, and stylistic refinement of this manuscript. All theoretical concepts, mechanical architecture, and scientific conclusions are the original intellectual property of the author.

References

- [1] L. Berezhiani and J. Khoury, “Dark Matter Superfluidity and Galactic Dynamics,” *Phys. Rev. D* **92**, 103510 (2015).
- [2] J. W. M. Bush, “Pilot-Wave Hydrodynamics,” *Annu. Rev. Fluid Mech.* **47**, 269-292 (2015).
- [3] A. Celletti and L. Chierchia, “KAM Stability and Celestial Mechanics,” *Mem. Am. Math. Soc.* **187**, 878 (2007).
- [4] Y. Couder and E. Fort, “Single-Particle Diffraction and Interference at a Macroscopic Scale,” *Phys. Rev. Lett.* **97**, 154101 (2006).
- [5] Gaia Collaboration et al., “Gaia Data Release 3. Summary of the content and survey properties,” *Astron. Astrophys.* **674**, A1 (2023).
- [6] P. Kervella, F. Arenou, F. Mignard, and F. Thévenin, “Stellar and substellar companions from Gaia EDR3: Proper motion anomaly and resolved common proper motion pairs,” *Astron. Astrophys.* **657**, A7 (2022).
- [7] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics*, 2nd ed. (Pergamon Press, Oxford, 1987).
- [8] J. M. Lotz et al., “Globular cluster distributions as a dynamical probe of dark matter,” arXiv preprint arXiv:2504.02476 (2025).
- [9] J. MacQueen, “Some methods for classification and analysis of multivariate observations,” *Proc. Fifth Berkeley Symp. Math. Statist. Prob.* **1**, 281-297 (1967).
- [10] A. Pikovsky, M. Rosenblum, and J. Kurths, *Synchronization: A Universal Concept in Nonlinear Sciences* (Cambridge University Press, Cambridge, 2001).
- [11] E. Vasiliev and H. Baumgardt, “Gaia EDR3 view on globular cluster kinematics and orbit anisotropy,” *Mon. Not. R. Astron. Soc.* **505**, 5978 (2021).