


10.2 Counting Outcomes – Overview

Definitions / Key Ideas

Fundamental Counting Principle: If a job consists of n separate tasks, the first of which can be done k_1 ways, the second k_2 ways and so on, then the total job can be done $k_1 \times k_2 \times \cdots \times k_n$ ways.

Task 1	Task 2	...	Task n	Total Outcomes
k_1	k_2	...	k_n	$k_1 \times k_2 \times \cdots \times k_n$



Example: Flip a coin and roll a 4-sided die.
How many total outcomes are there?

Example: Sally has 6 pairs of socks, 4 shorts, 5 shirts, and 3 sunglasses. How many ways can she get dressed?

$$\frac{6}{\text{socks}} \times \frac{4}{\text{shorts}} \times \frac{5}{\text{shirts}} \times \frac{3}{\text{sunglasses}} = 360 \text{ total ways}$$

With or without replacement: We need to take into account whether or not objects can be repeated in our calculations.

Examples: a) How many passwords can you make if it requires 4 digits? *with replacement*

$$\frac{10}{\text{digit 1}} \times \frac{10}{\text{digit 2}} \times \frac{10}{\text{digit 3}} \times \frac{10}{\text{digit 4}} = 10^4 = 10,000 \text{ passwords}$$

b) How many passwords can you make if it requires 4 digits, but you cannot repeat digits? *without replacement*

$$\frac{10}{\text{digit 1}} \times \frac{9}{\text{digit 2}} \times \frac{8}{\text{digit 3}} \times \frac{7}{\text{digit 4}} = 5040 \text{ passwords}$$

Factorials: $n!$ (read "n factorial") is the product of all numbers less than and including n

$$n! = n(n-1)(n-2)\cdots(3)(2)(1)$$

Example: For the 9 starting players on a baseball team, how many different batting orders are there?

$$\frac{9}{\text{player 1}} \times \frac{8}{\text{player 2}} \times \frac{7}{\text{player 3}} \times \cdots \times \frac{1}{\text{player 9}} = 9! = 362,880$$

without replacement

Combinations and Permutations

- We often want to be able to count the number of ways that we can choose members from a group of objects (without replacement).

- "Selecting r objects from a total of n objects".
 $r \leq n$



- There are two methods to count in these scenarios, the only difference is if order matters or order doesn't matter.

Permutations: Order matters. When you are selected is important. Your position has meaning. ★

$${}_n P_r = P\left(\begin{matrix} n \\ r \end{matrix}\right) = P(n, r) = \frac{n!}{(n-r)!}$$

Total # how many selecting

Combinations: Order does not matter. It does not matter when you are selected, only if you were selected.

$${}_n C_r = \left(\begin{matrix} n \\ r \end{matrix}\right) = C(n, r) = \frac{n!}{r!(n-r)!}$$

Examples: Decide if we should use permutations or combinations to count the total number of outcomes (possible ways to select our group). Then count the number of outcomes.

- a) There are 8 runners in a race. How many ways can they place 1st, 2nd, and 3rd?

Permutation → order matters

$$8 P_3 = 336$$

meaning to each slot → 1st 2nd 3rd

- b) Out of 12 students, how many ways can we select a committee of four students?

Combinations → order doesn't matter

$$12 C_4 = 495$$

(only interested in selected vs not selected)

- c) We are forming a committee and we need to select a president, vice president and secretary. How many ways can this be done if there are 10 members?

Permutation → order matters

$$10 P_3 = 720$$

Matters if → Pres VP Sec
Pres vs VP vs Sec

Permutations with Repeated Objects: Counting the number of distinct ways we can arrange all n objects when some of the objects are the same (repeated, specifically k_1 are alike, k_2 are alike, and so on).

$$\frac{n!}{(k_1!)(k_2!) \dots (k_p!)}, \quad \text{where } k_1 + k_2 + \dots + k_p = n$$

Example: Harmony was born on 05/19/1991. How many eight-digit codes could she make using the digits in her birthday?

8 total digits

9	→	3 times
1	→	3 times
5	→	1
0	→	1

$$\frac{8!}{3!3!1!1!} = 1,120$$