

# Galaxy Evolution Through Tidal Interactions: Evolution of the Observed and Mass Derived Rotation Curves of M31 and the Milky Way

COLTON QUIRK<sup>1</sup>

<sup>1</sup> *University of Arizona*

## 1. INTRODUCTION

The study of galaxy evolution is crucial to understanding the large-scale structure of the universe and the mechanisms that drive changes in galactic morphology and kinematics. One key aspect of galaxy evolution is the role of tidal interactions in shaping the observed and mass-derived rotation curves of galaxies. This project focuses on the evolution of the rotation curves of the Milky Way and M31 (the Andromeda Galaxy), particularly how their disk and bulge components respond to gravitational interactions over time. Tidal interactions are the gravitational forces which arise from interactions between spatially extended objects, not point-sources (E. R. Dimitri Gadotti 2016). As the two galaxies move towards each other the parts of the galaxy closest to each other will feel a different amount of gravitational force than the parts of the galaxy farther from each other. This will cause the structure of the galaxies to change. This project will study how the galaxies change by analyzing the rotation curves of each galaxy, for both the disk stars and the bulge stars. Rotation curves provide insight into the mass distribution of galaxies, distinguishing between visible and dark matter contributions. By examining how these curves evolve through tidal interactions, we can gain a clearer picture of the dynamical history of these galaxies and their future trajectories.

Understanding the evolution of rotation curves is fundamental to deciphering how galaxies grow, interact, and redistribute their mass. The discrepancy between observed and mass-derived rotation curves has been a longstanding issue in astrophysics, offering evidence for dark matter or alternative gravitational theories. Studying these curves in the Milky Way and M31 allows us to explore how tidal interactions—such as satellite mergers, close encounters, and long-range gravitational influences—reshape the internal dynamics of galaxies (P. L. T. Dusan Keres 2007). Since both the Milky Way and M31 are on a collision course, analyzing their past and present rotation curves provides a valuable window into how interactions impact galactic structure, angu-

lar momentum distribution, and the overall dark matter profile.

Our current understanding of galactic rotation curves stems from both observational and theoretical studies. Observations of spiral galaxies indicate that their rotation curves remain nearly flat at large radii, implying the presence of extended dark matter halos (E. R. Dimitri Gadotti 2016). Theoretical models and N-body simulations suggest that tidal interactions can redistribute mass within a galaxy, potentially altering both the baryonic and dark matter components (R. G. T. S. Jay Anderson 2012). In particular, major interactions, such as mergers or flybys, can lead to angular momentum transfer, bulge growth, and changes in the mass-to-light ratio, all of which impact the rotation curve.

Despite significant progress, many open questions remain regarding the evolution of rotation curves under tidal influences. How exactly do tidal interactions reshape the mass distribution of the disk and bulge over different timescales (R. A. F. B. W. Rachel S. Somerville 2015)? Do observed changes in rotation curves primarily reflect baryonic redistribution, or do they hint at modifications in the dark matter halo? Additionally, while we can model individual interactions, predicting long-term evolutionary trends remains challenging due to the complex interplay between baryons, dark matter, and external forces. Finally, as the Milky Way and M31 move toward their eventual merger, how will their respective rotation curves evolve, and what can this tell us about the fate of similar galactic interactions throughout cosmic history? Addressing these questions will enhance our understanding of galaxy evolution and the nature of dark matter.

## 2. PROPOSAL

### 2.1. Proposal

This project seeks to answer how the rotation curves of the Milky Way and the Andromeda galaxy change over time. For example, do the rotation curves maintain a flat profile throughout the merger? If they do not maintain a flat profile when and why does it deviate? As well this project will analyze how the disk and

bulge particles within the galaxies are affected by the merger process. By answering these questions we can gain significant insights into how mergers affect galaxy evolution.

## 2.2. Methods

This project will analyze the rotation curves of the Milky Way and the Andromeda Galaxy throughout their merger. We will use all 802 snapshots for both galaxies using the VLowRes data. For each snapshot we will construct four rotation curves. We will create rotation curves for both the disk type and bulge type particles, particle types 2.0 and 3.0 respectively. As well for each particle type we will construct two rotation curves, the mass-derived rotation curve, and the "observed" rotation curve.

The mass-derived rotation curves are calculated by taking radial slices of the galaxy and measuring the mass within that radial slice. Then we can calculate the velocity needed for a particle to have a circular orbit around that mass at that radial slice.

$$F = m\vec{a} = m\frac{v^2}{r} = \frac{GMm}{r^2} \quad (1)$$

We can solve this equation for the velocity of a circular orbit.

$$v_{\text{circ}} = \sqrt{\frac{GM}{r}} \quad (2)$$

This circular velocity would be calculated for each radial slice by measuring the amount of mass within that radial distance. See Figure 1 for an example of a mass derived rotation curve of M31.

The "observed" rotation curve is the line-of-sight velocity field when viewed edge-on. This will be constructed for each snapshot for each galaxy and particle type. This will be made using the same technique as used in Lab 7 of this course. That is we will take the galaxy and rotate it such that the angular momentum of the galaxy is aligned with the z-axis. Then we will plot the velocities of the particle versus their x-coordinate (relative to the center of mass of the galaxy) as a 2D

histogram. This will construct a rotation curve of the galaxy for a particular snapshot. See Figure 2 for an example of an "observed" rotation curve.

Thus the overall process for finding how the rotation curves change over time is shown below. This process will be completed for both galaxies.

1. Load in the data for a particular snapshot
2. Create two MassProfile objects, one for Disk Particles and the other for Bulge Particles
3. For each MassProfile object create a mass derived rotation curve using equation 2
4. For each MassProfile rotate the frame of the galaxy such that the angular momentum aligns with the z-axis
5. Then plot the x-coordinate versus the velocity of the particles as a 2d histogram
6. Save both of these profiles and continue to the next snapshot


These rotation curves will then be overplotted to show how the "observed" and mass-derived rotation curves align with each other. As well the rotation curves will then be added to an animation to show the evolution over time for both galaxies and particle types.

## 2.3. Hypothesis


We hypothesize that the rotation curves will not maintain a flat profile throughout the merger. As the tidal forces disrupt the rotation of the galaxies the rotation curves will be impacted. We also expect there to be some transfer between the rotation curves of the disc particles and bulge particles. As discussed in R. A. F. B. W. Rachel S. Somerville (2015) there is a transfer of particles between the disc and the bulge of galaxies during mergers. This will affect the distribution of the angular momentum of the galaxies during the merger, leading to variances in the rotation curves of the disc and bulge particles.

## REFERENCES

- Dimitri Gadotti, E. R. 2016, Galactic Bulges (Springer)
- Dusan Keres, P. L. T. 2007, The Astrophysical Journal
- Jay Anderson, R. G. T. S. 2012, The Astrophysical Journal
- Rachel S. Somerville, R. A. F. B. W. 2015, Monthly Notices of the Royal Astronomical Society



**Figure 1.** Mass-derived rotation curve for M31 at snap number 0. This was made for Homework 5.



**Figure 2.** "Observed" rotation curve for M31 at snap number 0. This was made for Lab 7.