

Notes: (Topic 3.6) Sinusoidal Function Transformations

Previously (in Topics 3.4 – 3.5), we learned how to identify the midline, amplitude, and period from the graph of a sinusoidal function. Now, we will learn how those properties affect the equations of sinusoidal functions.

To do this, let's consider how the midline, amplitude, and period affect the graph of a sinusoidal function through the lens of transformations that we learned in Unit 1.


The **midline** of a sinusoidal function is simply a **vertical translation**.

The **amplitude** of a sinusoidal function is the **vertical dilation**.

The **period** of a sinusoidal function is the result of a **horizontal dilation**.


Given a function written as $f(\theta) = a \sin(b\theta) + d$ or $k(\theta) = a \cos(b\theta) + d$, the graphs of f and k will have the following transformations:

1. A vertical dilation by a factor of a . (a represents the amplitude of the graph)
2. A vertical translation of d units. (d represents the midline of the graph)
3. A horizontal dilation by a factor of $\left|\frac{1}{b}\right|$. (The period is given by $P = \left|\frac{2\pi}{b}\right|$)




Graph of h

Example 1: The graph of the sinusoidal function h is shown in the figure above, along with information about the amplitude, midline, and period of the graph. The function h can be written as $h(\theta) = a \sin(b\theta) + d$. Find the values of the constants a , b , and d .



Graph of f

Example 2: The figure shows the graph of a trigonometric function f . Write an expression for $f(\theta)$.



Graph of g


Example 3: The figure shows the graph of a sinusoidal function g . What are the values of the period and amplitude of g ?

- (A) The period is π , and the amplitude is 2.
- (B) The period is π , and the amplitude is 4.
- (C) The period is 2π , and the amplitude is 2.
- (D) The period is 2π , and the amplitude is 4.

Example 4: The trigonometric function k has a maximum at the point $(0, 6)$. After this maximum, the next minimum occurs at the point $\left(\frac{\pi}{2}, -4\right)$. Which of the following could be an expression for $k(x)$?

- (A) $10\cos(\pi x) + 1$
- (B) $10\cos(2x) + 1$
- (C) $5\cos(\pi x) + 1$
- (D) $5\cos(2x) + 1$

Example 5: The trigonometric function f is given by $f(x) = 3\sin(2x) - 1$. Which of the following could be the graph of $f(x)$?




Phase Shift of a Sinusoidal Function

A horizontal translation of a sinusoidal function is called a **phase shift**.

The graph of $g(x) = \sin(x + c)$ is a **phase shift** of the graph of $f(x) = \sin(x)$ by $-c$ units.


The same results can be applied to the cosine function.



Graph of g

Example 6: The figure shows the graph of a trigonometric function g . Which of the following could be an expression for $g(x)$?


- (A) $3\cos(x) - 1$ (B) $-3\cos(x - \pi) - 1$ (C) $3\sin\left(x + \frac{\pi}{2}\right) - 1$ (D) $-3\sin\left(x - \frac{3\pi}{2}\right) - 1$



Graph of h

Example 7: The figure shows the graph of a trigonometric function h . Which of the following could be an expression for $h(x)$?

- (A) $2 \cos\left(2\left(x - \frac{\pi}{4}\right)\right) + 1$
- (B) $2 \cos(2(x - \pi)) + 1$
- (C) $2 \sin\left(2\left(x - \frac{\pi}{4}\right)\right) + 1$
- (D) $2 \sin(2(x - \pi)) + 1$



Example 8: The graph of h and its dashed midline for two full cycles is shown. Five points, F , G , J , K , and P are labeled on the graph. No scale is indicated, and no axes are presented.

The coordinates for the five points: F , G , J , K , and P are: $F(2, 16)$, $G(5, 11)$, $J(8, 6)$, $K(11, 11)$, $P(14, 16)$.

The function h can be written in the form $h(t) = a \sin(b(t + c)) + d$. Find values of constants a , b , c , and d .