

2025 Combinatorics Workshop

2025 조합론 학술대회

Monday, 18 August 2025 – Wednesday, 20 August 2025

Auditorium on the 2nd floor
IBS Science Culture Center
(IBS 과학문화센터)

55 Expo-ro, Daejeon, South Korea

<https://cw2025.combinatorics.kr/>



The electronic version of this booklet can be found at:
<https://cw2025.combinatorics.kr/>

The open-source \LaTeX template, `AMCOS_booklet`, used to generate this booklet is available
at https://github.com/maximelucas/AMCOS_booklet

Contents

About	5
History	5
Organizing committee	7
Advisory committee	7
Timetable	9
Monday, August 18, 2025	9
Tuesday, August 19, 2025	10
Wednesday, August 20, 2025	11
List of Abstracts	13
Monday 14:00-15:00	13
Monday 15:00-15:30	14
Monday 16:00-18:00	15
Tuesday 09:30-10:30	17
Tuesday 11:00-12:00	17
Tuesday 13:30-14:30	19
Tuesday 14:30-15:30	19
Tuesday 16:00-16:30	20
Tuesday 16:30-18:00	21
Wednesday 09:30-10:30	22
Wednesday 11:00-12:30	22
List of Participants	25
Useful Information	29
How to get to the IBS?	29
Accommodation	29
Lunch on Tuesday	29
Sponsors	31

Combinatorics Workshop (조합론 학술대회) is an annual conference for researchers in combinatorics and related areas in Korea. It was started in 2004 by the Yonsei University BK21 Research Group. Since 2013, this workshop has been advised by the committee of discrete mathematics of the Korean Mathematical Society. This year, it will occur at the Institute for Basic Science (IBS) in Daejeon from August 18 to August 20, 2025.

History

1. 2004 Combinatorics Workshop (August 16-17, 2004)
 - Venue: Yonsei University, Seoul
 - Host: Yonsei University BK21 Research Group
 - (Local) Organizers: Seung Kyung Park, Sang-Gu Lee, Sangwook Ree
2. 2005 Combinatorial Mathematics Conference (August 8-10, 2005)
 - Venue: Yonsei University, Seoul
 - Host: Yonsei University BK21 Research Group
 - (Local) Organizers: Dongsu Kim, Seung Kyung Park, Sang-Gu Lee, Jaeun Lee
3. 2006 Combinatorics Workshop (August 9-10, 2006)
 - Venue: Yeongnam University, Gyeongsan
 - Host: Department of Mathematics, Yeongnam University
 - (Local) Organizers: Jaeun Lee, Young Soo Kwon
4. 2007 Combinatorics Workshop (August 6-8, 2007)
 - Venue: KAIST, Daejeon
 - Host: KAIST Combinatorics Lab
 - (Local) Organizers: Heesung Shin, Jang Soo Kim
5. 2008 Workshop on Combinatorics (August 7-8, 2008)
 - Venue: Sungkyunkwan University, Suwon
 - Host: Sungkyunkwan University BK21 Division
 - (Local) Organizers: Gi-Sang Cheon, Sang-Gu Lee
6. 2009 Combinatorics Workshop (August 20-21, 2009)
 - Venue: KAIST, Daejeon
 - Host: KAIST ASARC
 - (Local) Organizers: Soon-Yi Kang, Sang-il Oum
7. 2010 Combinatorics Workshop (August 20-21, 2010)
 - Venue: Yeongnam University, Gyeongsan
 - Host: Department of Mathematics, Yeongnam University
 - (Local) Organizers: Jaeun Lee, Young Soo Kwon

8. 2011 Combinatorics Workshop (August 17-19, 2011)
 - Venue: Kangwon National University, Chuncheon
 - Host: Department of Mathematics Education, Kangwon National University
 - (Local) Organizers: Seunghyun Seo, Heesung Shin
9. 2012 Combinatorics Workshop (August 9-10, 2012)
 - Venue: Chonnam National University, Gwangju
 - Host: Department of Mathematics, Chonnam National University
 - (Local) Organizers: Hyeong-Kwan Ju, Sangwook Kim
10. 2013 Combinatorics Workshop (August 7-9, 2013)
 - Venue: CAMP (Center for Applications of Mathematical Principles), NIMS, Daejeon
 - Host: NIMS (National Institute for Mathematical Sciences)
 - (Local) Organizers: Dongsu Kim, Boram Park, Seunghyun Seo, Heesung Shin, Ae Ja Yee
11. 2014 Combinatorics Workshop (October 31-November 1, 2014)
 - Venue: Ajou University, Suwon
 - Host: Department of Mathematics, Ajou University
 - (Local) Organizers: Soojin Cho, Suyoung Choi, Boram Park
12. 2015 Combinatorics Workshop (July 13-16, 2015)
 - Venue: CAMP (Center for Applications of Mathematical Principles), NIMS, Daejeon
 - Host: NIMS (National Institute for Mathematical Sciences)
 - (Local) Organizers: JiYoon Jung, Seunghyun Seo, Heesung Shin, Jiang Zeng
13. 2016 Combinatorics Workshop (July 21-23, 2016)
 - Venue: KAIST, Daejeon
 - Host: KIAS CMC
 - (Local) Organizers: Soojin Cho, Dongsu Kim, Young Soo Kwon, Sang-il Oum
14. 2017 Combinatorics Workshop (August 16-17, 2017)
 - Venue: Sungkyunkwan University, Suwon
 - Host: Applied Algebra and Optimization Research Center (AORC), Sungkyunkwan University
 - (Local) Organizers: Gi-Sang Cheon, Jang Soo Kim, Young Soo Kwon, Heesung Shin
15. 2018 Combinatorics Workshop (August 17-18, 2018)
 - Venue: Seoul National University, Seoul
 - Host: IMDARC, Seoul National University
 - (Local) Organizers: Woong Kook, Seung Jin Lee
16. 2019 Combinatorics Workshop (August 13-15, 2019)
 - Venue: Hotel Skypark Songdo, Incheon
 - Host: Combinatorics Incheon and IBS Discrete Mathematics Group
 - (Local) Organizers: O-joung Kwon, Suil O, Sang-il Oum, Heesung Shin
17. 2020 Combinatorics Workshop (August 24, 2020)
 - Venue: Kyung Hee University, Seoul (online)
 - Host: Kyung Hee University and IBS Discrete Mathematics Group
 - (Local) Organizers: Jeong Han Kim, Sang June Lee, Sang-il Oum, Heesung Shin
18. 2021 Combinatorics Workshop (December 20-22, 2021)
 - Venue: The Bloomvista, Yangpyeong
 - Host: IBS Discrete Mathematics Group
 - (Local) Organizers: Jeong-Ok Choi, Sang-il Oum, Heesung Shin

19. 2022 Combinatorics Workshop (September 30-October 1, 2022)
 - Venue: GIST, Gwangju
 - Host: GIST and IBS Extremal Combinatorics and Probability Group
 - (Local) Organizers: Jeong-Ok Choi, Minki Kim, Sangwook Kim, Hong Liu
20. 2023 Combinatorics Workshop (August 16-18, 2023)
 - Venue: Yonsei University, Seoul
 - Host: Yonsei University
 - (Local) Organizers: JiSun Huh, Sangwook Kim, Joonkyung Lee, Jaebum Sohn
21. 2024 Combinatorics Workshop (August 19-21, 2024)
 - Venue: Chungbuk National University, Cheongju
 - Host: Chungbuk National University and IBS Discrete Mathematics Group
 - (Local) Organizers: Doowon Koh, Eunjeong Lee, Sang-il Oum, Meesue Yoo

Webpages: cwYYYY.combinatorics.kr for each year YYYY.

Organizing committee

- Sang-il Oum (엄상일), IBS Discrete Mathematics Group
- Jang Soo Kim (김장수), Sungkyunkwan University
- Seunghyun Seo (서승현), Kangwon National University

Advisory committee

- Committee of Discrete Mathematics, The Korean Mathematical Society
(Chair: Sang-il Oum)

Timetable

IS: Invited Speaker, CT: Contributed Talk

Monday, August 18, 2025

13:30–13:50	Registration		
13:50–14:00	Welcome remarks		
14:00–15:00	IS	Eun Jung Kim 김은정 KAIST & IBS DIMAG & CNRS	Algorithm and MSO logic on graphs with tree-like structure
15:00–15:30	CT	Yongho Shin 신용호 University of Wrocław	Learning-augmented online bipartite fractional matching
15:30–16:00	Coffee		
16:00–16:30	CT	Dabeen Lee 이다빈 KAIST	Combinatorial optimization through the lens of boolean polynomials and binary matroids
16:30–17:00	CT	Seonghyuk Im 임성혁 KAIST & IBS ECOPRO	Random perturbation of dense graphs
17:00–17:30	CT	Hyunwoo Lee 이현우 KAIST & IBS ECOPRO	On a Ramsey–Turán variant of Roth’s theorem
17:30–18:00	CT	Ingyu Baek 백인규 Yonsei University	On attainable homomorphism domination exponents
18:30–20:00	Dinner		

Tuesday, August 19, 2025

09:30–10:30	IS	Dongsu Kim 김동수 KAIST	Combinatorics of orthogonal polynomials in the q -Askey scheme
10:30–11:00	Coffee		
11:00–11:30	CT	Heesung Shin 신희성 Inha University	Combinatorial Bijections in Permutations, Trees, and Paths
11:30–12:00	CT	Younghan Yoon 윤영한 Ajou University	On the graph a -numbers
12:00–13:30	Lunch break		
13:30–14:30	IS	Ae Ja Yee 이애자 Pennsylvania State University	Partition statistics and the Littlewood decomposition
14:30–15:00	CT	Inseo Kim 김인서 Chungbuk National University	On toric Schubert varieties in flag varieties
15:00–15:30	CT	Seonkyung Kim 김선경 Kangwon National University	The largest size of an $(s, s + 2)$ -core partition with even or odd parts only
15:30–16:00	Coffee		
16:00–17:00	IS	Zhicong Lin Shandong University	Tree structures and combinatorics of bi- γ positivity
17:00–17:30	CT	Seunghun Lee 이승훈 IBS DIMAG	On the extension problem on the moment curve
17:30–18:00	CT	Seonghyeon Yu 유성현 Ajou University	Bier spheres and their full subcomplexes
18:00	Conference Banquet at IBS		

Wednesday, August 20, 2025

09:30–10:30	IS	O-joung Kwon 권오정 Hanyang University & IBS DIMAG	On problems related to digraph width parameters
10:30–11:00	Coffee		
11:00–11:30	CT	Myounghwan Lee 이명환 Hanyang University	Unavoidable butterfly minors in digraphs of large cycle rank
11:30–12:00	CT	Seokbeom Kim 김석범 KAIST & IBS DIMAG	Subtournaments of tournaments of large clique-width
12:00–12:30	CT	Seog-Jin Kim 김석진 Konkuk University	The square of every subcubic planar graph without 4-cycles and 5-cycles is 7-choosable
12:30–13:30	Pizza Lunch		

Monday 14:00-15:00

Algorithm and MSO logic on graphs with tree-like structure

Eun Jung Kim

IS

¹ KAIST

² IBS Discrete Mathematics Group

³ CNRS

A finite-state automaton is a computing machine (“algorithm”) of primitive form, which scans a given string over a finite alphabet, e.g. a string consisting of 0’s and 1’s, and decides if the given string is in some language. It is primitive in the sense that the machine does not have a memory, and the machine keeps track of the inner working of the algorithm by having a “state”, out of a finite number of possible states, which is updated according to a hard-wired rule as the machine reads the symbols on the input string one by one. A set of strings (“language”) recognized by a finite automaton is known as a regular language. It is well-known that a regular language is precisely described by a homomorphism from the set of all strings to a finite semigroup. A less known but nonetheless important is that a language is regular if and only if it is defined by so called Monadic Second Order Logic (Büchi-Elgot-Trakhtenbrot 1960), MSO logic in short. As a consequence, a language that can be defined using MSO logic admits an efficient algorithm for membership test, marking a regular language precisely as a language in which logical, algebraic and algorithmic behavior squarely meet. And these nice characterizations generalize to labelled trees, which extends the path-like shape inherent in strings (Thatcher and Wright, 1968). Courcelle’s theorem (Courcelle 1990, Courcelle-Makowski-Rotices 2000) radically broadened the applications of Büchi-Elgot-Trakhtenbrot theorem on the equivalence of “recognizability and definability” on strings. Many fundamental graph problems can be expressed using MSO logic. And many such problems can be efficiently solved via so-called dynamic programming algorithms when the input graph is tree-like, e.g. has constant treewidth or constant cliquewidth. Courcelle’s theorem revealed that the availability of dynamic programming algorithms for those problems is not a coincidence by showing that any MSO-definable graph problem has a nice algebraic property. This in turn leads to a meta-algorithm, which is a finite-state (tree) automaton in essence. Whether the opposite implication holds on graph with tree-like structures, thus allowing a nice characterization akin to those on strings and labelled trees, was asked by Courcelle himself (1990). The converse statement of Courcelle’s theorem turns out to hold. It is proved by

Bojańczyk and Pilipczuk (2016) that if such a nice algebraic property holds for a graph property, the property admits a logical description by MSO logic on graphs of bounded treewidth. As Courcelle’s theorem holds on graphs of bounded cliquewidth as well as on matroids of bounded branchwidth when representable over a finite field, it is conjectured that “recognizability equals definability” on these more general structures. In this talk, we overview how algorithms design is intrinsically intertwined with algebraic and logical aspects of problems on graphs with tree-like structure. We also give a brief overview on essential, powerful tools for tackling these conjectures.

Monday 15:00-15:30

Learning-augmented online bipartite fractional matching

Davin Choo¹, Billy Jin², Yongho Shin³

¹ Havard University

² Purdue University

³ University of Wrocław

Online bipartite matching is a fundamental problem in online optimization, extensively studied both in its integral and fractional forms due to its theoretical significance and practical applications, such as online advertising and resource allocation. Motivated by recent progress in learning-augmented algorithms, we study online bipartite fractional matching when the algorithm is given advice in the form of a suggested matching in each iteration. We develop algorithms for both the vertex-weighted and unweighted variants that provably dominate the naive “coin flip” strategy of randomly choosing between the advice-following and advice-free algorithms. Moreover, our algorithm for the vertex-weighted setting extends to the AdWords problem under the small bids assumption, yielding a significant improvement over the seminal work of Mahdian, Nazerzadeh, and Saberi (EC 2007, TALG 2012). Complementing our positive results, we establish a hardness bound on the robustness-consistency tradeoff that is attainable by any algorithm. We empirically validate our algorithms through experiments on synthetic and real-world data.

Monday 16:00-18:00

Combinatorial optimization through the lens of boolean polynomials and binary matroids

Ahmad Abdi¹, Dabeen Lee²

¹ London School of Economics

² KAIST

Combinatorial optimization is the problem of choosing an optimal combination of items from a discrete set of elements. One of the most pressing challenges in modern applications is to develop efficient methodologies for nonlinear combinatorial optimization. In this talk, we introduce a novel approach to solving nonlinear problems through the lens of boolean polynomials and binary matroids. Discovering previously unrecognized connections to binary matroids, we develop a new framework for linearizing the equivalent boolean polynomial optimization formulation. We provide a simple characterization for the convex hull of binary solutions to the linearization when the underlying binary matroid is a projective geometry, which contains all binary matroids over the same ground set as restrictions. We also show that if the matroid satisfies the sums of circuits property, then the convex hull coincides with the associated cocycle polytope. Moreover, taking the signs of the objective coefficients into account, we extend the sums of circuits property to idealness of binary clutters via signed binary matroids. To go further beyond the sums of circuits property, we present a sum-of-squares hierarchy that admits efficient semidefinite relaxations. Lastly, we explain extensions to discrete optimization over any finite fields.

Random perturbation of dense graphs

Jie Han¹, Seonghyuk Im^{2,3}, Bin Wang¹, Junxue Zhang¹

¹ Beijing Institute of Technology

² KAIST

³ IBS Extremal Combinatorics and Probability Group

A randomly perturbed graph is formed by taking the union of a deterministic graph G and a binomial random graph $G(n, p)$ on the same vertex set. Balogh, Treglown, and Wagner showed that for a fixed graph F , if $p \gg n^{-1/m_1(F)}$ and an n -vertex graph G has the minimum degree $\Omega(n)$, then the graph $G \cup G(n, p)$ contains an F -factor with high probability. We extend their result by proving that if F is not a forest, then it suffices to assume $e(G) = \Omega(n^2)$, relaxing the minimum degree condition. Additionally, we establish analogous results for powers of Hamilton cycles and for families of graphs with bounded maximum degree in a randomly perturbed graph.

On a Ramsey–Turán variant of Roth’s theorem

Matija Bucić¹, Micha Christoph², Jaehoon Kim³, Hyunwoo Lee^{3,4}, Varun Sivashankar¹

¹ Princeton University

² ETH Zürich

³ KAIST

⁴ IBS Extremal Combinatorics and Probability Group

A classical theorem of Roth states that the maximum size of a solution-free set of a homogeneous linear equation \mathcal{L} in \mathbb{F}_p is $o(p)$ if and only if the sum of the coefficients of \mathcal{L} is 0. In this paper, we prove a Ramsey–Turán variant of Roth’s theorem, with respect to a natural notion of “structured” sets introduced by Erdős and Sárközy in the 1970’s. Namely, we show that the following statements are equivalent:

- (a) Every solution-free set A of \mathcal{L} in \mathbb{F}_p with $\alpha(\text{Cay}_{\mathbb{F}_p}(A)) = o(p)$ has size $o(p)$.
- (b) There exists a non-empty *subset* of coefficients of \mathcal{L} with zero sum.

On attainable homomorphism domination exponents

Ingyu Baek¹, Joonkyung Lee¹

Yonsei University

For graphs H and F , the inequalities of the form $t(H, G) \geq t(F, G)^s$, where $t(H, G)$ denotes the homomorphism density of H in G , received considerable attention during the recent decades. For example, Sidorenko’s conjecture, one of the major open problems in graph theory, asks whether such inequalities for bipartite graphs H , $F = K_2$, and $s = e(H)$ hold for arbitrary graphs G . Towards more systematic study of such inequalities, Blekherman, Raymond, Razborov, and Wei recently defined the *domination exponent* $C(H, F)$, which denotes the infimum of $s > 0$ such that $t(H, G) \geq t(F, G)^s$ holds for all graphs G . The first open problem they asked is whether there exists a graphon W bounded away from 0 (and 1) such that $t(H, W) = t(F, W)^{C(H, F)}$. We answer this question in the negative, by showing that there exists infinitely many graphs H such that the equality in $t(H, W) \geq t(K_3, W)^{C(H, K_3)}$ is never attained by graphons W bounded away from 0 and 1. This suggests that using ‘sparse’ graphs G , e.g., the Ruzsa–Szemerédi graphs, is somewhat unavoidable in the systematic study of graph homomorphism inequalities.

Tuesday 09:30-10:30

Combinatorics of orthogonal polynomials in the q -Askey scheme

Dongsu Kim

IS

KAIST

Combinatorics of orthogonal polynomials has been studied from 1980's. There are various combinatorial interpretations for important orthogonal polynomials and their properties. Orthogonal polynomials in q -Askey scheme have q -factorials in their terms and coefficients. A nice combinatorial model to handle q -factorials has been introduced in Lecture hall graphs and the Askey scheme, by S. Corteel, B. Jonnadula, J. Keating, J. S. Kim, arXiv:2311.12761v2. This talk is a history of the combinatorics of orthogonal polynomials up to the above paper.

Tuesday 11:00-12:00

Combinatorial Bijections in Permutations, Trees, and Paths

*Heesung Shin*¹

Inha University

This presentation traces my two-decade journey exploring combinatorial bijections in enumerative combinatorics. My research establishes strong connections and provides combinatorial interpretations for algebraic identities between permutations, trees (rooted, ordered, labeled, p -ary), and lattice paths (Dyck, Schröder, Delannoy, F -paths). Key contributions include bijections for pattern-avoiding inversion sequences and permutations, refined enumerations of trees based on various properties like indegree sequences and maximal decreasing subtrees, and analysis of lattice paths with avoidance conditions. The work also covers bijections for parking functions and involutions on partitions for hook length symmetry. This talk emphasizes the power of bijections in solving problems and revealing new connections in combinatorial theory.

On the graph a -numbers

Suyoung Choi¹, Younghan Yoon¹

Ajou University

We introduce combinatorial invariants, called a -numbers, of finite simple graphs, motivated by toric topology. These invariants exactly correspond to the Betti numbers of the canonical real toric variety associated with graphs. In this talk, we explore two fundamental questions about the behavior of the a -numbers:

1. Are they monotone increasing under edge inclusion?
2. Do they form a unimodal sequence?

Tuesday 13:30-14:30

Partition statistics and the Littlewood decomposition

(in honor of Professor Dongsu Kim's retirement)

Ae Ja Yee

IS

Pennsylvania State University

Integer partitions carry various interesting statistics. Of those, the most loved and studied statistics are Dyson's rank and crank, which explain Ramanujan's mod 5, 7 and 11 partition congruences. In 1990, Garvan, Kim and Stanton found other cranks, which split the set of partitions into t equinumerous classes for $t = 5, 7, 11$, and thus give a combinatorial account for the three congruences of Ramanujan. In their proof, the Littlewood decomposition of partitions into t -core and t -quotient partitions is an essential component. Since then, their cranks along with Dyson's rank and crank have been adopted to prove other partition congruences and refinements. In this talk, I will discuss the Littlewood decomposition from a partition theory point of view and present some recent results on various partition statistics arising from the Littlewood decomposition. This talk is based on joint work with Hyunsoo Cho, Byungchan Kim and Eunmi Kim.

Tuesday 14:30-15:30

On toric Schubert varieties in flag varieties

Inseo Kim¹, Eunjeong Lee¹

Chungbuk National University

Let G be a simple algebraic group over \mathbb{C} , and let P be a parabolic subgroup of G . A *flag variety* G/P is a smooth projective homogeneous variety that admits an action of a maximal torus T of G , where T is contained in the parabolic subgroup P . Schubert varieties form an interesting family of T -invariant subvarieties of G/P , indexed by elements of a certain coset of the Weyl group of G . Note that not every Schubert variety is toric with respect to the induced T -action. In this talk, we consider *toric* Schubert varieties in G/P when G is of type A .

The largest size of an $(s, s + 2)$ -core partition with even or odd parts only

*Hyoeun Cho*¹, *Seong hyeon Hwan*², *Seonkyung Kim*², *Hayan Nam*³

¹ Yonsei University

² Kangwon National University

³ Konkuk University

Olsson and Stanton initiated the computation of the largest size of a simultaneous core partition. Since then, there have been various works on the largest size of a core partition. In particular, Nam and Yu computed the largest size of an $(s, s + 1)$ -core partition, where all the parts are of the same parity. In this paper, we evaluate the largest size of an $(s, s + 2)$ -core partition, where all the parts are of the same parity.

Tuesday 16:00-16:30

Tree structures and combinatorics of bi- γ positivity

Zhicong Lin

IS

Shandong University

Multiset Eulerian polynomials and $1/k$ -Eulerian polynomials are different generalizations of the classical Eulerian polynomials. I will talk about combinatorics of their bi- γ -positivity via tree structures.

Tuesday 16:30-18:00

On the extension problem on the moment curve

Seunghun Lee¹, Eran Nevo²

¹ IBS Discrete Mathematics Group

² Hebrew University

We show that for $2 \leq d \leq 4$, every finite geometric simplicial complex Δ with vertices on the moment curve in \mathbb{R}^d can be extended to a triangulation T of the cyclic polytope C where Δ, T and C all have same vertex set. Further, for $d \geq 5$ we construct complexes Δ for which no such triangulations T exist.

Bier spheres and their full subcomplexes

Suyoung Choi¹, Younghan Yoon¹, Seonghyeon Yu¹

Ajou University

In 1992, Thomas Bier introduced a combinatorial construction that yields a large family of simplicial $(m-2)$ -dimensional PL -spheres on $2m$ vertices. In algebraic topology, the full subcomplexes of a simplicial complex K often provide significant information about topological invariants of K or the topological space associated with K . In this talk, we will discuss full subcomplexes and the bigraded Betti numbers of Bier spheres.

Wednesday 09:30-10:30

On problems related to digraph width parameters

O-joung Kwon

IS

¹ Hanyang University

² IBS Discrete Mathematics Group

Treewidth is a well-known graph parameter that plays an important role in Robertson and Seymour's graph minors project. Several width parameters inspired by treewidth have been introduced for digraphs, including directed treewidth, directed pathwidth, DAG-width, Kelly-width, cycle rank, DAG-depth, and so on. Compared to undirected width parameters, finding unavoidable structures for digraph width parameters turns out to be very difficult. Recently, we provided a collection of three obstruction families for cycle rank in terms of butterfly minors. In this talk, we present this result and related open problems. This is joint work with Meike Hatzel, Myounghwan Lee, and Sebastian Wiederrecht.

Wednesday 11:00-12:30

Unavoidable butterfly minors in digraphs of large cycle rank

Meike Hatzel¹, O-joung Kwon^{2,1}, Myounghwan Lee², Sebastian Wiederrecht³

¹ IBS Discrete Mathematics Group

² Hanyang University

³ KAIST

Cycle rank is one of the depth parameters for digraphs introduced by Eggan in 1963. We show that there exists a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that every digraph of cycle rank at least $f(k)$ contains a *directed cycle chain*, a *directed ladder*, or a *directed tree chain* of order k as a butterfly minor. *Directed cycle chains* and *directed ladders* are strongly connected digraphs that are obtained from an undirected path by splitting its edges or vertices in a specific way. A *directed tree chain* of order k is obtained from a directed path $v_1 v_2 \cdots v_{2^k}$ by adding the edge $(v_{2^i j}, v_{2^i j + 2^{i-1}})$ for every $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, 2^{k-i}\}$. We also investigate a new connection between cycle rank and a directed analogue of the weak coloring number of graphs.

Subtournaments of tournaments of large clique-width

Seokbeom Kim^{1,2}, *Taite LaGrange*³, *Mathieu Rundström*³, *Sophie Spirkl*³

¹ KAIST

² IBS Discrete Mathematics Group

³ University of Waterloo

If a graph has large clique-width, what can we say about its induced subgraphs? From classical results on clique-width, it is not hard to deduce that the class of H -free graphs has bounded clique-width if and only if H is an induced subgraph of a four-vertex path. Here, a graph is H -free if it has no induced subgraph isomorphic to H . Motivated by this result, there have been studies on the clique-width of graphs excluding two or more graphs as induced subgraphs, as well as H -free graphs with additional structural properties. In this talk, we investigate an analogue of this question for tournaments, which are a special type of directed graph. Again, what can we say about subtournaments of a tournament with large (directed) clique-width? We present recent progress on this question along with some open problems.

The square of every subcubic planar graph without 4-cycles and 5-cycles is 7-choosable

*Ligang Jin*¹, *Yingli Kang*², *Seog-Jin Kim*³

¹ Zhejiang Normal University

² Jinhua University of Vocational Technology

³ Konkuk University

The square of a graph G , denoted G^2 , has the same vertex set as G and has an edge between two vertices if the distance between them in G is at most 2. Thomassen (2018) proved that $\chi(G^2) \leq 7$ if G is a subcubic planar graph. A natural question is whether $\chi_\ell(G^2) \leq 7$ or not if G is a subcubic planar graph. Recently, Kim and Lian (2024) proved that $\chi_\ell(G^2) \leq 7$ if G is a subcubic planar graph of girth at least 6. In this paper, we prove that $\chi_\ell(G^2) \leq 7$ if G is a subcubic planar graph without 4-cycles and 5-cycles, which improves the result of Kim and Lian.

List of Participants

1. **Ingyu Baek** 백인규
Yonsei University
2. **Jihyo Chae** 채지효
Yonsei University
3. **Yeonsu Chang** 장연수
Hanyang University
4. **Gi-Sang Cheon** 천기상
Sungkyunkwan University
5. **Qiyuan Chen**
Chinese Academy of Sciences
6. **Hyunsoo Cho** 조현수
Ewha Womans University
7. **Minho Cho** 조민호
KIAS
8. **Hojin Chu** 추호진
KIAS
9. **Younggwang Cho** 조영광
Sungkyunkwan University
10. **Wansoo Choi** 최완수
Seoul National University
11. **Sylvia Corteel**
CNRS & UC Berkeley
12. **Taehyun Eom** 엄태현
Chonnam National University
13. **Yijia Fang**
National University of Singapore
14. **Colin Geniet**
IBS Discrete Mathematics Group
15. **Cheolwon Heo** 허철원
SUNY Korea
16. **Taehee Hong** 홍태희
Seoul National University
17. **JiSun Huh** 허지선
University of Seoul
18. **Junhyuk Huh** 허준혁
University of Cambridge
19. **Tony Huynh**
IBS Discrete Mathematics Group
20. **Bokhee Im** 임복희
Chonnam National University
21. **Seonghyuk Im** 임성혁
KAIST & IBS ECOPRO
22. **Musung Kang** 강무성
Seoul National University
23. **Soon-Yi Kang** 강순이
Kangwon National University
24. **Dongsu Kim** 김동수
KAIST
25. **Eun Jung Kim** 김은정
KAIST & IBS DIMAG & CNRS
26. **Gunwoo Kim** 김건우
KAIST & IBS DIMAG

27. **Hyobeen Kim** 김효빈
Chonnam National University
28. **Inseo Kim** 김인서
Chungbuk National University
29. **Jang Soo Kim** 김장수
Sungkyunkwan University
30. **Jeewon Kim** 김지원
KAIST
31. **Jeongsoo Kim** 김정수
Chungbuk National University
32. **Sangwook Kim** 김상욱
Chonnam National University
33. **Seog-Jin Kim** 김석진
Konkuk University
34. **Seokbeom Kim** 김석범
KAIST & IBS DIMAG
35. **Seonkung Kim** 김선경
Kangwon National University
36. **O-joung Kwon** 권오정
Hanyang University & IBS DIMAG
37. **Young Soo Kwon** 권영수
Yeungnam University
38. **Jae Ha Kyung** 경제하
Chungbuk National University
39. **Chanhee Lee** 이찬희
Kangwon National University
40. **Dabeen Lee** 이다빈
KAIST
41. **Dohyeon Lee** 이도현
KAIST & IBS DIMAG
42. **Eunjeong Lee** 이은정
Chungbuk National University
43. **Hyunwoo Lee** 이현우
KAIST & IBS ECOPRO
44. **Jongseo Lee** 이종서
KAIST
45. **Myounghwan Lee** 이명환
Hanyang University
46. **Seok Hyeong Lee** 이석형
Seoul National University
47. **Seunghun Lee** 이승훈
IBS Discrete Mathematics Group
48. **Zhicong Lin**
Shandong University
49. **Sungmin Moon** 문성민
KAIST
50. **Sunyo Moon** 문선요
KIAS
51. **Hayan Nam** 남하얀
Konkuk University
52. **Semin Oh** 오세민
Kyungpook National University
53. **Sang-il Oum** 엄상일
IBS Discrete Mathematics Group
54. **Hyemi Park** 박혜미
Hanyang university

55. **Jeong Rye Park** 박정례
Kyungpook National University

56. **Jongyook Park** 박종욱
Kyungpook National University

57. **Homoon Ryu** 류호문
Ajou University

58. **Jaehyeon Seo** 서재현
Yonsei University

59. **Seunghyun Seo** 서승현
Kangwon National University

60. **Heesung Shin** 신희성
Inha University

61. **Yongho Shin** 신용호
University of Wrocław

62. **Jaebum Sohn** 손재범
Yonsei University

63. **Ae Ja Yee** 이애자
The Pennsylvania State University

64. **Jaeyeong Yoo** 유재영
Kangwon National University

65. **Meesue Yoo** 류미수
Chungbuk National University

66. **Semin Yoo** 유세민
IBS Discrete Mathematics Group

67. **Younghan Yoon** 윤영한
Ajou University

68. **Seonghyeon Yu** 유성현
Ajou University

69. **Changki Yun** 윤창기
Seoul National University

Useful Information

Talks will be held at the **Auditorium** on the 2nd floor of the IBS Science Culture Center. Wi-Fi will be available during the conference.

The location of the **conference dinner** on Monday will be announced later. The **conference banquet** on Tuesday will be held at the IBS main building on the ground floor.

How to get to the IBS?

The detailed travel instructions is available on <https://travel.dimag.kr>.

Accommodation

Designated hotels for the conference are as follows: Onoma Hotel, Lotte City Hotel Daejeon, Hotel ICC, I-HOTEL.

Lunch on Tuesday

Lunch on Tuesday will not be provided. Here is a list of restaurants near the IBS:

- **IBS Cafeteria** (1st floor of the IBS main building): 5,000 KRW per person.
- **Shinsaegae Department Store** (5-minute walk from the IBS): Various restaurants available on the basement floor and the 5th floor.
- **Il Forno (일뿐르노)**: Italian self-service restaurant on the 39th floor of the Shinsaegae Expo Tower, which is where the Onoma Hotel is located. Has a great view.

More information: <https://travel.dimag.kr/localinfo/>

Sponsors



