

Invited Lecture IL 08
Molecular Behavior of 1-Pentanol Guest in the NH₄F-doped Clathrate Hydrate
Authors and affiliation
Byeonggwon Lee ^{1,2} , Jeongtak Kim ^{3,4} , <u>Kyuchul Shin</u> ^{1,3,*} , Ki Hun Park ⁵ , Minjun Cha ⁵ , Saman Alavi ^{6,7,*} , John A. Ripmeester ⁷
¹ Dept. of Hydrogen & Renewable Energy, Kyungpook National University, Daegu 41566, Republic of Korea
² Radioactive Waste Treatment Research Team, Korea Atomic Energy Research Institute, Daejeon 34057, Republic of Korea
³ Dept. of Applied Chemistry, Kyungpook National University, Daegu 41566, Republic of Korea
⁴ Climate Change Research Division, Korea Institute of Energy Research, Daejeon 34129, Republic of Korea
⁵ Dept. of Energy and Resources Engineering, Kangwon National University, Gangwon-do 24341, Republic of Korea
⁶ Department of Chemistry and Biomolecular Sciences, University of Ottawa, Ottawa, Ontario K1N 6N5, Canada.
⁷ National Research Council of Canada, 100 Sussex Dr., Ottawa, Ontario K1A 0R6, Canada
Key Word (3 words)
Clathrate Hydrate, Hydrogen Bonding, Ammonium Fluoride
Abstract (less than 300 words)
<p>In pure water and even in the presence of help gases, 1-pentanol (1-PeOH) cannot form the clathrate hydrate structures both because of its size that is not appropriate for either the structure II $5^{12}6^4$ or structure H $5^{12}6^8$ cages and the presence of a hydrophilic OH group that tends to destabilize the clathrate hydrate framework. [1] Recently, the NH₄F doping in the clathrate framework was suggested to make the hydrate lattice more resistant to disruption by host-guest hydrogen bonding. [2,3] In this work, we report the synthesis of 1-pentanol + methanol binary structure II clathrate hydrate doped with NH₄F and demonstrate that the space filling of 1-pentanol as a guest molecule can be managed by the hydrogen bonding between doped NH₄F in the water framework and the hydroxyl group of 1-pentanol. The structural characterization of 1-PeOH + methanol hydrates was conducted by the powder X-ray diffraction (PXRD) pattern analysis and the solid state ¹³C NMR spectroscopy. The 1-PeOH guest conformations and molecular interactions in the clathrate cages and the guest conformations was also investigated by the solid state ¹³C NMR spectroscopy and molecular dynamics simulations. Additionally, the 1-PeOH hydroxyl group was observed to be incorporated in the clathrate hydrate water framework. The findings in this work show that doping the clathrate hydrate lattice with NH₄F and using methanol as a helper guest can be used for stabilizing heavier alcohol molecules which cannot be engaged in normal clathrate hydrates.</p>
References
[1] J. A. Ripmeester, C. I. Ratcliffe, J. Phys. Chem. 94 (1990) 8773– 8776.
[2] K. Shin, I. Moudrakovski, M. Davari, S. Alavi, C. I. Ratcliffe, J. A. Ripmeester, CrystEngComm 16 (2013) 7209-7217.
[3] K. Shin, I. Moudrakovski, C. I. Ratcliffe, J. A. Ripmeester, Angew. Chem. Int. Ed. 56 (2017) 6171-6175.
MTMS '21