

| |
|---|
| PA 07 |
| Volumetric behavior of HFO-1234ze(E) + acetone liquid mixture at 303.2 K. |
| Authors and affiliation |
| <p>Atuski Tokoro¹, Masaki Okada², Taka-aki Hoshina^{*1}, Tomoya Tsuji³, and Takeshi Furuya⁴</p> <p>¹ College of Industrial Technology, Nihon University, 1-2-1 Izumicho, Narashino, Chiba 275-8575, Japan.</p> <p>² College of Science and Technology, Nihon University, 7-24-1 Narashinodai, Funabashi, Chiba 274-8501, Japan.</p> <p>³ Universiti Teknologi Malaysia (UTM) Kuala Lumpur, Off Jalan Sultan Yahya Petra 54100 Kuala Lumpur, Malaysia</p> <p>⁴ The National Institute of Advanced Industrial Science and Technology (AIST), 1-1-1 Higashi, Tsukuba, Ibaraki 305-8565 Japan</p> <p>*E-Mail: hoshina.takaaki@nihon-u.ac.jp</p> |
| Key Word (3 words) |
| Density, Hydrofluoroorefin, Binary mixture |
| Abstract (less than 300 words) |
| <p>Hydrofluorocarbons (HFC) have been widely used as refrigerants, forming agents and spray propellants. However, hydrofluoroorefins (HFO) are now considered to be an alternative for HFC because of zero ozone depletion potential (ODP) and its quite low global warming potential (GWP) [1]. The authors have investigated bubble point pressure [2,4], liquid density [3] and dielectric constant [4], for the mixtures of conventional propellant, propane or dimethyl ether (DME) with some organic solvents. However, there are few experimental data for the mixtures containing (E)-1,3,3,3-Tetrafluoropropene (HFO-1234ze(E)) except for the bubble point pressure reported previously [2].</p> <p>In this study, the density was measured for a liquid mixture of HFO-1234ze(E) + acetone at 1.0 MPa and 303.2 K, and the volumetric behavior of mixture was investigated for the mole fraction dependence of the excess molar volume.</p> <p>Density of mixture was measured by oscillation of U-shaped tube densimeter (Anton Parr 512P). The density of mixture was estimated by Oscillation period of U-shaped tube filled with liquid mixture. The densimeter was calibrated with butane[5] and water[6] at 1.0 MPa and 303.2 K.</p> <p>The density of liquid mixture for HFO-1234ze(E) + acetone studied monotonously increased with increasing mole fraction of HFO-1234ze(E) at 1.0 MPa and 303.2 K. The excess molar volume of this liquid mixture studied showed to be negative in the whole range of mole fraction of HFO-1234ze(E), and a minimum was seen around 60 mol% of HFO-1234ze(E). This may suggests that an attractive force among HFO-1234ze(E) and acetone will be important role of volumetric behavior in liquid mixture of HFO-1234ze(E) + acetone at 1.0 MPa and 303.2 K.</p> |
| References |
| <p>[1] Muhre, G. <i>et al.</i>, IPCC 5th assessment report, 732 (2013).</p> <p>[2] Nakazawa, M. <i>et al.</i>, Fluid Phase Equilibria, 522, 112764 (2020).</p> <p>[3] Hoshina, T. <i>et al.</i>, The review of high pressure science and technology, 23, 319 (2013).</p> <p>[4] Koshiha, Y. <i>et al.</i>, Kagaku Kogaku Ronbunshu, 44, 223 (2018).</p> <p>[5] Miyamoto, H <i>et al.</i>, Int. J. Thermophys., 22, 459 (2001).</p> <p>[6] Wagner, W. <i>et al.</i>, J. Phys. Chem. Ref. Data, 18, 1537 (2002).</p> |
| MTMS '21 |