

Use of Scale Morphology for Discriminating Wild Stocks of Atlantic Striped Bass

R. ANNE RICHARDS¹ AND CHRISTINE ESTEVES

National Marine Fisheries Service, Northeast Fisheries Science Center
166 Water Street, Woods Hole, Massachusetts 02543, USA

Abstract.—We used scale morphology to discriminate wild Atlantic striped bass *Morone saxatilis* from the Hudson River, Chesapeake Bay, and the Roanoke River. Morphological features used were perimeter shape as described by Fourier analysis, rectangularity, width and spacing of the first 10 circuli, and patterns of partitioning of the scale interior. Performance of discriminant functions was evaluated by examining the percent correct classification of known-origin samples (precision) and the bias in resulting stock composition estimates (accuracy). Correct classification rates ranged from 57% to 84%, varying with the number of stocks and year-classes included. Lowest rates were for three stocks with multiple year-classes; highest rates were for single-year-class analysis of the most geographically distant stocks. Despite some poor classification rates, bias and variance in stock composition estimates were low in four of five discrimination problems, suggesting that corrections based on the classification matrix could be applied with some confidence to unknown samples. However, we used bootstrapping to evaluate bias stability; further evaluation of independent samples would be useful.

Three major anadromous stocks of Atlantic striped bass *Morone saxatilis* occur along the east coast of the United States, spawning in the Hudson River, Chesapeake Bay, and the Roanoke River (Berggren and Lieberman 1978). Another major spawning stock once occurred in the Delaware River, but its abundance has been low for most of this century (Chittenden 1971; Rago et al. 1993). During late spring, summer, and fall, subadult and adult striped bass are distributed in coastal waters from North Carolina to Canada (Merriman 1941; Chapoton and Sykes 1961; Boreman and Lewis 1987), where they have supported important fisheries since colonial times. The Hudson River and Chesapeake Bay stocks appear to be more highly migratory than the Roanoke River population (Merriman 1941; Vladikov and Wallace 1952; Hassler et al. 1981; Boreman and Lewis 1987), and probably make up the bulk of the coastal stock. Striped bass stocks exhibit pronounced differences in maturation rates (Specker et al. 1987; Hoff et al. 1988; Olsen and Rulifson 1992) and recruitment levels (Boreman and Austin 1985; Richards and Deuel 1987), so stock composition is an important consideration in management of this species.

Although striped bass have been the subject of numerous stock identification studies in the past (Waldman et al. 1988), the need for information on stock composition has increased in recent years

as managers have attempted to tailor regulations to rebuild some stocks and maintain others (ASMFC 1981, 1989, 1990; USDOC 1993). Important fishery and population monitoring programs are conducted in coastal areas where striped bass stocks mix (Richards and Deuel 1987; Rago et al. 1993); however, without stock discrimination these programs cannot provide population-specific estimates of vital rates. Previous studies of southern New England striped bass fisheries have shown both temporal and spatial variation in stock composition (Fabrizio 1987; Van Winkle et al. 1988; Wirgin et al. 1993), and stock composition monitoring concurrent with other biological sampling is needed.

The purpose of our study was to determine whether the Hudson River, Chesapeake Bay, and Roanoke River stocks of Atlantic striped bass can be discriminated on the basis of scale morphology. Scales are routinely collected for age determination of striped bass, so sampling for stock identification would entail no further effort if scale morphology could be used. In a previous study of striped bass scales (Richards and Esteves 1997, this issue), Fourier shape descriptors, rectangularity, and some aspects of internal structure showed stock-specific differences that might form the basis for discriminating these stocks.

Methods

Morphological variables considered in this study were Fourier coefficients (describing perimeter shape), rectangularity (a ratio shape descrip-

¹ Present address: Chesapeake Biological Laboratory, Post Office Box 38, Solomons, Maryland 20688, USA.

TABLE 1.—Scale morphology variables included in discriminant analyses of striped bass from three stocks.

Variable	Hudson River versus Chesapeake Bay versus Roanoke River		Hudson River versus Chesapeake Bay		Hudson River versus Roanoke River, age 2
	All ages	Age 2	All ages	Age 2	
Fourier coefficients					
Amplitude coefficient	7, 10, 11, 12	1, 3, 5	3, 5–11	4, 5, 9, 11	5, 8, 11
Phase coefficient	3, 10	2, 3, 4, 7, 9	2, 8, 9, 11, 12	7, 9, 12	2, 3, 4, 7
Ratio Shape	Rectangularity	Rectangularity		Rectangularity	Rectangularity
Circulus patterns		Variance of widths 1–5 Variance of spaces 6–10 Average of widths 6–10		Average of widths 1–5	
Angles	Left angle	Left, right angles	Right angle; asymmetry		Left, right angles

tor), circulus deposition patterns, and angular measurements of the scales' interior. The scales were from striped bass collected during the spawning season on spawning grounds in the Hudson River, Chesapeake Bay, and Roanoke River. Descriptions of sampling and data collection methods are provided by Richards and Esteves (1997). A subset of variables was selected for use in discriminant analysis. Variables were included in discriminant analyses if they could be corrected for variation with size according to the methods of Reist (1986), if they showed significant stock-of-origin effects in a multivariate analysis of variance (MANOVA), and if they were not highly correlated with other included variables ($r < 0.70$) (Bliss 1970). We eliminated size-corrected variables that showed significant year-class effects because we did not wish to discriminate on the basis of age structure differences among stocks.

We performed five sets of discriminant analyses: (1) a three-stock analysis (Hudson River, Chesapeake Bay, and Roanoke River) of all year-classes; (2) a three-stock analysis of only age-2 fish; (3) a Chesapeake Bay versus Hudson River analysis of all year-classes; (4) a Chesapeake Bay versus Hudson River analysis of age-2 fish; and (5) a Hudson River versus Roanoke River analysis of age-2 fish. We varied the number of stocks examined because several studies have had limited success in distinguishing Roanoke River and Chesapeake Bay stocks (Grove et al. 1976; Fabrizio 1987; Margraf and Riley 1993). Examining only a single year-class allowed us to include variables that showed significant year-class effects or that could not be corrected for size effects. Age-2 samples (1988 year-class) provided the largest sample size across all stocks.

We tested the data sets for equality of covariance matrices. We used linear discriminant analysis if covariance matrices were equal and quadratic discriminant analysis if covariance matrices were unequal. Prior probabilities of group membership were assumed equal rather than proportional to sample size. We estimated the classification matrix by using jackknifing. Analyses were conducted using Statistical Analysis System version 6.03 (SAS Institute 1988).

Performance of the discriminant functions was evaluated via two measures: (1) the proportion of samples correctly classified to stock of origin; and (2) bias in stock composition estimates. We used the "misallocation statistic" (c) of Prager and Fabrizio (1990) as an index of bias in the stock composition estimates. The statistic varies between 0 (no bias) and 1 (maximum possible bias). Proportionate stock membership was estimated as the sum over all fish of the predicted probabilities of group membership. We developed confidence intervals (CIs) for the misallocation statistic and for stock composition estimates by bootstrapping (Efron 1982). The CIs included 95% of the outcomes of 400 bootstrap replicates.

Results

Three-Stock Discrimination, All Ages

Eight variables were selected for discrimination of three stocks with all year-classes (Table 1): six Fourier coefficients (four amplitude and two phase), rectangularity, and one angle measurement (left). With use of linear discriminant analysis, the overall correct classification rate for the three stocks was 56.6% (Table 2). The highest correct classification rate was for the Roanoke River stock

TABLE 2.—Classification matrices from five sets of discriminant analysis of known-origin samples of striped bass. Numbers with asterisks represent samples classified to their correct stock.

Source stock	Assigned stock						N
	Chesapeake Bay		Hudson River		Roanoke River		
	Number	%	Number	%	Number	%	
Three stocks, all ages (56.6% correct overall)							
Chesapeake Bay	98*	53.3*	47	25.5	39	21.2	184
Hudson River	29	24.8	66*	56.4*	22	18.8	117
Roanoke River	7	15.6	6	13.3	32*	71.1*	45
Total	134		119		93		346
Three stocks, age 2 (60.5% correct overall)							
Chesapeake Bay	28*	56.0*	9	18.0	13	26.0	50
Hudson River	10	32.3	20*	64.5*	1	3.2	31
Roanoke River	9	27.3	3	9.1	21*	63.6*	33
Total	47		32		35		114
Chesapeake Bay versus Hudson River, all ages (71.8% correct overall)							
Chesapeake Bay	139*	75.5*	45	24.5			184
Hudson River	40	34.2	77*	65.8*			117
Total	179		122				301
Chesapeake Bay versus Hudson River, age 2 (75.3% correct overall)							
Chesapeake Bay	38*	76.0*	12	24.0			50
Hudson River	8	25.8	23*	74.2*			31
Total	46		35				81
Hudson River versus Roanoke River, age 2 (84.4% correct overall)							
Hudson River			27*	87.1*	4	12.9	31
Roanoke River			6	18.2	27*	81.8*	33
Total			33		31		64

(71.1%); only 56.4% of Hudson River and 53.3% of Chesapeake Bay fish were correctly identified. The discriminant function performed better in terms of bias in stock composition estimates ($c = 0.03$; 95% CI = 0.01–0.06). The contributions of the Hudson River and Roanoke River stocks were overestimated by 4% and 6%, respectively, and the Chesapeake Bay stock was underestimated by 10% (Table 3).

Three-Stock Discrimination, Age 2

Using only age-2 fish allowed us to include three Fourier coefficients that showed significant year-class effects and it eliminated the need to correct most variables for size effects. Other changes in the variable set occurred as a result of reevaluating the selection criteria on this smaller data set (Table 1). The final variable set for discrimination included three amplitude and five phase Fourier coefficients, rectangularity, two angle measurements, and three circulus pattern variables (variance in the width of the first five circuli, variance in the intercirculus spacing of circuli 6 through 10, and average width of circuli 6 through 10).

The overall correct classification rate based on

linear discriminant analysis was 60.5%. The rate for the Chesapeake Bay (56.0%) was lower than for the Hudson River (64.5%) and the Roanoke River (63.6%) (Table 2). Bias in stock composition estimates ($c = 0.13$; 95% CI = 0.02–0.28) was higher than for the three-stock problem involving all age-classes. The proportional contribution of the Chesapeake Bay stock was overestimated by 18%, and the Hudson River and Roanoke River contributions were underestimated by 7% and 11%, respectively (Table 3).

Chesapeake Bay versus Hudson River, All Ages

Fifteen variables were used in the Chesapeake Bay versus Hudson River, multi-age-class analysis: eight amplitude and five phase descriptors, one angle measurement, and angular asymmetry (Table 1). The overall correct classification rate by quadratic discriminant analysis was 71.8% (Hudson River, 65.8%; Chesapeake Bay, 75.5%; Table 2). Bias in the stock composition estimates was low ($c = 0.01$; 95% CI = 0–0.03) and only 1% of each stock was misallocated (Table 3).

TABLE 3.—Stock composition estimates (95% confidence limits are in parentheses) from five sets of discriminant analyses of striped bass from three stocks: P_j = true proportion in stock j ; \hat{P}_j = estimated proportion in stock j based on predicted probabilities of stock membership; Δ = difference between actual and estimated proportions; c = misallocation statistic, ranging from 0 to 1. Statistics were calculated from 400 bootstrap replicates.

Stock and statistic	Chesapeake Bay versus Hudson River versus Roanoke River		Chesapeake Bay versus Hudson River	
	All ages	Age 2	All ages	Age 2
Proportionate stock membership				
Chesapeake Bay				
P_j	0.53 (0.48, 0.58)	0.44 (0.35, 0.53)	0.61 (0.54, 0.66)	0.62 (0.51, 0.73)
\hat{P}_j	0.43 (0.39, 0.47)	0.62 (0.39, 0.80)	0.60 (0.55, 0.64)	0.64 (0.52, 0.78)
Δ	-0.10 (-0.09, -0.11)	0.18 (0.04, 0.27)	-0.01 (0.01, -0.02)	0.02 (0.01, 0.05)
Hudson River				
P_j	0.34 (0.29, 0.39)	0.27 (0.20, 0.36)	0.39 (0.33, 0.46)	0.38 (0.27, 0.49)
\hat{P}_j	0.38 (0.34, 0.42)	0.20 (0.05, 0.38)	0.40 (0.36, 0.45)	0.36 (0.22, 0.48)
Δ	0.04 (0.05, 0.03)	-0.07 (-0.15, -0.02)	0.01 (0.03, -0.01)	-0.02 (-0.05, -0.01)
Roanoke River				
P_j	0.13 (0.09, 0.16)	0.29 (0.21, 0.38)		
\hat{P}_j	0.19 (0.14, 0.24)	0.18 (0.07, 0.34)		
Δ	0.06 (0.05, 0.08)	-0.11 (-0.14, -0.04)		
Misallocation				
Misallocation statistic (c)	0.03 (0.01, 0.06)	0.13 (0.02, 0.28)	0.01 (0.0, 0.03)	0.03 (0.0, 0.12)

Chesapeake Bay versus Hudson River, Age 2

Nine variables were included in the single-year-class analysis of the Chesapeake Bay and Hudson River stocks: four amplitude and three phase coefficients, rectangularity, and average width of the first five circuli (Table 1). Quadratic discriminant analysis resulted in correct classification of 76.0% of Chesapeake Bay and 74.2% of Hudson River samples (75.3% overall; Table 2). Bias in the resulting stock composition estimates was low ($c = 0.03$; 95% CI = 0–0.12), and 2% of the samples were misallocated between stocks (Table 3).

Hudson River versus Roanoke River, Age 2

We used 10 variables in the analysis of Hudson River versus Roanoke River, including three amplitude coefficients, four phase coefficients, rectangularity, and right and left angles (Table 1). By quadratic discriminant analysis, 87.1% of the Hudson River and 81.8% of the Roanoke River fish were correctly identified, for an overall correct rate of 84.4% (Table 2). The stock composition estimates had low bias ($c = 0.04$; 95% CI = 0–0.14) and 3% of the samples were misallocated between the stocks (Table 3).

Discussion

The precision of discriminant analysis based on striped bass scale morphology varied widely

among comparisons. The two analyses that included all three stocks had correct rates averaging only 57% and 61%. Although better than random, such rates are too low for three-stock discrimination to be practical for management. Exclusion of the Roanoke River stock added about 15% to correct classification rates, bringing them into the range of other stock identification studies of Chesapeake Bay and Hudson River striped bass (Waldman et al. 1988). The best overall rate (84%) was achieved for the comparison of Hudson River with Roanoke River fish, which involved the most geographically distant stocks considered.

Although most stock identification studies focus on correct classification rates, the adequacy of a stock identification method depends also on the stock composition estimates produced. If correct classification rates are low, but misclassifications are balanced, the estimate of stock composition may be accurate even though individual fish are not always classified accurately. This point is well illustrated by our results for the Hudson–Chesapeake analysis of age-2 fish. The overall correct classification rate was only 75.3%; however, the estimates of stock composition were very close to the true values (Table 3). Because stock composition estimates are the ultimate goal of discrimination, the distribution of misallocations is a key issue (Prager and Fabrizio 1990). Stock compo-

TABLE 3.—Extended.

Stock and statistic	Hudson River versus Roanoke River, age 2
Proportionate stock membership	
Chesapeake Bay	
P_j	
\hat{P}_j	
Δ	
Hudson River	
P_j	0.48 (0.37, 0.60)
\hat{P}_j	0.51 (0.34, 0.66)
Δ	0.03 (0.03, 0.06)
Roanoke River	
P_j	0.52 (0.40, 0.63)
\hat{P}_j	0.49 (0.34, 0.66)
Δ	-0.03 (-0.06, 0.03)
Misallocation	
Misallocation statistic (c)	0.04 (0.0, 0.14)

sition estimates resulting from our discriminant analyses were quite accurate. In four of five problem sets, misallocation statistics (c) were 0.04 or lower, and most estimated stock compositions differed very little from true compositions.

Even more important than the observed pattern of misallocations, however, is potential variability in those patterns. Bias in stock composition estimates can be corrected by using a classification matrix for known-origin samples (e.g., by following methods outlined by Van Winkle et al. 1988). However, fish in an unknown sample will necessarily differ from those used to develop the discriminant function and its correction. Thus an important issue is consistency of bias from sample to sample. Consistency presumably depends on the representativeness of the known-origin samples; if the sample is adequate, variability in the bias estimates should be low. We attempted to address this issue for our samples by bootstrapping confidence intervals for the stock composition estimates and misallocation statistics. The confidence limits for the estimates were quite narrow, suggesting that bias correction could be applied with some assurance. In four of the five problem sets, confidence limits for the estimates showed a spread similar to the true (bootstrap) proportions, suggesting that the pattern of misallocations was fairly stable. These results are undoubtedly somewhat

optimistic, however, because bootstrap replicates do not represent completely independent samples.

Based on results of other stock identification studies that incorporated scale shape analysis (Jarvis et al. 1978; Casselman et al. 1981; DePontual and Prouzet 1987; Riley and Carline 1982; Margraf and Riley 1993), we expected to see higher rates of correct classification for Atlantic striped bass. However, rates reported in some previous studies may be optimistic. For example, Jarvis et al. (1978) reported 80–100% correct classification of walleyes *Stizostedion vitreum* from two eastern Lake Erie stocks. Correct classification rates were estimated by resubstitution (i.e., by using the same samples used to estimate the discriminant function), which is known to bias correct classification rates upward (Lachenbruch and Mickey 1968). For comparison with Jarvis et al., we estimated a discriminant function for our age-2 Chesapeake Bay and Hudson River samples using the first 12 harmonics of the Fourier series and compared correct rates obtained by resubstitution and jackknifing. With jackknifing, the overall correct classification rate was 65%; with resubstitution, the correct rate rose to 100%.

Two other studies of striped bass have achieved moderate success in discriminating stocks according to scale morphology. Margraf and Riley (1993) used Fourier analysis to discriminate Hudson River and Chesapeake Bay stocks with 80% correct classification. Ross and Pickard (1990) correctly classified 73% of wild and hatchery Pacific striped bass using Fourier coefficients, and 90% when circulus features were added to the analysis. Our results from using multiple scale features were similar to these, although circulus features were not as powerful in our analysis as in Ross and Pickard's. Greater differences in circulus patterns might be expected between wild and hatchery stocks than among wild stocks.

Why scale morphology does not provide a more distinct stock-specific signal for wild striped bass is not clear. If striped bass stocks return to natal estuarine areas to spawn, a high degree of genetic isolation is possible. However, mitochondrial DNA studies of Atlantic striped bass show unusually low levels of base sequence heterogeneity (Wirgin et al. 1990, 1993), suggesting that little genetic differentiation has occurred. Environmental effects may be more important in producing the subtle differences in scale morphology that allow some discrimination. We found significant year-class effects in 28% of our size-corrected vari-

ables, suggesting that environmental factors have a substantial influence.

In a study of otoliths from Atlantic cod *Gadus morhua*, Campana and Casselman (1993) concluded that variation in otolith shape was due primarily to differences in growth rates among stocks. If striped bass scale shape is similarly influenced by growth rate, it is not clear whether one can expect stock discrimination to be successful, because the evidence is contradictory regarding the degree and direction of growth rate differences among Atlantic striped bass stocks (Setzler et al. 1980; Coutant 1985; ASMFC 1990; Conover 1990).

Although discrimination based on scale morphology did not consistently provide high correct classification rates for wild Atlantic striped bass, it provided accurate estimates of stock composition in four of the five analyses we conducted. Thus the utility of scale morphology hinges on the stability of bias in stock composition estimates (i.e., on how reliably corrections can be made based on the classification matrix). Our results suggest the bias may be quite stable; however, this should be more stringently tested on independent samples of known-origin fish, because the bootstrapping technique we used is probably optimistic.

Acknowledgments

The samples for this study were collected by John Waldman as part of a larger study of stock identification methods for striped bass (Waldman et al. 1997). Funding was provided by the National Marine Fisheries Service through the Emergency Striped Bass Study (Section 7 of the Anadromous Fish Conservation Act, Public Law 96-118).

References

- ASMFC (Atlantic States Marine Fisheries Commission). 1981. Interstate fisheries management plan for the striped bass. ASMFC, Fisheries Management Report 1, Washington, D.C.
- ASMFC (Atlantic States Marine Fisheries Commission). 1989. Supplement to the striped bass FMP—amendment #4. ASMFC, Fisheries Management Report 15, Washington, D.C.
- ASMFC (Atlantic States Marine Fisheries Commission). 1990. Source document for the supplement to the striped bass FMP—Amendment #4. ASMFC, Fisheries Management Report 16, Washington, D.C.
- Berggren, T. J., and J. T. Lieberman. 1978. Relative contribution of Hudson, Chesapeake, and Roanoke striped bass, *Morone saxatilis*, stocks to the Atlantic coast fishery. U.S. National Marine Fisheries Service Fishery Bulletin 76:335–345.
- Bliss, C. I. 1970. Statistics in biology, volume 2. McGraw-Hill, New York.
- Boreman, J., and H. M. Austin. 1985. Production and harvest of anadromous striped bass stocks along the Atlantic coast. Transactions of the American Fisheries Society 114:3–7.
- Boreman, J., and R. R. Lewis. 1987. Atlantic coastal migration of striped bass. Pages 331–339 in M. J. Dadswell and five coeditors. Common strategies of anadromous and catadromous fishes. American Fisheries Society Symposium 1, Bethesda, Maryland.
- Campana, S. E., and J. M. Casselman. 1993. Stock discrimination using otolith shape analysis. Canadian Journal of Fisheries and Aquatic Sciences 50:1062–1083.
- Casselman, J. M., J. J. Collins, E. J. Crossman, P. E. Ihssen, and G. R. Spangler. 1981. Lake whitefish (*Coregonus clupeaformis*) stocks of the Ontario waters of Lake Huron. Canadian Journal of Fisheries and Aquatic Sciences 38:1772–1789.
- Chapoton, R. B., and J. E. Sykes. 1961. Atlantic coast migration of large striped bass as evidenced by fisheries and tagging. Transactions of the American Fisheries Society 90:13–20.
- Chittenden, M. E., Jr. 1971. Status of the striped bass, *Morone saxatilis*, in the Delaware River. Chesapeake Science 12:131–136.
- Conover, D. O. 1990. The relation between capacity for growth and length of growing season: evidence for and implications of countergradient variation. Transactions of the American Fisheries Society 119:416–430.
- Coutant, C. C. 1985. Striped bass, temperature, and dissolved oxygen: a speculative hypothesis for environmental risk. Transactions of the American Fisheries Society 114:31–61.
- DePontual, J., and P. Prouzet. 1987. Atlantic salmon, *Salmo salar* L., stock discrimination by scale-shape analysis. Aquaculture and Fisheries Management 18:277–289.
- Efron, B. 1982. The jackknife, the bootstrap and other resampling plans. Society for Industrial and Applied Mathematics, Philadelphia.
- Fabrizio, M. C. 1987. Growth-invariant discrimination and classification of striped bass stocks by morphometric and electrophoretic methods. Transactions of the American Fisheries Society 116:728–736.
- Grove, T. L., T. J. Berggren, and D. A. Powers. 1976. The use of innate tags to segregate spawning stocks of striped bass (*Morone saxatilis*). Pages 166–176 in M. Wiley, editor. Estuarine processes, volume 1. Academic Press, New York.
- Hassler, W. W., N. L. Hill, and J. T. Brown. 1981. The status and abundance of striped bass, *Morone saxatilis*, in the Roanoke River and Albemarle Sound, North Carolina, 1956–1980. North Carolina Department of Natural Resources and Community Development, Division of Marine Fisheries, Special Scientific Report 38, Morehead City.
- Hoff, T. B., J. B. McLaren, and J. C. Cooper. 1988. Stock

- characteristics of Hudson River striped bass. Pages 59–68 in L. W. Barnthouse, R. J. Klauda, D. S. Vaughan, and R. L. Kendall, editors. Science, law, and Hudson River power plants: a case study in environmental impact assessment. American Fisheries Society Monograph 4, Bethesda, Maryland.
- Jarvis, R. S., H. F. Klodowski, and S. P. Sheldon. 1978. New method of quantifying scale shape and an application to stock identification in walleye (*Stizostedion vitreum vitreum*). Transactions of the American Fisheries Society 107:528–534.
- Lachenbruch, P. A., and M. R. Mickey. 1968. Estimation of error rates in discriminant analysis. Technometrics 10:1–11.
- Margraf, F. J., and L. M. Riley. 1993. Evaluation of scale shape for identifying spawning stocks of coastal Atlantic striped bass (*Morone saxatilis*). Fisheries Research 18:163–172.
- Merriman, D. 1941. Studies on the striped bass (*Roccus saxatilis*) of the Atlantic coast. U.S. Fish and Wildlife Service Fishery Bulletin 50:1–77.
- Olsen, E. J., and R. A. Rulifson. 1992. Maturation and fecundity of Roanoke River–Albemarle Sound striped bass. Transactions of the American Fisheries Society 121:524–537.
- Prager, M. H., and M. C. Fabrizio. 1990. Comparison of logistic and discriminant analyses for stock identification of anadromous fish, with application to striped bass (*Morone saxatilis*) and American shad (*Alosa sapidissima*). Canadian Journal of Fisheries and Aquatic Sciences 47:1570–1577.
- Rago, P. J., R. M. Dorazio, R. A. Richards, D. G. Deuel, and C. D. Stephan. 1993. Emergency striped bass research study report for 1991. U.S. Department of the Interior and U.S. Department of Commerce, Washington, D.C.
- Reist, J. D. 1986. An empirical evaluation of coefficients used in residual and allometric adjustment of size covariation. Canadian Journal of Zoology 64:1363–1368.
- Richards, R. A., and D. G. Deuel. 1987. Atlantic striped bass: stock status and the recreational fishery. U.S. National Marine Fisheries Service Marine Fisheries Review 49(2):58–66.
- Richards, R. A., and C. Esteves. 1997. Stock-specific variation in scale morphology of Atlantic striped bass. Transactions of the American Fisheries Society 126:908–918.
- Riley, L. M., and R. F. Carline. 1982. Evaluation of scale shape for the identification of walleye stocks from western Lake Erie. Transactions of the American Fisheries Society 111:736–741.
- Ross, W. R., and A. Pickard. 1990. Use of scale patterns and shape as discriminators between wild and hatchery striped bass stocks in California. Pages 71–77 in N. C. Parker and five coeditors. Fish-marking techniques. American Fisheries Society Symposium 7, Bethesda, Maryland.
- SAS Institute. 1988. SAS/STAT TM user's guide, release 6.03 edition. SAS Institute, Cary, North Carolina.
- Setzler, E. M., and eight coauthors. 1980. Synopsis of biological data on striped bass, *Morone saxatilis* (Walbaum). NOAA Technical Report NMFS Circular 433.
- Specker, J. L., D. L. Berlinsky, H. D. Bibb, and J. F. O'Brien. 1987. Oocyte development in striped bass: factors influencing estimates of age at maturity. Pages 162–174 in M. J. Dadswell and five coeditors. Common strategies of anadromous and catadromous fishes. American Fisheries Society, Symposium 1, Bethesda, Maryland.
- USDOC (U.S. Department of Commerce). 1993. Spotlight: Atlantic striped bass. Pages 25–32 in Our living oceans. USDOC, NOAA Technical Memorandum NMFS-F/SPO-15, Washington, D.C.
- Van Winkle, W., K. D. Kumar, and D. S. Vaughan. 1988. Relative contributions of Hudson River and Chesapeake Bay striped bass stocks to the Atlantic coastal population. Pages 255–266 in L. W. Barnthouse, R. J. Klauda, D. S. Vaughan, and R. L. Kendall, editors. Science, law, and Hudson River power plants: a case study in environmental impact assessment. American Fisheries Society, Monograph 4, Bethesda, Maryland.
- Vladykov, V. D., and D. H. Wallace. 1952. Studies of the striped bass, *Roccus saxatilis* (Walbaum), with special reference to the Chesapeake Bay region during 1936–1938. Bulletin of the Bingham Oceanographic Collection, Yale University 14:132–177.
- Waldman, J. R., J. Grossfield, and I. Wirgin. 1988. Review of stock discrimination techniques for striped bass. North American Journal of Fisheries Management 8:410–425.
- Waldman, J. R., R. A. Richards, W. B. Schill, I. Wirgin, and M. C. Fabrizio. 1997. An empirical comparison of stock identification techniques applied to striped bass. Transactions of the American Fisheries Society 126:369–385.
- Wirgin, I. I., and five coauthors. 1993. Mitochondrial DNA variation in striped bass (*Morone saxatilis*) from Canadian rivers. Canadian Journal of Fisheries and Aquatic Sciences 50:80–87.
- Wirgin, I. I., P. Silverstein, and J. Grossfield. 1990. Restriction endonuclease analysis of striped bass mitochondrial DNA: the Atlantic coastal migratory stock. Pages 475–491 in N. C. Parker and five coeditors. Fish-marking techniques. American Fisheries Society Symposium 7, Bethesda, Maryland.

Received October 16, 1996

Accepted May 1, 1997