

Southeast Alaska Pink Salmon Preseason Forecast

Please direct any questions to: Sara Miller (sara.miller@alaska.gov).

Last updated: August 2023

Background ¹

Pink salmon *Oncorhynchus gorbuscha* runs in Southeast Alaska support a valuable commercial fishery (Clark et al. 2006), although annual abundance varies tremendously. An average of 33 million pink salmon per year were harvested in Southeast Alaska in the last 10 years (2012–2021), with a range of 8 million (2018 and 2020) to 95 million (2013) fish. Although the inseason management of these stocks is focused on monitoring daily harvest and fishing effort, and using aerial survey counts to assess whether adequate numbers of pink salmon are present to meet escapement goals (Piston and Heintz 2020), the fishing industry benefits from meaningful preseason forecasts in order to plan appropriately for the harvest, processing, transportation, and marketing of these fish. Pink salmon runs are notoriously difficult to forecast (Adkison 2002; Haeseker et al. 2005; Shevlyakov and Koval 2012; Radchenko 2020) due to the species’ propensity to respond dramatically to changes in the marine environment (Farley et al. 2020), the odd- and even-year cycles of abundance (Heard 1991; Ruggerone et al. 2003; Krkošek et al. 2011), and the fact that there is only one cohort (i.e., only one age class) in the fishery each year (Heard 1991). With only a 2-year life cycle, information about cohort strength is not available from jacks or siblings as with other species of salmon (Adkison and Peterman 2000; Haeseker et al. 2007); sibling recruit modeling is not possible with pink salmon. Beginning in the late 1960s, the Alaska Department of Fish and Game (ADF&G) implemented programs to improve pink salmon stock assessment but met with limited success in forecasting pink salmon runs. Forecast accuracy (i.e., absolute percent error) from 1981 through 2006 was 57%. Past forecasts relied primarily on measures of pink salmon spawning abundance or success (Jones and Hofmeister 1985; Hofmeister and Jones 1989; Hofmeister and Blick 1993), both of which are poorly known and explain little of the variation in annual recruitment, which is largely determined in the early marine environment (Parker 1968; Mortensen et al. 2000; Willette et al. 2001).

NOAA Alaska Fisheries Science Center, Auke Bay Laboratories (National Oceanic and Atmospheric Administration [NOAA]) initiated the Southeast Alaska Coastal Monitoring (SECM) project in 1997 (Orsi et al. 1997; Murphy et al. 1999) to identify relationships between the year-class strength of juvenile salmon and the biophysical parameters influencing their growth and survival, their prey and predator interactions, their habitat utilization, and their stock interactions in marine waters (Orsi et al. 2000, 2005, 2009). Through this project, standardized monthly sampling and trawl surveys have been conducted annually from May to August (the August survey was dropped in 2020) to collect ecosystem (oceanographic data such as temperature and salinity profiles of the water column, surface water samples, and zooplankton samples) and catch per unit effort (CPUE) data associated with juvenile salmon at 8 stations in Icy Strait, a major migration corridor in northern Southeast Alaska (Orsi et al. 2001, 2006, 2012). The environmental and oceanographic data provided through the SECM project has become one of the longest continuous time series of its kind for the North Pacific. A major finding of the SECM survey is that relative abundance of juvenile pink salmon in June and July was highly correlated to harvest of adults in the subsequent year. As a result, NOAA used peak juvenile pink salmon CPUE and environmental information collected during SECM surveys to forecast the Southeast Alaska pink salmon harvest starting in 2004 (Murphy et al. 1999; Wertheimer et al. 2009a, 2010a, 2011–2015, 2017, 2018; Murphy et al. 2019b).

In the past, ADF&G and NOAA produced separate Southeast Alaska preseason pink salmon forecasts. In 2007, ADF&G began adjusting their simple trend forecasts with juvenile pink salmon abundance data from the SECM survey (Heintz et al. 2007; Piston and Heintz 2014, 2017; Appendix A). Forecast accuracy (i.e., absolute percent error) improved from 57% (1981 to 2006 preseason forecasts) to 31% (2007 through 2017

¹The background section was taken directly from Miller, S. E., J. M. Murphy, S. C. Heintz, A. W. Piston, E. A. Fergusson, R. E. Brenner, W. W. Strasburger, and J. H. Moss. 2022. Southeast Alaska pink salmon forecasting models. Alaska Department of Fish and Game, Fishery Manuscript No. 22-03, Anchorage.

preseason forecasts; the 2018 preseason forecast was based on the average of 5 recent even-year harvests and did not use juvenile abundance indices from the SECM survey; Appendix B). The largest absolute percent error between the forecast and the actual harvest occurred in the 1987, 1988, 2006, and 2018 forecasts. In 2018, ADF&G and NOAA scientists collaborated to create a joint preseason forecast for 2019 (Piston et al. 2021a). The SECM project and Southeast Alaska pink salmon harvest forecasts are now conducted cooperatively by NOAA and ADF&G using the ADF&G research vessel (R/V) Medeia (Piston et al. 2021a, 2022). The current method (2019 to 2022 preseason forecasts) is to forecast the adult pink salmon harvest in a multiple linear regression model with peak monthly (June or July) juvenile pink salmon CPUE and a temperature index (Piston et al. 2019, 2020, 2021a, 2022). The temperature index is based on the overall average 20 m integrated water column temperature recorded during May–July or May–August at 8 stations in Icy Strait as part of the annual SECM survey (Icy Strait Temperature Index [ISTI]; Murphy et al. 2019a). Together, both agencies continue to examine alternative variables and statistical methods to improve annual forecasts.

Assessment

The current assessment follows Miller et al. 2022.

Advisory Announcements

- 2023 forecast
- 2022 forecast
- 2021 forecast
- 2020 forecast
- 2019 forecast

Assessment reports

- 2023 forecast report
- 2022 forecast report
- 2021 forecast report
- 2020 forecast report

Timeline

~end of August: Andy Piston or Teresa Fish will send a data sheet with the Icy Strait CPUE data, the ISTI temperature (May, June, July average), and harvest. It is optimal to rerun the last ten years of harvest data to update prior harvests in the dataset. Andy (or Teresa) will coordinate with NOAA for the CPUE and ISTI data.

Early October: Review preliminary model runs with NOAA and ADF&G Southeast Alaska Pink Salmon Forecasting Team.

~end of October: Final write-up due with re-run of final model with updated harvest data for the current year.

mid- November: Advisory Announcement due. This is the primary responsibility of the SEAK pink salmon manager and biologist. The biometrician reviews the announcement.

Southeast Alaska Pink Salmon Forecasting Team:

- Teresa Fish (ADF&G; Pink and chum salmon project leader)
- Andy Piston (ADF&G; CF Salmon Research Supervisor)
- Jim Murphy (NOAA; statistician)
- Wess Strasburger (NOAA; biologist)
- Sara Miller (ADF&G, biometrician)
- Emily Fergusson (NOAA; biologist)
- Andrew Gray (NOAA)

References

- Adkison, M. D. 2002. Preseason forecasts of pink salmon harvests in Southeast Alaska using Bayesian model averaging. *Alaska Fishery Research Bulletin* 9(1):1–8.
- Adkison, M. D., and R. M. Peterman. 2000. Predictability of Bristol Bay, Alaska sockeye salmon *Oncorhynchus nerka* returns 1 to 4 years in the future. *North American Journal of Fisheries Management* 20(1):69–80.
- Clark, J. H., A. McGregor, R. D. Mecum, P. Krasnowski, and A. M. Carroll. 2006. The commercial salmon fishery in Alaska. *Alaska Fishery Research Bulletin* 12(1):1–146.
- Farley Jr., E. V., J. M. Murphy, K. Cieciel, E. M. Yasumiishi, K. Dunmall, T. Sformo, and P. Rand. 2020. Response of pink salmon to climate warming in the northern Bering Sea. *Deep Sea Research Part II: Topical Studies in Oceanography* 177:104830.
- Haeseker, S. L., R. M. Peterman, Z. Su, and C. C. Wood. 2005. Retrospective evaluation of preseason forecasting models for pink salmon. *North American Journal of Fisheries Management* 25(3):897–918.
- Haeseker, S. L., B. Dorner, R. M. Peterman, and Z. Su. 2007. An improved sibling model for forecasting chum salmon and sockeye salmon abundance. *North American Journal of Fisheries Management* 27(2):634–642.
- Heard, W.R. 1991. Life history of pink salmon (*Oncorhynchus gorbuscha*). Pages 119–230 [In] C. Groot and L. Margolis, editors. *Pacific salmon life histories*. University of British Columbia Press, Vancouver, B.C.
- Heinl, S., X. Zhang, and H. Geiger. 2007. Pages 48–51 [In] D. M. Eggers. Run forecasts and harvest projections for 2007 Alaska salmon fisheries and review of the 2006 season. Alaska Department of Fish and Game, Special Publication No. 07-01, Anchorage.
- Hofmeister, K., and J. Blick. 1993. Pages 43–45 [In] H. J. Geiger, and H. Savikko, editors. Preliminary forecasts and projections for 1993 Alaska salmon fisheries and review of the 1992 season. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 5J93-04, Juneau.
- Hofmeister, K. T., and D. Jones. 1989. Pages 33–35 [In] H. J. Geiger, and H. Savikko, editors. Preliminary forecasts and projections for 1989 Alaska salmon fisheries. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 5J89-01, Juneau.
- Jones, J. D., and K. T. Hofmeister. 1985. Pages 29–31 [In] D. M. Eggers, editor. Preliminary forecasts and projections for 1985 Alaska salmon fisheries. Alaska Department of Fish and Game, Division of Commercial Fisheries, Informational Leaflet No. 244, Juneau.
- Krkošek, M., R. Hilborn, R. M. Peterman, and T. P. Quinn. 2011. Cycles, stochasticity and density dependence in pink salmon population dynamics. *Proceedings of the Royal Society B: Biological Sciences* 278(1714):2060–2068.

- Miller, S. E., J. M. Murphy, S. C. Heinl, A. W. Piston, E. A. Fergusson, R. E. Brenner, W. W. Strasburger, and J. H. Moss. 2022. Southeast Alaska pink salmon forecasting models. Alaska Department of Fish and Game, Fishery Manuscript No. 22-03, Anchorage.
- Mortensen, D., A. Wertheimer, C. M. Taylor, and J. Landingham. 2000. The relation between early marine growth of pink salmon, *Oncorhynchus gorbuscha*, and marine water temperature, secondary production, and survival to adulthood. *Fishery Bulletin* 98(2):319–335.
- Murphy, J. M., A. L. J. Brase, and J. A. Orsi. 1999. Survey of juvenile Pacific salmon in the northern region of southeastern Alaska, May–October 1997. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-AFSC-105.
- Murphy, J. M., E. A. Fergusson, A. Piston, S. Heinl, A. Gray, and E. Farley. 2019a. Southeast Alaska pink salmon growth and forecast models. NPAFC Technical Report 15:75–81. (Available at <http://www.npafc.org>)
- Murphy, J. M., A. C. Wertheimer, E. Fergusson, A. Piston, S. Heinl, C. Waters, J. Watson, and A. Gray. 2019b. 2018 pink salmon harvest forecast models from Southeast Alaska coastal monitoring surveys. NPAFC Doc. 1848. 19 pp. National Oceanic and Atmospheric Administration (NOAA), National Marine Fisheries Service (NMFS), Alaska Fisheries Science Center, Auke Bay Laboratories, and Alaska Department of Fish and Game (Available at <http://www.npafc.org>)
- Orsi, J. A., J. M. Murphy, and A. L. J. Brase. 1997. Survey of juvenile salmon in the marine waters of southeastern Alaska, May–August 1997. (NPAFC Doc. 277) Auke Bay Laboratory, Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA, U.S. Dept. Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626, USA. (Available at <http://www.npafc.org>)
- Orsi, J. A., M. V. Sturdevant, J. M. Murphy, D. G. Mortensen, and B. L. Wing. 2000. Seasonal habitat use and early marine ecology of juvenile Pacific salmon in southeastern Alaska. *North Pacific Anadromous Fish Commission Bulletin* 2: 111–122. (Available at www.npafc.org)
- Orsi, J. A., M. V. Sturdevant, A. C. Wertheimer, B. L. Wing, J. M. Murphy, D. G. Mortensen, E. A. Fergusson, and B. K. Krauss. 2001. Survey of juvenile salmon in the marine waters of southeastern Alaska, May–September 2000. (NPAFC Doc. 536) Auke Bay Laboratory, Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA, U.S. Department of Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626, USA. (Available at www.npafc.org)
- Orsi, J. A., E. A. Fergusson, M. V. Sturdevant, B. L. Wing, A. C. Wertheimer, and W. R. Heard. 2005. Survey of juvenile salmon and associated epipelagic ichthyofauna in the marine waters of southeastern Alaska, May–August 2004. (NPAFC Doc. 871) Auke Bay Laboratory, Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA, U.S. Department of Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626, USA. (Available at www.npafc.org)
- Orsi, J. A., E. A. Fergusson, M. V. Sturdevant, B. L. Wing, A. C. Wertheimer, and W. R. Heard. 2006. Annual survey of juvenile salmon and ecologically-related species in the marine waters of southeastern Alaska, May–August 2005. (NPAFC Doc. 955) Auke Bay Lab., Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA, 11305 Glacier Highway, Juneau, AK 99801-8626, USA. (Available at www.npafc.org)
- Orsi, J. A., E. A. Fergusson, M. V. Sturdevant, B. L. Wing, A. C. Wertheimer, and W. R. Heard. 2009. Annual survey of juvenile salmon, ecologically-related species, and environmental factors in the marine waters of southeastern Alaska, May–August 2008. (NPAFC Doc. 1181) Auke Bay Lab, Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA, 11305 Glacier Highway, Juneau, AK 99801-8626, USA. (Available at <http://www.npafc.org>)
- Orsi, J. A., E. A. Fergusson, M. V. Sturdevant, W. R. Heard, and E. V. Farley, Jr. 2012. Annual survey of juvenile salmon, ecologically-related species, and biophysical factors in the marine waters of southeastern Alaska, May–August 2011. (NPAFC Doc. 1428, Rev. 1) Auke Bay Lab., Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA, NMFS, 17109 Point Lena Loop Road, Juneau, 99801, USA. (Available at <http://www.npafc.org>)

- Parker, R. R. 1968. Marine mortality schedules of pink salmon of the Bella Coola River, central British Columbia. *Journal of the Fisheries Research Board of Canada* 25(4):757–794.
- Piston, A., and S. Heinl. 2014. Pages 50–54 [In] A. R. Munro, and C. Tide, editors. Run forecasts and harvest projections for 2014 Alaska salmon fisheries and review of the 2013 season. Alaska Department of Fish and Game, Special Publication No. 14-10, Anchorage.
- Piston, A., and S. Heinl. 2017. Pages 48–51 [In] R. E. Brenner, and A. R. Munro, editors. Run forecasts and harvest projections for 2017 Alaska salmon fisheries and review of the 2016 season. Alaska Department of Fish and Game, Special Publication No. 17-08, Anchorage.
- Piston, A. W., and S. C. Heinl. 2020. Pink salmon stock status and escapement goals in Southeast Alaska through 2019. Alaska Department of Fish and Game, Special Publication No. 20-09, Anchorage.
- Piston, A., S. Heinl, S. Miller, R. Brenner, J. Murphy, J. Watson, A. Gray, and E. Fergusson. 2019. Pages 46–49 [In] R. E. Brenner, A. R. Munro, and S. J. Larsen, editors. Run forecasts and harvest projections for 2019 Alaska salmon fisheries and review of the 2018 season. Alaska Department of Fish and Game, Special Publication No. 19-07, Anchorage.
- Piston, A., S. Heinl, S. Miller, R. Brenner, J. Murphy, J. Watson, A. Gray, and E. Fergusson. 2020. Pages 42–45 [In] R. E. Brenner, S. J. Larsen, A. R. Munro, and A. M. Carroll, editors. 2020. Run forecasts and harvest projections for 2020 Alaska salmon fisheries and review of the 2019 season. Alaska Department of Fish and Game, Special Publication No. 20-06, Anchorage.
- Piston, A., S. Heinl, S. Miller, R. Brenner, J. Murphy, J. Moss, W. Strasburger, E. Fergusson, and A. Gray. 2021a. Pages 42–45 [In] R. E. Brenner, S. J. Larsen, A. R. Munro, and A. M. Carroll, editors. Run forecasts and harvest projections for 2021 Alaska salmon fisheries and review of the 2020 season. Alaska Department of Fish and Game, Special Publication No. 21-07, Anchorage.
- Piston, A., S. Heinl, S. Miller, R. Brenner, J. Murphy, J. Moss, W. Strasburger, E. Fergusson, and A. Gray. 2022. Pages 45–48 [In] R. E. Brenner, S. J. Donnellan, and A. R. Munro, editors. Run forecasts and harvest projections for 2022 Alaska salmon fisheries and review of the 2021 season. Alaska Department of Fish and Game, Special Publication No. 22-11, Anchorage.
- Radchenko, V. 2020. Lost in the Ocean: Where have pink salmon been during our quest for salmon in the Gulf of Alaska? Pages 4–11 [In] North Pacific Anadromous Fish Commission Newsletter No. 47, January 2020.
- Ruggerone, G. T., M. Zimmermann, K. W. Myers, J. L. Nielsen, and D. E. Rogers. 2003. Competition between Asian pink salmon (*Oncorhynchus gorbuscha*) and Alaskan sockeye salmon (*O. nerka*) in the North Pacific Ocean. *Fisheries Oceanography* 12(3):209–219.
- Shevlyakov, E. A., and M. V. Koval. 2012. Forecast and production dynamics of the pink salmon of Kamchatka. North Pacific Anadromous Fish Commission Technical Report 8:121–125. (Available at www.npafc.org)
- Wertheimer, A.C., J.A. Orsi, E.A. Fergusson, and M. V. Sturdevant. 2009a. Forecasting pink salmon harvest in Southeast Alaska from juvenile salmon abundance and associated environmental parameters: 2008 returns and 2009 forecast. NPAFC Document 1202. (Available at <http://www.npafc.org>)
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2010a. Forecasting pink salmon harvest in Southeast Alaska from juvenile salmon abundance and associated environmental parameters: 2009 Harvest and 2010 Forecast. NPAFC Doc. 1278. (Available at <http://www.npafc.org>)
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2011. Forecasting pink salmon harvest in Southeast Alaska from juvenile salmon abundance and associated environmental parameters: 2010 Returns and 2011 Forecast. NPAFC Doc. 1343. (Available at <http://www.npafc.org>)
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2012. Forecasting pink salmon harvest in Southeast Alaska from juvenile salmon abundance and associated biophysical parameters: 2011 returns and 2012 forecast. NPAFC Doc. 1414, Rev. 1. (Available at <http://www.npafc.org>)

- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2013. Forecasting pink salmon harvest in Southeast Alaska from juvenile salmon abundance and associated environmental parameters: 2012 Returns and 2013 Forecast. NPAFC Doc. 1486 24 pp. (Available at <http://www.npafc.org>)
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2014. Forecasting pink salmon harvest in southeast Alaska from juvenile salmon abundance and associated biophysical parameters: 2013 returns and 2014 forecast. NPAFC Doc. 1555. (Available at <http://www.npafc.org>)
- Wertheimer, A. C., J. A. Orsi, and E. A. Fergusson. 2015. Forecasting pink salmon harvest in southeast Alaska from juvenile salmon abundance and associated biophysical parameters: 2014 returns and 2015 forecast. NPAFC Doc. 1618. (Available at <http://www.npafc.org>)
- Wertheimer, A.C., J.A. Orsi, and E.A. Fergusson. 2017. Forecasting pink salmon harvest in southeast Alaska from juvenile salmon abundance and associated biophysical parameters: 2015 returns and 2016 forecast. NPAFC Doc. 1740. (Available at <http://www.npafc.org>)
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and J. M. Murphy. 2018. Forecasting pink salmon harvest in Southeast Alaska from juvenile salmon abundance and associated biophysical parameters: 2016 returns and 2017 forecast. NPAFC Doc. 1772. (Available at <http://www.npafc.org>)
- Willette, T. M., R. T. Cooney, V. Patrick, D. M. Mason, G. I. Thomas, and D. Scheel. 2001. Ecological processes influencing mortality of juvenile pink salmon (*Oncorhynchus gorbuscha*) in Prince William Sound, Alaska. *Fisheries Oceanography* 10(1):14–41.