

2021 Preseason Pink Salmon Forecast

pink_cal_mixpool

Sara Miller, Rich Brenner, Jim Murphy

November 3, 2020—draft

Objective

To forecast the Southeast Alaska (SEAK) pink salmon harvest in 2021.

Executive Summary

Forecasts were developed using an approach described in Murphy et al. 2019. A multiple regression model was developed using monthly peak juvenile pink salmon CPUE (standardized catch based on 20 minute trawl set) for the June and July surveys and associated environmental parameters. The model used was:

$$E(y) = \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_1 X_2 + \epsilon,$$

where y is $\ln(\text{harvest})$, β_1 is the coefficient for $\ln(\text{CPUE}+1)$, β_2 is the coefficient for the environmental covariate water temperature (i.e., May through July average temperature in the upper 20 m at eight stations in Icy Strait; 'ISTI'), β_3 is the interaction term, and ϵ represents the normal error term that is lognormal.

Leave-one-out cross validation (hindcast) and model performance metrics such as Mean and Median Absolute Percentage Error (MAPE, MEAPE), and mean absolute scaled error (MASE) (Hyndman and Kohler 2006) were then used to evaluate forecast accuracy of alternative models. Statistical analyses were performed with the R Project for Statistical computing version 3.6.3 (R Core Team 2020).

Based on an additive model with CPUE and temperature, the SEAK pink salmon harvest in 2021 is predicted to be in the moderate range with a point estimate of 29.8 million fish (80% prediction interval: 20.1 to 44.1 million fish).

Analysis

Three hierarchical models were investigated. The full model was:

$$E(y) = \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_1 X_2,$$

where X_1 is $\ln(\text{CPUE}+1)$ and X_2 is the average temperature in Icy Strait in May, June, and July at eight stations in Icy Strait (Icy Strait and Upper Chatham transects), and β_3 is the interaction term between CPUE and the temperature indice. In the past, the ISTI variable was the average temperature in the upper 20 m during May through August at eight stations in Icy Strait (Icy Strait and Upper Chatham transects). The regression coefficients CPUE and temperature (ISTI) were significant in the first two models

(m1, m2). The interaction term was not significant in model m3 (Table 1). Therefore, only the first two models will be considered further.

Table 1: Parameter estimates for the three potential models.

model	term	estimate	std.error	statistic	p.value
m1	(Intercept)	2.280	0.204	11.169	0.000
m1	CPUE	0.460	0.073	6.317	0.000
m2	(Intercept)	6.793	0.971	6.997	0.000
m2	CPUE	0.521	0.053	9.814	0.000
m2	ISTI	-0.513	0.109	-4.701	0.000
m3	(Intercept)	4.001	2.565	1.560	0.135
m3	CPUE	1.664	0.976	1.706	0.104
m3	ISTI	-0.208	0.281	-0.739	0.469
m3	CPUE:ISTI	-0.124	0.106	-1.174	0.255

The model summary results using the metrics AICc, MAPE, MEAPE, and MASE (Hyndman and Kohler 2006) are shown in Table 2. For all of these metrics, the smallest value is the preferred model. The difference (Δ_i) between a given model and the model with the lowest AICc value and the metric MASE were the primary statistics for choosing appropriate models in this analysis. Models with $\text{AICc}\Delta_i \leq 2$ have substantial support, those in which $4 \leq \text{AICc}\Delta_i \leq 7$ have considerably less support, and models with $\text{AICc}\Delta_i > 10$ have essentially no support (Burnham and Anderson 2004). These two metrics (AICc, MASE) suggest that model m2 is the preferred models. If temperature is actually altering how CPUE is related to abundance it makes sense to restrict the temperature data to the CPUE months in the forecast model (June and July). The month of May is included as there are important migratory dynamics prior to the time juveniles are actually sampled in Icy Strait. Model m2 (based on average temperature in May through July) was used to forecast the 2021 pink salmon harvest.

Table 2: Summary of model outputs and forecast error measures

model	AdjR2	AICc	MAPE	MEAPE	MASE
CPUE	0.639	29	0.111	0.09	0.379
CPUE+ISTI	0.820	15	0.077	0.06	0.256




Figure 1: Relationship between a) $\ln(\text{CPUE}+1)$ and $\ln(\text{harvest})$ and b) temperature in May through July (ISTI) and $\ln(\text{harvest})$.

Results

The best regression model based on AICc, the MASE metric, significant coefficients in the models, and the argument for restricting temperature to the months when CPUE is sampled was model m2 (i.e. the model containing CPUE, and a May through July temperature variable). The adjusted R^2 value for model m2 was 0.82 indicating overall a good model fit.

Conclusion

The SEAK pink salmon harvest in 2021 is predicted to be in the moderate range with a point estimate of 29.8 million fish (80% prediction interval: 20.1 to 44.1 million fish).




Figure 2: SEAK harvest (millions) a) by year and b) by the fitted values from model m2. The line in figure b) is a one to one line. The predicted 2021 forecast is symbolized as a grey circle with an 80% prediction interval (20.1 to 44.1 million fish) in figure a).

Table 3: Detailed output for model m2. Juvenile year 2012 (year 2013) shows the largest standardized residual.

year	SEAKCatch_log	CPUE	resid	hat_values	Cooks_distance	std_resid	fitted
1998	3.750	2.477744	0.416	0.048	0.036	1.472	3.333
1999	4.354	5.622380	-0.499	0.343	0.785	-2.123	4.853
2000	3.011	1.597723	-0.223	0.097	0.024	-0.812	3.234
2001	4.205	3.729985	-0.039	0.121	0.001	-0.143	4.244
2002	3.813	2.868826	0.158	0.048	0.005	0.558	3.656
2003	3.961	2.784664	-0.077	0.167	0.006	-0.291	4.038
2004	3.813	3.077820	0.214	0.058	0.012	0.760	3.599
2005	4.079	3.899407	0.011	0.102	0.000	0.039	4.069
2006	2.451	2.040345	-0.170	0.244	0.049	-0.677	2.621
2007	3.802	2.572781	0.169	0.059	0.008	0.602	3.633
2008	2.766	1.167639	-0.043	0.103	0.001	-0.157	2.809
2009	3.638	2.323473	-0.300	0.231	0.140	-1.182	3.938
2010	3.178	2.094192	0.106	0.066	0.003	0.377	3.073

year	SEAKCatch_log	CPUE	resid	hat_values	Cooks_distance	std_resid	fitted
2011	4.076	3.687782	0.182	0.089	0.014	0.659	3.894
2012	3.059	1.305918	0.033	0.103	0.001	0.120	3.026
2013	4.551	3.161036	0.588	0.084	0.136	2.119	3.963
2014	3.616	1.923429	0.370	0.059	0.036	1.315	3.247
2015	3.558	3.426619	-0.354	0.070	0.040	-1.269	3.913
2016	2.912	2.201588	-0.079	0.096	0.003	-0.288	2.992
2017	3.547	3.905705	0.002	0.256	0.000	0.010	3.544
2018	2.092	0.310436	-0.472	0.212	0.301	-1.835	2.564
2019	3.049	1.171558	0.224	0.103	0.026	0.818	2.825
2020	2.088	1.141971	-0.216	0.242	0.078	-0.855	2.303

References

- Burnham, K. P., and Anderson, D. R. 2004. Multimodel inference: Understanding AIC and BIC in model selection. *Sociological Methods and Research* 33: 261-304.
- Hyndman, R. J. and A. B. Koehler. 2006. Another look at measures of forecast accuracy. *International Journal of Forecasting* 22: 679-688.
- Murphy, J. M., E. A. Fergusson, A. Piston, A. Gray, and E. Farley. 2019. Southeast Alaska pink salmon growth and harvest forecast models. *North Pacific Anadromous Fish Commisison Technical Report No. 15*: 75-81.
- R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <http://www.r-project.org/index.html>