



Nonlinear convection-diffusion model of lung capillary perfusion and gas exchange

Bastián Herrera Cuesta

Instituto de Ingeniería Biológica y Médica

Pontificia Universidad Católica de Chile

Santiago, Chile

Abstract

Pulmonary capillary perfusion and gas exchange are the fundamental physical processes that allow for respiration to occur at the microscale. Present-day computational simulations of these phenomena are often based on low-dimensional mathematical models on idealized alveolar geometries, where the chemical reactions between O_2 - CO_2 and haemoglobin are simplified [1, 2]. However, these models fail to capture the complex chemical reactions that take place in pulmonary capillary blood, and overlook the effect of porous alveolar structure. In our study, we develop a coupled perfusion and gas exchange model that reflects physiological gas and Hb dynamics in pulmonary capillaries. To this end, we formulate a system of two coupled nonlinear convection-diffusion boundary value problems. We derive the equations from continuum balance laws, incorporating an experimentally-validated relationship between gas partial pressures and Hb saturations through a fully coupled Hill equation-like model [3]. We numerically solve these problems using a finite element scheme in a simple slab capillary domain, on a range of blood flow velocity fields and blood pH values. Moreover, we perform numerical simulations in a 3D alveolar domain reconstructed from μ -CT rat lung images, which were morphologically manipulated in order to simulate various degrees of emphysema. Numerical perfusion experiments agree with expected blood pressure drops and velocity fields in the lung capillary domain [4]. Additionally, numerical gas exchange simulations reconstruct physiological O_2 and CO_2 partial pressure and Hb saturation dynamics in the capillary domain, and reproduce the ~ 100 ms equilibrium time of O_2 [5]. Further, these simulations replicate Bohr and Haldane effects in blood, and provide a fast and effective framework for estimating whole-lung diffusing capacities for O_2 and CO_2 , whose values show a marked decrease during emphysema. We envision that this model broadens the applicability of computational lung simulations in clinical settings, specially during exercise and pathological conditions that alter perfusion dynamics and the overall lung gas exchanging function.

References

- [1] ZURITA, PABLO; HURTADO, DANIEL, *Computational modeling of capillary perfusion and gas exchange in alveolar tissue*. Computer Methods in Applied Mechanics and Engineering, Vol. **399**. (2022).
- [2] STEWART, NEIL J.; PARRA-ROBLES, JUAN; WILD, JIM M., *Finite element modeling of ^{129}Xe diffusive gas exchange NMR in the human alveoli*, Journal of Magnetic Resonance **271**, (2016). 21-33.
- [3] DASH, RANJAN K.; KORMAN, BEN; BASSINGTHWAIGHTE, JAMES B., *Simple Accurate Mathematical Models of Blood HbO_2 and $HbCO_2$ Dissociation Curves at Varied Physiological Conditions: Evaluation and Comparison with Other Models*, European Journal of Applied Physiology **116**, (2016). 97-113.

- [4] WEST, JOHN B.; LUKS, ANDREW, *West's Respiratory Physiology: The Essentials*, 11th edition, Wolters Kluwer, (2021).
- [5] TABUCHI, ARATA; STYP-REKOWSKA, BEATA; SLUTSKY, ARTHUR S.; WAGNER, PETER D.; PRIES, AXEL R.; KUEBLER, WOLFGANG M., *Precapillary oxygenation contributes relevantly to gas exchange in the intact lung*, American Journal of Respiratory and Critical Care Medicine **188**, (2013). 474-481.