

COMPUTATIONAL PHYSICS

Problem Set 7

Patrick Anker

October 27, 2017

PROBLEM 1: *Analytically solve the heat equation*

$$\frac{\partial u(x, t)}{\partial t} = \alpha \frac{\partial^2 u(x, t)}{\partial x^2} \quad (1)$$

for a rod of length L with Neumann boundary conditions

$$\frac{\partial u(0, t)}{\partial x} = \frac{\partial u(L, t)}{\partial x} = 0$$

using the method of separation of variables.

For the separation of variables, we assume that $u(x, t)$ is of the form

$$u(x, t) = f(x)g(t)$$

By plugging this substitution into the 1D heat equation (Eq. 1), we get this expression:

$$f(x) \frac{\partial g}{\partial t} = \alpha g(t) \frac{\partial^2 f}{\partial x^2}$$

If we divide this equation by $u(x, t)$, the equation becomes a useful form:

$$\begin{aligned} \frac{1}{u} \left(f(x) \frac{\partial g}{\partial t} \right) &= \frac{1}{u} \left(\alpha g(t) \frac{\partial^2 f}{\partial x^2} \right) \\ \frac{\dot{g}}{g} &= \alpha \frac{f''}{f} \end{aligned} \quad (2)$$

where the dot implies a time derivative and the double prime implies two spacial derivatives. For this to be a physical solution – that is, if an expression that involves dimensions of time were to equal an expression that involves dimensions of space – the two sides of Eq. 2 must be constant:

$$\frac{\dot{g}}{g} = \alpha \frac{f''}{f} = c$$

We now can work with each side individually. The time-dependent function $g(t)$ is the easiest to work with; simply use the method “separate-integrate:”

$$\begin{aligned} \frac{dg/dt}{g} &= c \\ \frac{dg}{g} &= c dt \\ \Rightarrow g(t) &= G_0 e^{ct} \end{aligned}$$

Since the Neumann boundary conditions only apply to the spacial dependence, this form of $g(t)$ is completely specified. k will be determined via the spacial dependence:

$$\alpha \frac{f''}{f} = c$$

$$f'' = \frac{c}{\alpha} f$$

Solutions to this differential equation follow the form:

$$f(x) = Ae^{x\sqrt{c/\alpha}} + Be^{-x\sqrt{c/\alpha}}$$

From the boundary conditions, we can limit this further:

$$\frac{\partial u(0, t)}{\partial x} = \frac{\partial u(L, t)}{\partial x} = 0$$

$$g(t)f'(0) = g(t)f'(L) = 0$$

$$f'(0) = f'(L) = 0$$

$$\rightarrow A\sqrt{\frac{c}{\alpha}} - B\sqrt{\frac{c}{\alpha}} = 0 \Rightarrow A = B$$

Now let's consider the other boundary $f'(L) = 0$:

$$A\sqrt{\frac{c}{\alpha}}e^{L\sqrt{c/\alpha}} - A\sqrt{\frac{c}{\alpha}}e^{-L\sqrt{c/\alpha}} = 0$$

For this to work, either c is (trivially) 0 or c is negative; this would yield oscillatory behavior. Let $k^2 = \left|\frac{c}{\alpha}\right|$ and substitute in:

$$Ake^{ikL} - Ake^{-ikL} = 0$$

$$2iAk \sin(kL) = 0$$

$$\sin(kL) = 0$$

$$\rightarrow kL = n\pi$$

$$k^2 = \frac{n^2\pi^2}{L^2}$$

$$\Rightarrow c = -\frac{\alpha n^2\pi^2}{L^2}$$

This \sin showed up for the *derivative* of $f(x)$. Furthermore, this is just for *one* eigenvector. The total spectrum of solutions make up the open domain where $n \in (-\infty, \infty) \& n \in \mathbb{Z}$. This means the full solution $u(x, t)$ is:

$$u(x, t) = \sum_{n=-\infty}^{\infty} C_n \exp\left[-\frac{\alpha n^2 \pi^2}{L^2} t\right] \cos\left(\frac{n\pi}{L} x\right) \quad (3)$$

where C_n are just some constants that can be chosen at will.

* * *