


Identifying Models of Trait-Mediated Community Assembly

using random forests and
approximate Bayesian computation

Megan Ruffley, Katie Peterson,

Bob Week, David C. Tank,
Luke J. Harmon




- Megan Ruffley (she/her)
- University of Idaho
- Institute for Bioinformatics and Evolutionary Studies (**Ibest**)
- Bioinformatics and Computation Biology
- Plants
- Roller Derby



David C. Tank

Jack Sullivan

Luke J. Harmon



Interested in identifying neutral and non-neutral processes of assembly




“The processes that determine the patterns of the number and composition of co-occurring species have been central to community ecology for decades...” Chase 2003.

Plant successional theory

Clements and Gleason

Are communities a result of
deterministic (Clements 1938)
or *stochastic* (Gleason 1927)
processes?




Frederic Clements (1916)

“Succession is deterministic”

-stable ‘climax’ community that is in perfect balance with the climate

“Communities are super-organisms”

-Communities are dependent on the species interactions among members




FREDERICK EDWARD CLEMENTS
1874-1945




Henry Gleason (1917)

- “Each species is an independent entity”
 - distribution of each species is due to genetics, physiology, life history, and relations to abiotic and biotic factors
 - communities are just a consequence of independent interactions among species and their environment



(A) Groups of species are tightly associated, and are supplanted by other groups of tightly associated species. (B) Individual species independently respond to environmental conditions. Each curve on the graphs represents the abundance of a single species.



**note on
phylogeographic
concordance

(A) Groups of species are tightly associated, and are supplanted by other groups of tightly associated species. (B) Individual species independently respond to environmental conditions. Each curve on the graphs represents the abundance of a single species.

Assembly “rules”

- Jared Diamond 1975 – the community is a product of the species pool, the environment, and the interactions amongst species.
- Problem was, if all of these things seems similar, why were communities behaving differently? **cough** (gleason)

Assembly “rules”

- The whole debate begins again in a slightly different context.
- 1.) 1:1 match between species and environment, even if the timing of species entering a community is different
 - (Neill 1975, Tilman et al. 1986, Sommer 1991, Law & Morton 1996)
- 2.) depends on what species get there first (multiple state equilibria)
 - (Luh and Pimm 1993, Law 1999, Robinson and Dickerson 1988, Drake 1991, Samuels & Drake 1997)

Assembly “rules”

- The whole debate begins again in a slightly different context.
 - 1.) 1:1 match between species and environment
species entering a community.
 - (Neill 1975, Tilman 1982, Law 1999, Morton 1996)
 - 2.) different species get there first (multiple state equilibria)
 - (Law 1999, Tilman 1993, Law 1999, Robinson and Dickerson 1988, Drake 1991, Samuels & Drake 1997)
- **Evidence has been found for both arguments**


Measure Community Structure

- Species/genus ratios
 - Moreau 1948, Simberloff 1970
- Abundance distributions
 - Simpson 1949, Magurran 1988
- Higher Taxonomic diversity metrics
 - Faith 1992, Gotelli & Colwell 2001
- Functional traits information
 - MacArthur & Levins 1967; Weiher *et al.* 1999; McGill *et al.* 2006

Measure Community Structure

- Species/genus ratios
 - Moreau 1948, Simberloff 1970
- Abundance distributions
 - Simpson 1949, Magurran 1988
- Higher Taxonomic diversity metrics
 - Faith 1992, Gotelli & Colwell 2001
- Functional traits information
 - MacArthur & Levins 1967; Weiher *et al.* 1999; McGill *et al.* 2006

Null Models in Community Ecology (Gotelli & Graves)



Phylogenetic Community Ecology

- We should use phylogenies in community ecology!
 - Brooks & McLennan 1991, Losos 1996, Thompson et al. 2001
- “*...deterring by the unfamiliarity of molecular techniques and phylogenetic methods and the accompanying terminology*”


Phylogenetic Community Ecology

- The phylogeny encapsulates information about the functional traits important for assembly
 - **assumption
- Is the distribution of species in a community nonrandom with respect to phylogeny?

Phylogenetic Community Ecology

- Is the distribution of species in a community nonrandom with respect to phylogeny?


Habitat Filtering – species fail to establish in community due to incompatibility with relevant environmental factors (Bazzaz 1991)



Phylogenetic Community Ecology

- Is the distribution of species in a community nonrandom with respect to phylogeny?

Habitat Filtering – species fail to establish in community due to incompatibility with relevant environmental factors (Bazzaz 1991)




Competitive Exclusion – species fail to establish in a community when they cannot out-compete another species for their desired niche (MacArthur & Levins 1967)

Phylogenetic Community Ecology

Phylogenetic dispersion metrics are commonly used to detect signal for habitat filtering and competitive exclusion.

-Mean Phylogenetic Distance (MPD)





Assumption: functional traits of importance are phylogenetically conserved.

Regional Species Pool

Environmental Filter

Competitive Exclusion

Local Species Pool




Community Assembly Inference; dispersion metrics

- Phylogenetic Community Structure

Grandcolas 1998, Webb et al. 2000, 2002,
Kraft et al. 2007



- Trait Community Structure

Weiher et al. 1999, McGill et al. 2006,
Cornwell et al. 2006, Kraft et al. 2007




Traits are phylogenetically conserved

Habitat Filtering =
phylogenetic underdispersion





Traits are phylogenetically conserved

Habitat Filtering =
phylogenetic underdispersion




Competitive Exclusion =
phylogenetic overdispersion




Traits are phylogenetically conserved

**Habitat Filtering =
phylogenetic underdispersion**



**Competitive Exclusion =
phylogenetic overdispersion**




Traits are phylogenetically convergent

**Habitat Filtering =
phylogenetic overdispersion**


**Competitive Exclusion =
phylogenetic underdispersion**

Traits are phylogenetically conserved

**Habitat Filtering =
phylogenetic underdispersion**




**Competitive Exclusion =
phylogenetic overdispersion**




Traits are phylogenetically convergent

**Habitat Filtering =
phylogenetic overdispersion**



**Competitive Exclusion =
phylogenetic underdispersion**



Community Assembly Inference; dispersion metrics


- Phylogenetic Community Structure

Grandcolas 1998, Webb et al. 2000, 2002,
Kraft et al. 2007

- Trait Community Structure


Weiher et al. 1999, McGill et al. 2006,
Cornwell et al. 2006, Kraft et al. 2007

Weiher and Keddy 1995, Kraft et al.
2007, 2010, Cavender-Bares et al.
2009; Kemble 2009; Mayfield and
Levine 2010; Gerhold et al. 2015



Current Community Assembly Inference Methods

- Relying on often-violated assumptions
- Statistical significance != Biological significance
- Cannot simultaneously compare model support
- Not widely comparable across communities



CAMI development objectives

- 1. Implement a model-based inference procedure*

CAMI development objectives

1. *Implement a model-based inference procedure*
 - *Compare support for neutral and non-neutral models simultaneously*

CAMI development objectives

1. Implement a model-based inference procedure

- *Compare support for neutral and non-neutral models simultaneously*
- **Cannot write down a likelihood, so we will need to use an approximate approach based on simulations*

CAMI development objectives

1. *Implement a model-based inference procedure*
 - *Compare support for neutral and non-neutral models simultaneously*
***Cannot write down a likelihood, so we will need to use an approximate approach based on simulations*
2. *Parameterize the strength of the non-neutral assembly processes*

CAMI development objectives


1. Implement a model-based inference procedure

- *Compare support for neutral and non-neutral models simultaneously*
- ***Cannot write down a likelihood, so we will need to use an approximate approach based on simulations*

2. Parameterize the strength of the non-neutral assembly processes

Objective 1: Implement a model-based inference procedure


1.1 Regional community phylogeny: N, λ, μ




N	regional species pool
λ	speciation rate
μ	extinction rate
σ^2	rate of character change
α	strength of constraints (OU only)
n	local species pool
t_E	effect of environmental filtering
t_C	effect of competitive exclusions

Objective 1: Implement a model-based inference procedure

1.1 Regional community phylogeny: N, λ, μ




1.2 Trait evolution:
 σ^2, α



N	regional species pool
λ	speciation rate
μ	extinction rate
σ^2	rate of character change
α	strength of constraints (OU only)
n	local species pool
t_E	effect of environmental filtering
t_C	effect of competitive exclusions


Objective 1: Implement a model-based inference procedure

1.1 Regional community phylogeny: N, λ, μ



N	regional species pool
λ	speciation rate
μ	extinction rate
σ^2	rate of character change
α	strength of constraints (OU only)
n	local species pool
t_E	effect of environmental filtering
t_C	effect of competitive exclusions

1.2 Trait evolution: σ^2, α



1.3 Local community assembly: n, t_E, t_C


- Neutral
- Environmental Filtering
- Competitive Exclusion

Objective 2: Parameterize the strength of the non-neutral assembly processes


Objective 2: Parameterize the strength of the non-neutral assembly processes

Environmental Filtering

$$P = e^{-\frac{(Z_i - Z_j)^2}{\tau}}$$




Objective 2: Parameterize the strength of the non-neutral assembly processes




Objective 1: Implement a model-based inference procedure

1.1 Regional community phylogeny: N, λ, μ




N	regional species pool
λ	speciation rate
μ	extinction rate
σ^2	rate of character change
α	strength of constraints (OU only)
n	local species pool
t_E	effect of environmental filtering
t_C	effect of competitive exclusions

1.2 Trait evolution: σ^2, α



1.3 Local community assembly: n, t_E, t_C




$P > \text{unif}(0,1)$

- ✓
- ✗
- ✓
- ✗
- ✗
- ✓
- ✗

- Neutral
- Environmental Filtering
- Competitive Exclusion

Objective 1: Implement a model-based inference procedure



Simulation #	SS.1	SS.2	SS.3	SS.30	Model
1	29.3201204	0.047795881	2.202369339	0.024039567	-0.551807157	Neutral
2	58.01018705	0.05683677	4.296006276	0.014001377	-0.141285853	Neutral
3	28.35028897	0.057748823	2.565336481	0.02472052	0.543737366	Neutral
4	54.4519145	0.058161893	4.462506196	0.006577184	-0.297716652	Neutral
5	29.52415799	0.0449925	1.332280042	-0.027307941	-0.424597011	Neutral
6	13.8002455	0.043015337	3.827085961	0.052868925	0.00038215	Filtering
7	68.51513963	0.05221236	3.773466742	0.006122832	-0.373019012	Filtering
8	6.409247373	0.067539483	1.27370363	0.056550081	-0.419361409	Filtering
9	7.869379132	0.044893904	1.621625795	0.005944114	-0.225941885	Filtering
10	56.74215972	0.042376631	3.870139564	-0.010551762	1.612683176	Filtering
11	32.21489305	0.039088122	4.436192932	-0.005017495	-0.608370405	Competition
12	35.31406132	0.055897014	1.802319164	0.00665565	0.575928644	Competition
...	15.17747744	0.052302964	2.833102254	-0.007315425	-0.343133625	Competition
10,000	176.8646799	0.046759559	3.404803136	0.003965272	-0.492954288	Competition

Objective 1: Implement a model-based inference procedure

Model-selection using approximate approaches

- randomForests (*Breiman 2001, Breiman and Cutler 2007*)**
- Approximate Bayesian Computation (ABC; *Beaumont, et al. 2002, Csilléry, et al. 2012*)

<i>Simulation #</i>	<i>SS.1</i>	<i>SS.2</i>	<i>SS.3</i>	<i>SS.30</i>	<i>Model</i>
1	29.3201204	0.047795881	2.202369339	0.024039567	-0.551807157	Neutral
2	58.01018705	0.05683677	4.296006276	0.014001377	-0.141285853	Neutral
3	28.35028897	0.057748823	2.565336481	0.02472052	0.543737366	Neutral
4	54.4519145	0.058161893	4.462506196	0.006577184	-0.297716652	Neutral
5	29.52415799	0.0449925	1.332280042	-0.027307941	-0.424597011	Neutral
6	13.8002455	0.043015337	3.827085961	0.052868925	0.00038215	Filtering
7	68.51513963	0.05221236	3.773466742	0.006122832	-0.373019012	Filtering
8	6.409247373	0.067539483	1.27370363	0.056550081	-0.419361409	Filtering
9	7.869379132	0.044893904	1.621625795	0.005944114	-0.225941885	Filtering
10	56.74215972	0.042376631	3.870139564	-0.010551762	1.612683176	Filtering
11	32.21489305	0.039088122	4.436192932	-0.005017495	-0.608370405	Competition
12	35.31406132	0.055897014	1.802319164	0.00665565	0.575928644	Competition
...	15.17747744	0.052302964	2.833102254	-0.007315425	-0.343133625	Competition
10,000	176.8646799	0.046759559	3.404803136	0.003965272	-0.492954288	Competition

Results

- 1. Power Analysis*
- 2. Parameter Estimation*

Results

1. Power Analysis


- **Dispersion Metrics**

Phylogenetic data

Functional trait data

- **randomForest (RF)**
- **ABC**

Results: Power Analysis




Results

1. Power Analysis

2. Parameter Estimation

-Estimate τ_E and t_C

Results: Parameter Estimation




CAMI introductory tutorial

- compphylo.github.io/Oslo2019/CAMI_files/CAMI_1.html



Results: Lava-flow Islands in Craters of the Moon National Monument and Preserve



Katie Peterson
Dr. Christine Parent lab
University of Idaho



Results: Lava-flow Islands in Craters of the Moon National Monument and Preserve



Results: Lava-flow Islands in Craters of the Moon National Monument and Preserve



Smith, S. A., and J. W. Brown. 2018. Constructing a broadly inclusive seed plant phylogeny. *American Journal of Botany* 105(3): 302-314.




- Spermatophyta phylogeny
79,881 tips
- Discarded 79,768 tips (99.9%)

- 8 kipuka communities
- 113 species in regional community
- 18-60 species in local community



Try it with your community!

<https://github.com/ruffleymr/CommunityPhylogeneticExample>


Results: Lava-flow Islands in Craters of the Moon National Monument and Preserve




Maximum vegetative height



*Results: Lava-flow Islands in Craters of the Moon
National Monument and Preserve*



Results: Lava-flow Islands in Craters of the Moon National Monument and Preserve




CAMI tutorial 2: San Juan Islands

- compphylo.github.io/Oslo2019/CAMI_files/CAMI_2.html



Results: San Juan Islands in the Pacific Northwest

Diversity and Distributions, (*Diversity Distrib.*) (2016) **22**, 318–331




Deconstructing Darwin's Naturalization Conundrum in the San Juan Islands using community phylogenetics and functional traits

Hannah E. Marx^{1*}, David E. Giblin², Peter W. Dunwiddie² and David C. Tank¹




- 322 regional species
- 50 island communities
- 25-150 species in each
- 3 traits


Results: San Juan Islands in the Pacific Northwest





Results: San Juan Islands in the Pacific Northwest



Results: San Juan Islands in the Pacific Northwest



Results: San Juan Islands in the Pacific Northwest



CAMI Conclusions

- Simulate phylogenetic and phenotypic community assembly data under neutral and non-neutral models of assembly
- Implements a model-based inference procedure with parameter estimation

**CAMI: Community Assembly
Model Inference**

R package and docs available now @
<https://github.com/ruffleymr/CAMI>

CAMI: Community Assembly Model Inference

R package and docs available now @ www.github.com/ruffleymr/CAMI

Received: 2 January 2017 | Accepted: 21 July 2017
DOI: 10.1111/2041-210X.12918

RESEARCH ARTICLE

Methods in Ecology and Evolution  BRITISH ECOLOGICAL SOCIETY

ecolottery: Simulating and assessing community assembly with environmental filtering and neutral dynamics in R

François Munoz¹  | Matthias Grenié² | Pierre Denelle² | Adrien Taudière² |
Fabien Laroche^{2,3} | Caroline Tucker^{2,4} | Cyrille Violle² 

APPLICATION

VirtualCom: a simulation model for eco-evolutionary community assembly and invasion

Tamara Münkemüller^{1*} and Laure Gallien^{1,2,3}



Article |  Full Access |

A new modeling approach estimates the relative importance of different community assembly processes

Fons van der Plas , Thijs Janzen, Alejandro Ordóñez, Wimke Fokkema ... See all authors 

REVIEW

Methods in Ec

Inferring community assembly processes from macroscopic patterns using dynamic eco-evolutionary models and Approximate Bayesian Computation (ABC)

Mikael Pontarp^{1,2,3}  | Åke Bränström^{4,5} | Owen L. Petchey² 

Identifying Models of Trait-Mediated Community Assembly

using random forests and
approximate Bayesian computation

Megan Ruffley, Katie Peterson,

Bob Week, David C. Tank,
Luke J. Harmon

