

Information Compression #6

Basics of Krylov subspace methods

Youhei Yamaji

Department of Applied Physics, The University of Tokyo

1. The largest and smallest eigenvalues
2. Sparse matrix generated by Hamiltonian
3. Krylov subspace method



Classification of Information Compression in Linear Algebra by Memory Costs

(1) A matrix can be stored

- SVD for dense matrix
- Compressed sensing (so far)

(2) Although a matrix cannot be stored, vectors can be stored

- SVD for sparse matrix
- Krylov subspace method

(3) A vector cannot be stored

- Matrix product/tensornetwork states

This Week's Information Compression Algorithm

Main focus:

Algorithms that calculate
specified eigenvalues and eigenvectors
of huge* *sparse* matrices

*You may not store your matrix A or
you may not pay $O(L^3)$ * cost

$$A \in \mathbb{R}^{L \times L}$$

Especially the largest and smallest eigenstates

Largest and Smallest Eigenvalues

1. Ground state of quantum many-body system


$$\langle O \rangle = \frac{\vec{u}^\dagger O \vec{u}}{\vec{u}^\dagger \vec{u}}$$

The ground state is important:

- Room temperature is often enough low
and well described by zero-temperature wave function
- Interest in ground states (at zero temperature)
 - Low-temperature phase such as superfluid phase
 - Zero-temperature phase transitions
(quantum phase transition)

Low-Temperature Phases

Phase diagram of ^3He



D. D. Osheroff, R. C. Richardson, and D. M. Lee,
Phys. Rev. Lett. 28, 885 (1972).

Erkki Thuneberg

<http://ltl.tkk.fi/research/theory/helium.html>

Largest and Smallest Eigenvalues

2. Principle component analysis for huge data

Eigenvalue problem of covariance matrices


K. Pearson,
Philosophical Magazine 2, 559 (1901)

$$\begin{bmatrix} \sum_{\ell} (x - \bar{x})^2 & \sum_{\ell} (x - \bar{x})(y - \bar{y}) \\ \sum_{\ell} (y - \bar{y})(x - \bar{x}) & \sum_{\ell} (y - \bar{y})^2 \end{bmatrix}$$

Largest and Smallest Eigenvalues

2. Principle component analysis for huge data

K. Pearson, Philosophical Magazine 2, 559 (1901)



Higher dimension
→ Data compression

Category of Numerical Linear Algebra

You need to choose algorithm depending on whether your matrix is 1) sparse/dense and
2) stored/not stored in memory

For a matrix that is dense and stored, you can find standard subroutines with $O(L^3)^*$ cost in LAPACK

* L is the linear dimension of your matrix A

$$A \in \mathbb{R}^{L \times L}$$

Largest and Smallest Eigenvalues

Ground state of quantum many-body system

Typically, sparse and not stored

Principle component analysis for huge data

Eigenvalue problem of covariance matrices

Dense/sparse and stored/not stored

-Partial SVD/low-rank approximation


will discussed in 8th lecture

Sparse Matrix Generated by Hamiltonian

Quantum Many-Body Problems

Quantum dots

F. R. Braakman, et al., Nat. Nano. 8, 432 (2013)




Quantum dot:

- A quantum box can confine a single electron
- Utilized for single electron transistor, quantum computers

Three-body problem:

→ Number of states = 2^3 (factor 2 from spin)



States represented by
superposition

$$\mathcal{F} = \left\{ \sum_{n_0=0,1} \sum_{n_1=0,1} \sum_{n_2=0,1} C_{n_0 n_1 n_2} |n_0\rangle \otimes |n_1\rangle \otimes |n_2\rangle : C_{n_0 n_1 n_2} \in \mathbb{C} \right\}$$

Quantum Many-Body Problems

Mutual Interactions



1. Operators acting on a single qubit

A two dimensional representation of Lie algebra SU(2)

$$[\hat{S}_j^x, \hat{S}_j^y] = i\hat{S}_j^z$$

$$[\hat{S}_j^y, \hat{S}_j^z] = i\hat{S}_j^x$$

$$[\hat{S}_j^z, \hat{S}_j^x] = i\hat{S}_j^y$$

-Commutator $[\hat{A}, \hat{B}] \equiv \hat{A}\hat{B} - \hat{B}\hat{A}$

$$\hat{S}_j^x |0\rangle = \frac{1}{2} |1\rangle$$

$$\hat{S}_j^x |1\rangle = \frac{1}{2} |0\rangle$$

$$\hat{S}_j^y |0\rangle = \frac{i}{2} |1\rangle$$

$$\hat{S}_j^y |1\rangle = -\frac{i}{2} |0\rangle$$

$$\hat{S}_j^z |1\rangle = \frac{1}{2} |1\rangle$$

$$\hat{S}_j^z |0\rangle = -\frac{1}{2} |0\rangle$$

Quantum Many-Body Problems

Mutual Interactions



Fock space of N qubits:

$$\mathcal{F} = \left\{ \sum_{n_0=0,1} \sum_{n_1=0,1} \cdots \sum_{n_{N-1}=0,1} C_{n_0 n_1 \cdots n_{N-1}} |n_0\rangle \otimes |n_1\rangle \otimes \cdots \otimes |n_{N-1}\rangle \right\}$$

$(C_{n_0 n_1 \cdots n_{N-1}} \in \mathbb{C})$

2. Operators acting on N-qubit Fock space:

$$\hat{S}_j^a, \hat{S}_j^a \hat{S}_{j+1}^a : \mathcal{F} \rightarrow \mathcal{F}$$

$$\hat{S}_j^a \doteq \overbrace{1 \otimes \cdots \otimes 1}^{j-1} \otimes \hat{S}_j^a \otimes \overbrace{1 \otimes \cdots \otimes 1}^{N-j}$$

$$\hat{S}_j^a \hat{S}_{j+1}^a \doteq \overbrace{1 \otimes \cdots \otimes 1}^{j-1} \otimes \hat{S}_j^a \otimes \hat{S}_{j+1}^a \otimes \overbrace{1 \otimes \cdots \otimes 1}^{N-j-1}$$

Quantum Many-Body Problems

Quantum entanglement

Example: Two qubits



- Superposition
- Utilized for quantum teleportation
cf.) EPR “paradox”

Mutual interactions between two qubits

$$\hat{H} = J \sum_{a=x,y,z} \hat{S}_0^a \hat{S}_1^a \quad (J \in \mathbb{R}, J > 0)$$

→ Superposition



$$|1\rangle \otimes |0\rangle - |0\rangle \otimes |1\rangle$$

Hamiltonian Matrix

Example: N qubits



N-qubit Fock space:

$$\mathcal{F} = \left\{ \sum_{n_0=0,1} \sum_{n_1=0,1} \cdots \sum_{n_{N-1}=0,1} C_{n_0 n_1 \cdots n_{N-1}} |n_0\rangle \otimes |n_1\rangle \otimes \cdots \otimes |n_{N-1}\rangle \right\}$$

$(C_{n_0 n_1 \cdots n_{N-1}} \in \mathbb{C})$

Mutual interactions among N qubits:

Hamiltonian operator

$$\hat{H} : \mathcal{F} \rightarrow \mathcal{F}$$

$$\hat{H} = J \sum_{j=0}^{N-1} \sum_{a=x,y,z} \hat{S}_j^a \hat{S}_{\text{mod}(j+1,N)}^a$$

Vectors in Fock Space

Correspondence between spin and bit

$$\begin{aligned} |\uparrow\rangle &= |1\rangle \\ |\downarrow\rangle &= |0\rangle \end{aligned}$$

2^N -dimensional Fock space:

$$\mathcal{F} = \left\{ \sum_{n_0=0,1} \sum_{n_1=0,1} \cdots \sum_{n_{N-1}=0,1} C_{n_0 n_1 \cdots n_{N-1}} |n_0\rangle \otimes |n_1\rangle \otimes \cdots \otimes |n_{N-1}\rangle \right\} \\ (C_{n_0 n_1 \cdots n_{N-1}} \in \mathbb{C})$$

Decimal representation of orthonormalized basis

$$|I\rangle_d = |n_0\rangle \otimes |n_1\rangle \otimes |n_2\rangle \otimes \cdots \otimes |n_{N-1}\rangle \quad I = \sum_{\nu=0}^{N-1} n_\nu \cdot 2^\nu$$

Wave function as a vector

$$|\phi\rangle = \sum_{n_0=0}^1 \sum_{n_1=0}^1 \cdots \sum_{n_{N-1}=0}^1 C_{n_0 n_1 \cdots n_{N-1}} |n_0\rangle \otimes |n_1\rangle \otimes \cdots \otimes |n_{N-1}\rangle$$

$$v(I) = C_{n_0 n_1 \cdots n_{N-1}} \quad v(0 : 2^N - 1)$$

Vectors and Matrices in Fock Space

Inner product of vectors

$$\begin{aligned} & (\langle n_0| \otimes \langle n_1| \otimes \cdots \otimes \langle n_{N-1}|) \times (|n'_0\rangle \otimes |n'_1\rangle \otimes \cdots \otimes |n'_{N-1}\rangle) \\ &= \langle n_0|n'_0\rangle \times \langle n_1|n'_1\rangle \times \cdots \times \langle n_{N-1}|n'_{N-1}\rangle \end{aligned}$$

$$\langle n| \times |n'\rangle = \langle n|n'\rangle = \delta_{n,n'}$$

$$\langle \phi' | \phi \rangle = \sum_{n_0=0}^1 \sum_{n_1=0}^1 \cdots \sum_{n_{N-1}=0}^1 C'^*_{n_0 n_1 \cdots n_{N-1}} C_{n_0 n_1 \cdots n_{N-1}}$$

$$|\phi'\rangle = \sum_{n_0=0}^1 \sum_{n_1=0}^1 \cdots \sum_{n_{N-1}=0}^1 C'_{n_0 n_1 \cdots n_{N-1}} |n_0\rangle \otimes |n_1\rangle \otimes \cdots \otimes |n_{N-1}\rangle$$

$$|\phi\rangle = \sum_{n_0=0}^1 \sum_{n_1=0}^1 \cdots \sum_{n_{N-1}=0}^1 C_{n_0 n_1 \cdots n_{N-1}} |n_0\rangle \otimes |n_1\rangle \otimes \cdots \otimes |n_{N-1}\rangle$$

Hamiltonian matrix

$$H_{II'} = \langle I | \hat{H} | I' \rangle$$

Orthonormalized basis: $|I\rangle, |I'\rangle \in \mathcal{F} \quad \langle I | I' \rangle = \delta_{I,I'}$

Sparse Matrix

- Particle or orbital number: N
 - Fock space dimension: $\exp[N \times \text{const.}]$
 - # of terms in Hamiltonian: Polynomial of N
- # of matrix elements of Hamiltonian matrix:
(Polynomial of N) $\times \exp[N \times \text{const.}]$

For sufficiently large N ,
(Polynomial of N) $\times \exp[N \times \text{const.}]$
 $\ll (\exp[N \times \text{const.}])^2$

Then, the Hamiltonian matrix is **sparse**

An Example of Hamiltonian Matrix

$$\hat{H} = J \sum_{i=0}^{N-1} \hat{S}_i^z \hat{S}_{i+1}^z - \Gamma \sum_{i=0}^{N-1} \hat{S}_i^x$$

-Non-commutative

$$\left[\sum_{i=0}^{N-1} \hat{S}_i^z \hat{S}_{i+1}^z, \sum_{i=0}^{N-1} \hat{S}_i^x \right] \neq 0$$


→ Quantum fluctuations
or Zero point motion

-Sparse

of elements $\propto O(2^N)$

-Solvable

-Hierarchical matrix?



Computational and Memory Costs

Matrix-vector product of dense matrix

$$v_i = \sum_{j=0}^{N_H-1} A_{ij} u_j$$

Computational: $O((\text{Fock space dimension})^2)$

Memory: $O((\text{Fock space dimension})^2)$


Matrix-vector product of
large and sparse matrix

Computational: $O(\text{Fock space dimension})$

Memory: $O(\text{Fock space dimension})$

Hamiltonian is not stored in memory

Krylov Subspace Method for Sparse and Huge Matrices



Alexey Krylov

Aleksey Nikolaevich Krylov

1863-1945

Russian naval engineer and applied mathematician

Krylov subspace

$$A \in \mathbb{C}^{L \times L}$$

$$\mathcal{K}_n(A, \vec{b}) = \text{span}\{\vec{b}, A\vec{b}, \dots, A^{n-1}\vec{b}\}$$

Numerical cost to construct K_n : $\mathcal{O}(\text{nnz}(A) \times n)$

Numerical cost to orthogonalize K_n : $\mathcal{O}(L \times n^2)$

Cornelius Lanczos 1950

Walter Edwin Arnoldi 1951

*nnz: Number of non-zero
entries/elements

An Algorithm for Eigenvalue Problems of Large & Sparse Matrix: Power Method

Min. Eigenvalue of hermitian

Initial vector: $|v_1\rangle = \sum_{n=0} c_n |n\rangle$

Parameter: $\max_n \{E_n\} \leq \Lambda$

$$\hat{H}|n\rangle = E_n|n\rangle$$

$$\langle n'|n\rangle = \delta_{n',n}$$

$$E_0 \leq E_1 \leq \dots$$

$$\lim_{m \rightarrow +\infty} \frac{(\Lambda - \hat{H})^m |v_1\rangle}{\sqrt{\langle v_1 | (\Lambda - \hat{H})^{2m} | v_1 \rangle}} = |0\rangle$$

$$(\Lambda - \hat{H})^m |v_1\rangle = \sum_n (\Lambda - E_n)^m c_n |n\rangle$$

$$\lim_{m \rightarrow +\infty} \frac{\sum_{n>0} (\Lambda - E_n)^{2m} |c_n|^2}{(\Lambda - E_0)^{2m} |c_0|^2} = 0$$

Advanced Algorithm: Krylov Subspace Method

Krylov subspace method:

Finding approximate eigenstates in a Krylov subspace

$$\mathcal{K}_m(\hat{H}, |v_1\rangle) = \text{span}\{|v_1\rangle, \hat{H}|v_1\rangle, \dots, \hat{H}^{m-1}|v_1\rangle\}$$

Construction and orthogonalization of Krylov subspaces

Shift invariance:

$$\mathcal{K}_m(\hat{H}, |v_1\rangle) = \mathcal{K}_m(\hat{H} + z\mathbf{1}, |v_1\rangle)$$

Krylov subspace method:

- Lanczos method (symmetric/hermitian),
Arnoldi method (general matrix)
- Conjugate gradient method (CG method)
(many variation)

Lanczos Method

Initial : $\beta_1 = 0, |v_0\rangle = 0$

for $j = 1, 2, \dots, m$ **do**

$$|w_j\rangle = \hat{H}|v_j\rangle - \beta_j|v_{j-1}\rangle$$

$$\alpha_j = \langle w_j | v_j \rangle$$

$$|w_j\rangle \leftarrow |w_j\rangle - \alpha_j|v_j\rangle$$

$$\beta_{j+1} = \sqrt{\langle w_j | w_j \rangle}$$

$$|v_{j+1}\rangle = |w_j\rangle / \beta_{j+1}$$

Lanczos Method

$$\alpha_j = \langle v_j | \hat{H} | v_j \rangle$$

$$\beta_j = \langle v_{j-1} | \hat{H} | v_j \rangle = \langle v_j | \hat{H} | v_{j-1} \rangle$$

Orthogonalization

$$|v_j\rangle = \frac{\hat{H}|v_{j-1}\rangle - \sum_{\ell=1}^{j-1} |v_\ell\rangle \langle v_\ell | \hat{H} | v_{j-1} \rangle}{\langle v_j | \hat{H} | v_{j-1} \rangle}$$

$$\langle v_\ell | \hat{H} | v_{j-1} \rangle = \begin{cases} 0 & (\ell \leq j-3) \\ \beta_{j-1} & (\ell = j-2) \\ \alpha_{j-1} & (\ell = j-1) \end{cases}$$

Lanczos Method

Initial : $\beta_1 = 0, |v_0\rangle = 0$

for $j = 1, 2, \dots, m$ **do**

$$|w_j\rangle = \hat{H}|v_j\rangle - \beta_j|v_{j-1}\rangle$$

$$\alpha_j = \langle w_j | v_j \rangle$$

$$|w_j\rangle \leftarrow |w_j\rangle - \alpha_j|v_j\rangle$$

$$\beta_{j+1} = \sqrt{\langle w_j | w_j \rangle}$$

$$|v_{j+1}\rangle = |w_j\rangle / \beta_{j+1}$$

Lanczos Method

$$\alpha_j = \langle v_j | \hat{H} | v_j \rangle$$

$$\langle v_j | v_k \rangle = \delta_{j,k}$$

$$\beta_j = \langle v_{j-1} | \hat{H} | v_j \rangle = \langle v_j | \hat{H} | v_{j-1} \rangle$$

Hamiltonian projected onto m D Krylov subspace

$$H_m = \begin{pmatrix} \alpha_1 & \beta_2 & & & & 0 \\ \beta_2 & \alpha_2 & \beta_3 & & & \\ & \beta_3 & \alpha_3 & \ddots & & \\ & & \ddots & \ddots & \beta_{m-1} & \\ & & & \beta_{m-1} & \alpha_{m-1} & \beta_m \\ 0 & & & & \beta_m & \alpha_m \end{pmatrix}$$

Eigenvalues of projected Hamiltonian

→ Approximate eigenvalues of original Hamiltonian

Lanczos Method: # of Vectors Required

Initial : $\beta_1 = 0, |v_0\rangle = 0$

for $j = 1, 2, \dots, m$ **do**

$$|w_j\rangle \leftarrow \hat{H}|v_j\rangle - \beta_j|v_{j-1}\rangle$$

$$\alpha_j = \langle w_j | v_j \rangle$$

$$|w_j\rangle \leftarrow |w_j\rangle - \alpha_j|v_j\rangle$$

$$\beta_{j+1} = \sqrt{\langle w_j | w_j \rangle}$$

$$|v_{j+1}\rangle = |w_j\rangle / \beta_{j+1}$$

$$|v_{j-1}\rangle \rightarrow |w_j\rangle, |v_j\rangle$$

$$|w_j\rangle, |v_j\rangle$$

$$|w_j\rangle, |v_j\rangle$$

$$|w_j\rangle, |v_j\rangle$$

$$|w_j\rangle \rightarrow |v_{j+1}\rangle, |v_j\rangle$$

Convergence of Lanczos Method

Yousef Saad,

Numerical Methods for Large Eigenvalue Problems (2nd ed)

The Society for Industrial and Applied Mathematics 2011

Assumption: $\lambda_1 > \lambda_2 > \dots > \lambda_n$

Convergence theorem for the largest eigenvalue

$$0 \leq \lambda_1 - \lambda_1^{(m)} \leq (\lambda_1 - \lambda_n) \left[\frac{\tan \theta(|v_1\rangle, |0\rangle)}{C_{m-1}(1 + 2\gamma_1)} \right]^2$$
$$\sim 4(\lambda_1 - \lambda_n) [\tan \theta(|v_1\rangle, |0\rangle)]^2 e^{-4\sqrt{\gamma_1}m}$$

$$\gamma_1 = \frac{\lambda_1 - \lambda_2}{\lambda_2 - \lambda_n}$$

$$C_k(t) = \frac{1}{2} \left[\left(t + \sqrt{t^2 - 1} \right)^k + \left(t + \sqrt{t^2 - 1} \right)^{-k} \right]$$

Distribution of Eigenvalues of Hermitian Matrices

An important relationship between distribution or density of states and statistical mechanics

$$P(E) = \frac{\rho(E)e^{-\beta E}}{\int dE' \rho(E')e^{-\beta E'}} \sim \frac{\exp[-(E - \langle E \rangle)^2 / 2CT^2]}{\sqrt{2\pi CT^2}}$$

$$k_B = 1$$

$$C = \frac{\langle \hat{H}^2 \rangle - \langle \hat{H} \rangle^2}{T^2}$$


$$\langle \hat{H}^m \rangle \sim \int E^m P(E) dE$$

An Example of Density of State

24 site cluster of Kitaev model
(frustrated $S=1/2$ spins)

A. Kitaev, Annals Phys. 321, 2 (2006).

$$2^{24} = 16,777,216$$



Example of Dense Matrix: Random Symmetric Matrices

Eugene P. Wigner, Annals of Mathematics, 2nd Series, 67, 325 (1958)

Wigner's random matrix $(A)_{ij} = a_{ij}$

$$\underline{a_{ij} = a_{ji}} \quad (\text{Not necessarily sparse})$$

$$\int p_{ij}(a) da = 1$$

$$p_{ij}(+a) = p_{ij}(-a)$$

$$\langle a_{ij}^n \rangle = \int p_{ij}(a) a^n da \leq B_n$$

$$\langle a_{ij}^2 \rangle = \int p_{ij}(a) a^2 da = 1$$

Example of Dense Matrix: Random Symmetric Matrices

Eugene P. Wigner, Annals of Mathematics, 2nd Series, 67, 325 (1958)

Density of states of $L \times L$ symmetric random matrix

$$A\vec{v} = E\vec{v}$$

$$\sigma(E) = \begin{cases} \frac{\sqrt{4L - E^2}}{2\pi L} & (E^2 < 4L) \\ 0 & (E^2 > 4L) \end{cases}$$

Comment:

Sparse matrices in quantum many-body problems show smaller density of states than random matrices around the both ends of the distribution

- Sparse around maximum/minimum eigenvalues
- Lanczos method may work well

Approximate SVD by Krylov Subspace Method

Low-rank approximation by *block* Krylov subspace

C. Musco & C. Musco,
NIPS'15 Proceedings of 28th International Conference on
Neural Information Processing Systems 1, 1396 (2015)

$$\|A - ZZ^T A\|_2 \leq (1 + \epsilon) \|A - A_k\|_2 \quad \text{Operator norm defined by 2-norm (Spectral norm)}$$

$$A \in \mathbb{R}^{L \times M} \quad Z \in \mathbb{R}^{L \times k} \quad \text{rank } k \leq L, M$$

$$q = \mathcal{O}(\ln d / \sqrt{\epsilon})$$

random matrix $\Pi \in \mathbb{R}^{M \times k}$

$$\mathcal{K}_{q+1} = \text{span}\{A\Pi, (AA^T)A\Pi, \dots, (AA^T)^q A\Pi\}$$

$$Q \in \mathbb{R}^{N \times qk} \quad \text{Orthogonalized basis set of the block Krylov subspace}$$

$$M = Q^T A A^T Q \in \mathbb{R}^{qk \times qk}$$

U_k : the top k singular vectors of M

$$Z = QU_k$$

$(\Pi)_{ij}$: Random number generated by $e^{-x^2/2} / \sqrt{\pi}$

Next Week

第7回: 物質科学における情報圧縮

7 Information compression in condensed matter physics

Important References

Yousef Saad,
Numerical Methods for Large Eigenvalue Problems (2nd ed)
The Society for Industrial and Applied Mathematics 2011

Next Week

- 第1回： 現代物理学における巨大なデータ
- 第2回： 現代物理学と情報圧縮
- 第3回： 情報圧縮の数理1 (線形代数の復習)
- 第4回： 情報圧縮の数理2 (特異値分解と低ランク近似)
- 第5回： 情報圧縮の数理3 (スパース・モデリングの基礎)
- 第6回： 情報圧縮の数理4 (クリロフ部分空間法の基礎)

11/15 第7回： 物質科学における情報圧縮

11/15 #7 Information compression in materials science

- 第8回： データ解析の高速化：スパース・モデリングの物質科学への応用
- 第9回： データ空間の圧縮：クリロフ部分空間法の物質科学への応用
- 第10回： 高度なデータ圧縮：情報のエンタングルメントと行列積表現
- 第11回： 行列積表現の固有値問題への応用
- 第12回： テンソルネットワーク表現への発展
- 第13回： テンソルネットワーク繰り込みによる情報圧縮