




Dynamic causal modelling for fMRI

Jakob Heinze
Translational Neuromodeling Unit (TNU),
Institute for Biomedical Engineering
University and ETH Zürich

CP Course 2019, Zürich, Switzerland



Specialisation vs. Integration

Functional Specialisation

 u_1  $u_1 \times u_2$


«**Where**, in the brain, did my experimental manipulation have an effect?»

Functional Integration



«**How** did my experimental manipulation propagate through the network?»

Structural, functional & effective connectivity



anatomical/structural


- presence of physical connections

→ *DWI, tractography, tracer studies (animals)*

functional

- statistical dependency between regional time series

→ *correlations, ICA*



Sporns 2007, Scholarpedia

effective

- direct influences between neuronal populations


→ *DCM*

Context-independent

Mechanism - free

Mechanistic

A reminder – generative models



Observed data (fMRI)


$$y = g(x, \theta) + \varepsilon$$

Hidden states (Brain activity)

$$\frac{dx}{dt} = f(x, u, \theta) + \omega$$

Dynamic causal modelling

EEG,
MEG




Model inversion:
Estimating
neuronal
mechanisms


Forward model:
Predicting
measured activity

$$y = g(x, \theta) + \varepsilon$$

DCM for EEG
→ Next lecture
→ Dario Schöbi



State equation:
Describing neuronal
dynamics (and
hemodynamics)



$$\frac{dx}{dt} = f(x, u, \theta) + \omega$$



University of
Zurich^{UZH}



Translational Neuromodeling Unit

ETH zürich

Dynamic causal modelling



ACADEMIC
PRESS

Available online at www.sciencedirect.com



NeuroImage 19 (2003) 1273–1302

NeuroImage

www.elsevier.com/locate/ynimng


Dynamic causal modelling

K.J. Friston,* L. Harrison, and W. Penny


The Wellcome Department of Imaging Neuroscience, Institute of Neurology, Queen Square, London WC1N 3BG, UK

Received 18 October 2002; revised 7 March 2003; accepted 2 April 2003

DCM for fMRI - Overview



DCM for fMRI - Overview




Neuronal state equations


$$\frac{dx}{dt} = f(x, u) \approx f(x_0, 0) + \frac{\partial f}{\partial x} x + \frac{\partial f}{\partial u} u + \frac{\partial^2 f}{\partial x \partial u} u x + \frac{\partial^2 f}{\partial x^2} \frac{x^2}{2} + \dots$$

A C B D

bilinear model




nonlinear model





Neuronal state equations

DCM effective connectivity parameters are rate constants



$$\frac{dx_1}{dt} = a_{11}x_1 \longrightarrow$$


$$x_1(t) = x_1(0) \cdot \exp(a_{11}t)$$



If $x_1 \rightarrow x_2$ is 0.10s^{-1} , this means that, per unit time, the increase in activity in x_2 corresponds to 10% of the current activity in x_1

Neuronal state equations

Interim summary: bilinear neuronal state equation



$$\frac{dx}{dt} = \underbrace{\left(A + \sum_{j=1}^m u_j B^{(j)} \right)}_{\text{connectivity}} x + \underbrace{Cu}_{\text{External inputs}}$$

State change

External inputs

Current state

connectivity


$$\theta = \{A, B^{(1)}, \dots, B^{(m)}, C\}$$

Endogenous connectivity

Modulatory connectivity

Driving inputs



DCM for fMRI - Overview



The hemodynamic response

Neuronal dynamics only indirectly observable via hemodynamic response

- ↑ neuronal activity
- ↑ blood flow
- ↑ oxygenated Hb
- ↑ T2*
- ↑ fMRI signal



The hemodynamic model


6 parameters:

$$\theta^h = \{\kappa, \gamma, \tau, \alpha, \rho, \varepsilon\}$$


Important for model fitting,
but typically of no interest
for statistical inference.

Region specific HRF

→ Parameters computed
separately for each region



DCM for fMRI - Overview



The BOLD signal equation

Resting blood
volume

Deoxyhemoglobin
content

Blood
volume



$$\lambda(q, \nu) = \frac{\Delta S}{S_0} \approx V_0 \left[k_1(1 - q) + k_2 \left(1 - \frac{q}{\nu} \right) + k_3(1 - \nu) \right]$$

BOLD-Signal Parameters:

$$k_1 = 4.3\vartheta_0 E_0 TE$$

$$k_2 = \varepsilon r_0 E_0 TE$$

$$k_3 = 1 - \varepsilon$$

$$V_0 = 0.04 \quad E_0 = 0.32 - 0.4$$

At 1.5 Tesla

$$\vartheta_0 \approx 40.3 \text{ s}^{-1}$$

$$r_0 \approx 25 \text{ s}^{-1}$$

$$TE \approx 0.04 \text{ s}$$

$$\varepsilon \approx 1.28$$

At 3 Tesla

$$\vartheta_0 \approx 80.6 \text{ s}^{-1}$$

$$r_0 \approx 110 \text{ s}^{-1}$$

$$TE \approx 0.035 \text{ s}$$

$$\varepsilon \approx 0.47$$

At 7 Tesla


$$\vartheta_0 \approx 188 \text{ s}^{-1}$$

$$r_0 \approx 340 \text{ s}^{-1}$$

$$TE \approx 0.025 \text{ s}$$

$$\varepsilon \approx 0.026$$

From neural activity to the BOLD signal: Summary




BOLD signal is a **direct function** of v and q

$$y = \frac{\Delta S}{S_0} = g(v, q) + \varepsilon$$




Simulation example: What can DCM explain?

Example: single node




$$\frac{dx}{dt} = Ax + u_2 B^{(2)}x + Cu_1$$

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} a_{11} & 0 \\ 0 & a_{22} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + u_2 \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} c_{11} & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$






Simulation example: What can DCM explain?

Example: two connected node




$$\frac{dx}{dt} = Ax + u_2 B^{(2)}x + Cu_1$$

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} a_{11} & 0 \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + u_2 \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} c_{11} & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$






Simulation example: What can DCM explain?

Example: modulation of connection




$$\frac{dx}{dt} = Ax + u_2 B^{(2)}x + Cu_1$$

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} a_{11} & 0 \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + u_2 \begin{bmatrix} 0 & 0 \\ b_{21}^{(2)} & 0 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} c_{11} & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$






Simulation example: What can DCM explain?

Example: modulation of inhibitory self-connection



$$\frac{dx}{dt} = Ax + u_2 B^{(2)}x + Cu_1$$

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} a_{11} & 0 \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + u_2 \begin{bmatrix} 0 & 0 \\ 0 & b_{22}^{(2)} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} c_{11} & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$$




DCM for fMRI

A simple model of
a neural network

... described as a
dynamical system

... causes the data
(BOLD signal).

...



Neural node



Input




Connections

$$\dot{x} = f(x, u, \theta)$$

$$y = g(x, \theta) + \varepsilon$$

Let the system run with input (u) and parameters (θ), and you will get a BOLD signal time course y that you can compare to the measured data.

DCM for fMRI - Overview




Bayes' theorem

$$\text{posterior } p(\theta|y, m) = \frac{\text{likelihood prior}}{\text{model evidence}}$$


$p(y|\theta, m)p(\theta|m)$

likelihood prior

model evidence



Reverend Thomas Bayes
(1702-1761)





The likelihood function for DCM

$$p(y(t)|\theta, m) = \mathcal{N}(y(t); g(\theta^n, \theta^h, u), \theta^\sigma)$$


likelihood

Assume data is normally distributed around the prediction from the dynamical model (Gaussian noise)

The likelihood function for DCM

$$p(y(t)|\theta, m) = \mathcal{N}(y(t); g(\theta^n, \theta^h, u), \theta^\sigma)$$


likelihood



The likelihood function for DCM



$$p(y(t)|\theta, m) = \mathcal{N}(y(t); g(\theta^n, \theta^h, u), \theta^\sigma)$$

likelihood



The likelihood function for DCM

$$p(y(t)|\theta, m) = \mathcal{N}(y(t); g(\theta^n, \theta^h, u), \theta^\sigma)$$





Priors

$$p(\theta|y, m) = \frac{p(y|\theta, m)p(\theta|m)}{p(y|m)}$$

prior

Neuronal parameters:

- self-connections: principled (to “ensure” that the system is stable)
- other parameters (between—region connections, modulation, inputs): shrinkage priors


Hemodynamic parameters:

- empirical

Noise prior:

- assume relatively noisy data
(in SPM12 → set DCM.options.hE = 0; DCM.options.hC = 1)

Model estimation: running the machinery




Inversion – variational Free Energy approximation to model evidence

model evidence


$$\ln p(y) = \underbrace{\text{KL}[q||p]}_{\substack{\text{divergence} \\ \geq 0 \\ (\text{unknown})}} + \underbrace{F(q, y)}_{\substack{\text{neg. free energy} \\ (\text{easy to evaluate} \\ \text{for a given } q)}}$$

When $F(q, y)$ is maximized, $q(\theta)$ is our best estimate of the true posterior.




Model estimation: running the machinery

Variational Bayes



Example: Model Selection

- Specific sensory stimuli lead to unusual, additional experiences
- Grapheme-color synesthesia: **color**
- Involuntary, automatic; stable over time, prevalence ~4%
- Potential cause: aberrant **cross-activation/coupling** between brain areas
 - grapheme encoding area (FGA)
 - color area V4
 - superior parietal lobule (SPL)




Hubbard, 2007

Bottom-up or Top-down “cross-activation”?


Bottom-up

(Ramachandran & Hubbard, 2001)




Projectors

ABC




Top-down


(Grossenbacher & Lovelace, 2001)



Associators



Example: DCM for physiologically plausible feature extraction





What questions can we answer using DCM?

Model comparison

What is the functional architecture of a network of brain regions?

→ Synesthesia

Are optimal models different between groups?

→ Synesthesia

Which connections are modulated by experimental manipulations?

Parameter inference

Are parameters different between individuals/groups?

Use parameters as physiologically informed summary statistics

→ Generative embedding

... and of course many more!



DCM software note

Basic functionality for DCM for fMRI is provided within

SPM

<https://www.fil.ion.ucl.ac.uk/spm/>



Limitations

- Local minima:
 - Variational approximation can get stuck in local minima of free energy
- Size of networks:
 - Standard inversion gets prohibitively slow for large networks (more than 10 nodes).
- Regularization through fixed priors:
 - Current regularization depends on priors only. Regularization based on group (all subjects) would be better → empirical Bayes.

Recent additions to DCM for fMRI

- Massively parallel dynamic causal modelling

➤ **mpdcm**

Eduardo Aponte



- Regression dynamic causal modelling

➤ **rDCM**


Stefan Frässle



- Hierarchical unsupervised generative embedding



➤ **HUGE**

Yu Yao




Available in TAPAS:
www.translationalneuromodeling.org/tapas

Massively parallel DCM - mpdcm



$$\begin{aligned} \dot{x} &= f(x, u_1, \theta_1) \\ \dot{x} &= f(x, u_2, \theta_2) \\ &\vdots \\ \dot{x} &= f(x, u_n, \theta_n) \end{aligned} \quad \left. \right\} \text{mpdcm_integrate(dcms)} \quad \left[\begin{array}{c} y_1 \\ y_2 \\ \vdots \\ y_n \end{array} \right]$$





- Fast inversion of DCMs
 - MCMC based inversion possible
- **Thermodynamic Integration** (alternative to Free Energy)



Summary – Generative model



Summary – Generative Model




Summary - Bayesian System Identification

Neural (and hemo-) dynamics

Observer function


Inference on model structure

Inference on parameters



$$p(y | \theta, m) = N(g(\theta), \Sigma(\theta))$$
$$p(\theta, m) = N(\mu_\theta, \Sigma_\theta)$$

$$p(y | m) = \int p(y | \theta, m) p(\theta) d\theta$$
$$p(\theta | y, m) = \frac{p(y | \theta, m) p(\theta, m)}{p(y | m)}$$





Thank you!

Many thanks to Stefan Frässle,
Klaas Enno Stephan, Hanneke den Ouden
and Jean Daunizeau for many of the slides!

List with suggested DCM literature in Appendix of this presentation!



DCM literature (1)

- Aponte EA, Raman S, Sengupta B, Penny WD, Stephan KE, Heinze J (2016). mpdcm: A Toolbox for Massively Parallel Dynamic Causal Modeling. *Journal of Neuroscience Methods* 257: 7-16.
- Brodersen KH, Schofield TM, Leff AP, Ong CS, Lomakina EI, Buhmann JM, Stephan KE (2011) Generative embedding for model-based classification of fMRI data. *PLoS Computational Biology* 7: e1002079.
- Brodersen KH, Deserno L, Schlagenhauf F, Lin Z, Penny WD, Buhmann JM, Stephan KE (2014) Dissecting psychiatric spectrum disorders by generative embedding. *NeuroImage: Clinical* 4: 98-111
- Daunizeau J, David, O, Stephan KE (2011) Dynamic Causal Modelling: A critical review of the biophysical and statistical foundations. *NeuroImage* 58: 312-322.
- Daunizeau J, Stephan KE, Friston KJ (2012) Stochastic Dynamic Causal Modelling of fMRI data: Should we care about neural noise? *NeuroImage* 62: 464-481.
- **Friston KJ, Harrison L, Penny W (2003) Dynamic causal modelling. *NeuroImage* 19:1273-1302.**
- Friston KJ, Mattout J, Trujill-Barreto, Ashburner J, Penny W (2007) Variational free energy and the Laplace approximation. *NeuroImage* 34: 220-234.
- Friston K, Stephan KE, Li B, Daunizeau J (2010) Generalised filtering. *Mathematical Problems in Engineering* 2010: 621670.
- Friston KJ, Li B, Daunizeau J, Stephan KE (2011) Network discovery with DCM. *NeuroImage* 56: 1202–1221.
- Friston K, Penny W (2011) Post hoc Bayesian model selection. *Neuroimage* 56: 2089-2099.
- Friston KJ, Kahan J, Biswal B, Razi A (2014) A DCM for resting state fMRI. *Neuroimage* 94:396-407.
- Frässle S, Yao Y, Schöbi S, Aponte EA, Heinze J, Stephan KE (in press) Generative models for clinical applications in computational psychiatry. *Wiley Interdisciplinary Reviews: Cognitive Science*.
- **Frässle S, Lomakina EI, Razi A, Friston KJ, Buhmann JM, Stephan KE (2017) Regression DCM for fMRI. *NeuroImage* 155:406-421.**



DCM literature (2)

- Kiebel SJ, Kloppel S, Weiskopf N, Friston KJ (2007) Dynamic causal modeling: a generative model of slice timing in fMRI. *NeuroImage* 34:1487-1496.
- Li B, Daunizeau J, Stephan KE, Penny WD, Friston KJ (2011). Stochastic DCM and generalised filtering. *NeuroImage* 58: 442-457
- Marreiros AC, Kiebel SJ, Friston KJ (2008) Dynamic causal modelling for fMRI: a two-state model. *NeuroImage* 39:269-278.
- Penny WD, Stephan KE, Mechelli A, Friston KJ (2004a) Comparing dynamic causal models. *NeuroImage* 22:1157-1172.
- Penny WD, Stephan KE, Mechelli A, Friston KJ (2004b) Modelling functional integration: a comparison of structural equation and dynamic causal models. *NeuroImage* 23 Suppl 1:S264-274.
- Penny WD, Stephan KE, Daunizeau J, Joao M, Friston K, Schofield T, Leff AP (2010) Comparing Families of Dynamic Causal Models. *PLoS Computational Biology* 6: e1000709.
- Penny WD (2012) Comparing dynamic causal models using AIC, BIC and free energy. *NeuroImage* 59: 319-330
- **Raman S, Deserno L, Schlagenhauf F, Stephan KE (2016). A hierarchical model for integrating unsupervised generative embedding and empirical Bayes. *Journal of Neuroscience Methods* 269: 6-20.**
- Rigoux L, Stephan KE, Friston KJ, Daunizeau J (2014). Bayesian model selection for group studies – revisited. *NeuroImage* 84: 971-985.
- Stephan KE, Harrison LM, Penny WD, Friston KJ (2004) Biophysical models of fMRI responses. *Curr Opin Neurobiol* 14:629-635.
- Stephan KE, Weiskopf N, Drysdale PM, Robinson PA, Friston KJ (2007) Comparing hemodynamic models with DCM. *NeuroImage* 38:387-401.
- Stephan KE, Harrison LM, Kiebel SJ, David O, Penny WD, Friston KJ (2007) Dynamic causal models of neural system dynamics: current state and future extensions. *J Biosci* 32:129-144.



DCM literature (3)

- Stephan KE, Weiskopf N, Drysdale PM, Robinson PA, Friston KJ (2007) Comparing hemodynamic models with DCM. *NeuroImage* 38:387-401.
- Stephan KE, Kasper L, Harrison LM, Daunizeau J, den Ouden HE, Breakspear M, Friston KJ (2008) Nonlinear dynamic causal models for fMRI. *NeuroImage* 42:649-662.
- Stephan KE, Penny WD, Daunizeau J, Moran RJ, Friston KJ (2009a) Bayesian model selection for group studies. *NeuroImage* 46:1004-1017.
- Stephan KE, Tittgemeyer M, Knösche TR, Moran RJ, Friston KJ (2009b) Tractography-based priors for dynamic causal models. *NeuroImage* 47: 1628-1638.
- **Stephan KE, Penny WD, Moran RJ, den Ouden HEM, Daunizeau J, Friston KJ (2010) Ten simple rules for Dynamic Causal Modelling. *NeuroImage* 49: 3099-3109.**
- Stephan KE, Mathys C (2014). Computational approaches to psychiatry. *Current Opinion in Neurobiology* 25: 85-92.
- **Yao Y, Raman SS, Schiek M, Leff A, Frässle S, Stephan KE (2018). Variational Bayesian Inversion for Hierarchical Unsupervised Generative Embedding (HUGE). *NeuroImage*, 179: 604-619**