





Anxiety Disorders

Computational Psychiatry Course 2021

13th September

Helen Schmidt

Outline

- Introduction
- Case studies: Generalized anxiety, Social anxiety, Agoraphobia, Panic disorders,
 Specific phobias combined with symptoms
- Comorbidities
- Differential diagnosis
- Diagnostics
- Pathomechanism
- Treatment
- Prognosis

Clinical perspective

Introduction

Anxiety -> Fear

Fear

- alarm response to a present threat
- > trigger to fight or flight (survival)
- symptoms: sweating, trembling, heart palpitation, nausea

Anxiety

- future-oriented mood state
- preparation for possible negative events
- → avoidance
- Fear symptoms + muscle tension

Anxiety disorders

- Cluster of various types of anxiety disorders
- Significant and uncontrollable feelings of anxiety
- Huge impact on social, occupational & personal life

Epidemiology

- ► Lifetime prevalence: 15-20% (2:1 = f:m)
- ► Prevalence post-COVID infection: 47%
- ► High burden of disease (quality of life, chronicity, disability)
- ► Starts in childhood (mean age of onset: 11 years)
- ► High heritability (30-70%)

Overview

disorders Phobias Others (ICD-10 F41) (ICD-10 F40) Agoraphobia Generalized Anxiety Social Panic Phobia Disorder Specific Phobia

Anxiety

ICD-10 classification

Generalized anxiety disorder (GAD)


Generalized anxiety disorder (GAD)

- Uncontrollable thoughts
- Persistent worries
- ► Different areas: finance, future, health, etc.
- Loss of control & low self-efficacy
- Low self-confidence

- Physical tension
- Insomnia
- → Prevalence: 6%



Overview



Social phobia

Social phobia


- Restricted to social situations
- Examples: Talking to an audience,
 - Eating in groups, Waiting in line
- Afraid of being judged/ embarrassed
- Low self-confidence
- Physical symptoms (blushing, sweating, jitter,...)

- → Isolated or generalized
- → Prevalence: 13%



Overview

Anxiety disorders



Agoraphobia


Agoraphobia

- ► Fear of places or situations where escape is difficult or no help is available
- > > 2 situations: open spaces (e.g. parking lot, street), crowd or
- enclosed places (shops, public transportation)
- Trigger physical symptoms/ panic attacks
- ► Avoidance of gatherings, public places, traveling alone → social isolation

→ Prevalence: 3%



Overview



Panic disorder

Panic disorder

→ Panic attack:

- Intensive, recurrent fear (10-30 minutes)
- Sudden, unexpected beginning, uncontrollable
- Cognition: Misperception of bodily signals
 - Fear to get crazy, faint or to die
- Physical symptoms: heart palpitation, sweating, chest pain, shortness of breath
- Trigger: physical exertion, emotional stress

→ Panic disorder:

- Several panic attacks over 1 months
- Avoidance



Overview Anxiety disorders Phobias Others Agoraphobia Generalized Anxiety Social Panic Phobia Disorder Specific Phobia

https://www.goodtherapy.org/learn-about-therapy/issues/agoraphobia https://www.therecoveryvillage.com/mental-health/generalized-anxiety-disorder/

Specific phobias

Restricted to specific objects/situations:


- Heights (Akrophobia)
- Animals (Zoophobia), eg. Arachnophobia
- Flight (Aviophobia)
- Claustrophobia (confined spaces, e.g. MRI, elevator)
- Dentist, blood, needle
- → Considerable suffering: Avoidance
- → Prevalence: 5%





Combination of anxiety disorders

Severity of symptoms







anxiety disorders

Symptoms

Symptoms


Levels of anxiety

- 1) Vegetative reaction
- 2) Emotional reaction
- 3) Cognition
- 4) Motoric, behavior

Comorbidities

Comorbidities

- = additional disorders
- High comorbidity rate within various anxiety disorders
- Depression
- Alcohol or substance addiction (self-medication)
- Personality disorders:



Co-treatment of comorbidity

Impact on progression

Differential diagnosis

Differential diagnosis

= symptom overlap

Psychiatric disorders

- Affective disorders
- Schizophrenia
- OCD (Obsessivecompulsive disorder)
- PTSD (Post-traumatic stress disorder)

Substance-induced

- Amphetamines/Cocaine
- Steroids

Withdrawal from


- Alcohol
- Benzodiazepine
- Opioids


Somatic diseases

- Cardiac diseases (Arrhytmia, heart attack)
- Pulmonary diseases (Asthma, embolism)
- Allergic reaction
- Thyroid dysfunction
- Dizziness, Low blood-sugar, etc.

Diagnostics

Diagnostics





Exclusion of somatic diseases

- Laboratory tests
- Electrocardiogram (ECG)
- Drug screenings

Pathomechanism

Pathomechanism

Neurobiology:

genetics, neurotransmitters

Environmental factors: stress, e.g. job loss, separation


Cognition

Childhood experiences, personality traits



Learning theory: Mowrer's 2-process model

1. Process: Fear → acquired by classical conditioning



Learning theory: Mowrer's 2-process model

2. Process: Avoidance → maintained by operant conditioning



Strong fear of dogs -> You avoid all situations with dogs (parks, photos,..)




Reward (no dogs in your life)




Avoidance strengthens your fear >
Negative Reinforcement

Dysfunctional cognition



Epigenetics




Risk

Intermediate phenotypes of anxiety
Anxiety disorders


Resilience

Adaptive functioning

Neural circuit of anxiety




Amygdala





Treatment

Treatment




Antidepressant





Selective Serotonin-/Noradrenaline Reuptake Inhibitor (SSRI/SNRI)



Standardmedication for anxiety

- Except for specific phobias

Caution: AD can trigger somatic symptoms

→ Start with low dose

Psychotherapy

Cognitive Behavioral Therapy (CBT)

1) Psychoeducation

2) Cognitive restructuring

3) Interoceptive Exposure

4) In-vivo Exposure


5) Coping-strategies



Social Phobia & Panic Disorder




1.) Psychoeducation (Information): Vicious circle of anxiety




2.) Cognitive restructuring

«uncontrollable» thoughts -> associated with physical symptoms



3.) Interoceptive exposure



Force somatic symptoms ->

Panic sensation

→ Repetition: until symptoms are no longer perceived as «dangerous»

4.) In-vivo Exposure

→ Confrontation with situation

2 Options:

«Stimulus flooding»

- max. trigger of anxiety symptoms
- task: stay in situation → wait until «habituation»
 - anxiety reaction decreases

«Systematic desensitization» / «Graduated exposure»

gradually approach: reduction of anxiety symptoms






5) Coping strategies & emotion regulation

→ Relaxation exercises (PMR), breathing techniques, mindfulness, meditation



Virtual Reality (VR)



Exposure therapy

- Indication: various anxiety disorders
- Advantage: time saving, availability
- Side-effect: cybersickness



https://www.youtube.com/watch?v=qhsJD-2loZE


Prognosis

Prognosis & current situation

Anxiety

- High number of unreported cases
- Long latency period until final diagnosis
- ► 50% of anxiety patients are untreated
- Chronic progession, associated with comorbidities

Open problems that CP should address



Diagnostics

- Early detection (e.g. high-risk individuals, people with mild symptoms/ personality traits) → prevention strategies
- Identify subgroups
- Prediction of anxiety progression

Treatment

- Prediction of treatment selection
- Prediction of treatment outcome

CP References

Bishop, S. J., & Gagne, C. (2018). Anxiety, Depression, and Decision Making: A Computational Perspective. *Annu Rev Neurosci, 41*, 371-388. doi:10.1146/annurev-neuro-080317-062007

Browning, M., Behrens, T. E., Jocham, G., O'Reilly, J. X., & Bishop, S. J. (2015). Anxious individuals have difficulty learning the causal statistics of aversive environments. *Nature Neuroscience*, 18(4), 590-596. doi:10.1038/nn.3961

Paulus, M. P., & Yu, A. J. (2012). Emotion and decision-making: affect-driven belief systems in anxiety and depression. *Trends Cogn Sci, 16*(9), 476-483. doi:10.1016/j.tics.2012.07.009

Lecture References

Belmaker, R. H., & Agam, G. (2008). Major depressive disorder. N Engl J Med, 358(1), 55-68. doi:10.1056/NEJMra073096

Deng, J., Zhou, F., Hou, W., Silver, Z., Wong, C. Y., Chang, O., .Zuo, Q. K. (2021). The prevalence of depression, anxiety, and sleep disturbances in COVID-19 patients: a meta-analysis. *Ann N Y Acad Sci, 1486*(1), 90-111. doi:10.1111/nyas.14506

Domschke, K., Kapfhammer, H.-P., & Deckert, J. (2016). Angststörungen. In H.-J. Möller, G. Laux, & H.-P. Kapfhammer (Eds.), *Psychiatrie, Psychosomatik, Psychotherapie: Band 1: Allgemeine Psychiatrie Band 2: Spezielle Psychiatrie* (pp. 1-77). Berlin, Heidelberg: Springer Berlin Heidelberg.

Schiele, M. A., & Domschke, K. (2018). Epigenetics at the crossroads between genes, environment and resilience in anxiety disorders. *Genes Brain Behav, 17*(3), e12423. doi:10.1111/gbb.12423

World Health, O. (2004). ICD-10: international statistical classification of diseases and related health problems: tenth revision. In (2nd ed ed.). Geneva: World Health Organization.

Yang, X., Fang, Y., Chen, H., Zhang, T., Yin, X., Man, J., Lu, M. (2021). Global, regional and national burden of anxiety disorders from 1990 to 2019: results from the Global Burden of Disease Study 2019. *Epidemiology and psychiatric sciences, 30*, e36-e36. doi:10.1017/S2045796021000275

Zhang, W. H., Zhang, J. Y., Holmes, A., & Pan, B. X. (2021). Amygdala Circuit Substrates for Stress Adaptation and Adversity. *Biol Psychiatry*, 89(9), 847-856. doi:10.1016/j.biopsych.2020.12.026







Dr. med. Helen Schmidt, MPH
Consultant Psychiatrist and Psychotherapist

Medical Head

Translational Neuromodeling Unit (TNU)
schmidt@biomed.ee.ethz.ch

University & ETH Zurich

Dept. of Information Technology and Electrical Engineering
Institute of Biomedical Engineering and Neuroinformatics
Wilfriedstrasse 6
8032 Zürich

www.tnu.ethz.ch