

Schizophrenia (for non-clinicians)

04.09.2023, CPC Zurich

Dr. med. Jakob Siemerku
Translational Neuromodeling Unit
University of Zurich & ETH Zurich

siemerku@biomed.ee.ethz.ch



Why is it important to know about schizophrenia?

- One of the “major” psychiatric disorders
- 0.5% prevalence (Saha, 2005)
- Severe mental disorder
- Functional impairment
- Social dimension
- Many unsolved questions



Terminology

- **Schizophrenia:** Diagnosis
- **Psychotic Disorder/Psychosis:** Collective term for conditions with psychotic symptoms, used when course and causal factor(s) are unknown
- **Psychotic:** Symptoms related to difficulties in reality testing (delusions, hallucination, disorganized behaviour, ...); can occur in many psychiatric conditions
- Terms often used incorrectly and interchangeably

Important side-note

- Always use „person/patient with schizophrenia“, never use „schizophrenic“
- “Schizophrenia” perceived as stigmatizing, most prefer to use the term „psychosis“



Clinical Manifestation





- 37yo librarian
- Messy apartment
- Has been talking “weirdly” for weeks, but now unable to talk
- Apathy
- Slowed down
- Lost her job months ago

- 19yo student of physics
- Neighbour:
 - Spying on him
 - Plotting to murder him
 - CIA-affiliate
 - Hears N.'s voice
 - N. can read his mind
- Started six weeks ago
- Did not attend classes for ½ year



Symptom Categories



A 2x3 grid of interlocking puzzle pieces. The top row contains three pieces: pink (labeled 'positive'), light red (labeled 'mood'), and grey (labeled 'motor'). The bottom row contains three pieces: blue (labeled 'negative'), orange (labeled 'disorganization'), and light green (labeled 'cognition'). The pieces are set against a light blue background with faint outlines of other puzzle pieces.

positive

mood

motor

negative

disorga-
nization

cognition

Symptom Categories

positive

- “failure in reality testing”
- delusion
- hallucination
- ego-boundary distortion

negative

disorga-
nization

cognition



Symptom Categories



positive

mood

motor

negative

- loss of affective and driving functions
- impaired affective experience/expression
- loss of motivation & initiative
- reduced social drive
- slowed thinking

Symptom Categories



mood

motor



- overlap with negative symptoms
- anxiety, emotional arousal, and depression very frequent

Symptom Categories

positive

mood

- loosening of association
- disorganized speech
- disorganized behaviour

negative

disorga-
nization

cognition

Symptom Categories

"I'm blued in like blingo."

positive

mood

- loosening of association
- disorganized speech
- disorganized behaviour

negative

disorga-
nization

cognition

Symptom Categories

- excessive purposeless activities
- stereotypies
- Catatonic symptoms:
 - Inactivity/mutism ↔ agitation

motor

negative

disorga-
nization

cognition

Symptom Categories

positive

mood

motor

- general cognitive deficits
- pronounced in domains like working memory

negative

disorga
nization

cognition

Diagnostic Criteria (DSM 5)

MAIN CRITERIA	<ul style="list-style-type: none">• ≥ 2 symptom (categories) present• AND ≥ 1 core symptom• NO OTHER CAUSE!
TIME	<ul style="list-style-type: none">• ≥ 1 month main criteria• ≥ 6 months symptoms/functional impairment
SYMPTOMS	<ul style="list-style-type: none">• (core) delusions• (core) hallucinations• (core) disorganized speech • negative symptoms• disorganized or catatonic behaviour

(APA, 2013)



- 37yo librarian
 - Messy apartment
 - Has been talking “weirdly” for weeks, but now unable to talk
 - Apathy
 - Slowed down
 - Lost her job months ago
-
- Exam & lab results ok
 - Starts talking, uses “blue” in many contexts like “I’m blued in like blingo.”
- Schizophrenia

- 19yo student of physics
 - Neighbour:
 - Spying on him
 - Plotting to murder him
 - CIA-affiliate
 - Hears N.'s voice
 - N. can read his mind
 - Started six weeks ago
 - Did not attend classes for ½ year


 - Exam, lab & MRI results ok
- Schizophrenia




Course, Risk Factors, Outcome




Course of Schizophrenia




Course: Episodes & Outcome



Risk Factors across Lifespan



Clinical Care





- Aripiprazole → no improvement
- Risperidone → improvement, but motor side-effects
- Clozapine → function improves much
- Discharge to live with parents after 3 months

- 37yo librarian
 - Messy apartment
 - Has been talking “weirdly” for weeks, but now unable to talk
 - Apathy
 - Slowed down
 - Lost her job months ago
-
- Exam & lab results ok
 - Starts talking, uses “blue” in many contexts like “I’m blued in like blingo.”
- Schizophrenia

- 19yo student of physics
- Neighbour:
 - Spying on him
 - Plotting to murder him
 - CIA-affiliate
 - Hears N.'s voice
 - N. can read his mind
- Started six weeks ago
- Did not attend classes for ½ year

• Exam, lab & MRI results ok



→ Schizophrenia



- Risperidone → remission after 20d
- Day care clinic for 6wks
- Returns to uni after 9wks



Antipsychotics

Mesolimbic Pathway
Untreated Schizophrenia



“hyperactive” dopaminergic system

Mesolimbic Pathway
D2 Antagonist




“normalized” dopaminergic system

Antipsychotics

Overall change in symptoms

SMD (95% CrI)



(adapted from Leucht et al., 2013; Huhn et al., 2019)

Antipsychotics

- 50% respond to 1st line treatment
Response \neq remission, recovery or cure
- No prediction which antipsychotic is effective \rightarrow “*trial & error*”
in CH: 24 licensed, ~14 relevant for the treatment of Schizophrenia
- Major problem: discontinuation of treatment
 - Side effects
 - Poor insight

Treatment Strategies

- **Building trust, therapeutic relationship & working alliance**
- Management and prevention of side-effects
- Early treatment with antipsychotics



Treatment Strategies

- Activation & social support
- Psychoeducation
- Low-threshold service
- Cognitive Behavioural Therapy

(SGPP, 2016; DGPPN, 2012; NICE, 2014; APA, 2006)

Treatment Strategies

(Schizophrenia Canada, 2018)



**RECOVERY
IS POSSIBLE.
THE SOONER
THE TREATMENT,
THE BETTER.**

National Schizophrenia & Psychosis Awareness Day
May 24th, 2018
www.earlypsychosisintervention.ca




Pathophysiology





- **Phenotypic expression**
 - Continuum with affective disorders (Crow, 1986)
 - Failure of filter mechanisms (Hemsley and Zawada, 1976)
 - Internal monitoring deficiency (positive symptoms) & action initiation failure (negative symptoms) (Frith & Done, 1988)
 - Deficit vs. non-deficit SZ (Carpenter et al., 1988)
 - Aberrant salience syndrome (Kapur, 2003)
 - Dysconnection hypothesis (Stephan, 2009; Konrad & Winterer, 2008; dysmyelination: Segal et al., 2007)
 - Lateralization deficiency/language processing and distinction of thoughts and speech output (Crow, 2000)
- **Pathophysiological**
 - Abnormal transcallosal inter-hemispheric interaction → delusions of alien control (Nasrallah, 1985)
 - Dysfunction of inhibitory circuits (reduced power in the gamma range bands) (Kwon, 1999)
 - Corollary discharge (Feinberg, 1978; Frith & Done, 1988)
 - Hyperdopaminergic models (Carlsson, 1977; Randrup & Munkvad, 1967; Snyder, 1976); prefrontal-limbic DA imbalance (Weinberger, 1987), phasic-tonic FA imbalance model (Grace, 1991); common pathway hypothesis (Seeman, 2010)
 - NMDAR-hypofunction (Olney & Farber, 1995)
 - Altered GABAergic transmission (altered neural synchrony/cognitive deficits; reductions in GABAergic neurons) (Benes & Berreta, 2001)
 - Cholinergic hypotheses (Tandon and Greden, 1989)
 - Inflammation – kynurenic acid as endogenous NMDAR antagonist (tryptophan metabolism)
- **Pathogenesis**
 - Early developmental models – disruptions intruterine/early postnatal (neuronal proliferation, migration, differentiation, elimination, neurogenesis) → impaired neuronal structure, abnormal brain maturation (Murray, 2002)
 - Late developmental models – deviations in later emerging processes such as synaptic/axonal pruning/neuronal apoptosis and/or myelination)
 - Neurodegeneration → atrophic processes
 - Acceleration of aging → cortico-limbic glutamatergic activity because of reduced inhibition by GABAergic interneurons → excitotoxicity
 - Disturbed excitatory/inhibitory balance
- **Etiological**
 - Polygenic/multifactorial (Gottesman & Shields, 1967) → heritability, heterogeneity; copy number variations
 - Infectious diseases
 - Gene-environment interaction → two-hit-hypothesis (first genetic risk and early developmental alterations; then environmental factor) // epigenetic factors
 - „By-product“ of evolution of language (Crow, 2000)


Dopamine Hypothesis



Antipsychotics


Overall change in symptoms

SMD (95% CrI)



(adapted from Leucht et al., 2013; Huhn et al., 2019)

Genetic Evidence



(adapted from Greenwood et al., 2012)

Genome-wide association study of >35,000 cases:

Dopamine

- DRD2

Glutamate/NMDAr


- GRM3
- GRIN2A
- GRIA1
- SRR

Acetylcholine

- CHRNA3
- CHRNA5
- CHRNA4

(SWGPGC, 2014)


NMDAR × neuromodulator interactions



- phosphorylation of NMDARs
- modulation of membrane potential
- influence on receptor trafficking

Dysconnection Hypothesis

- NMDAR × neuromodulator interaction dysfunction:
 - Disturbed modulation (*precision*) of *prediction* and *Prediction Error* signals
 - Non-optimal inference



(Friston et al., 2016;
Stephan et al., 2016)


Dysconnection Hypothesis

„Overweight“ of Prediction Errors: Trait phenomena?

- ↓ Susceptibility to illusions
- Aberrant Salience, Delusional Mood
- ...

„Overweight“ of predictions: State phenomena?

- Delusions
- Acoustic hallucinations
- ...



(Adams et al., 2013; Frith & Friston, 2013; Holzman, 2000; Koethe et al, 2006; Umbricht & Krljes, 2005)

(Friston et al., 2016;
Stephan et al., 2016)


Dysconnection Hypothesis

„Overweight“ of Prediction Errors:
Trait phenomena?

- ↓ Susceptibility to illusions
- Aberrant Salience, Delusional Mood
- ...

„Overweight“ of predictions:
State phenomena?

- Delusions
- Acoustic hallucinations
- ...




(Friston et al., 2016;
Stephan et al., 2016)

(Adams et al., 2013; Frith & Friston, 2013; Powers et al., 2017, Sterzer et al, 2018)

To summarize...

- Severe mental disorder
- Functional impairment
- Heterogeneous
 - Risk factors – pathophysiology – symptoms – course - treatment response
- Antipsychotics are effective, but
 - focus on recovery!
- Computational Psychiatry → new approaches to treatment?

“Lost Years” by Bas Labruyère



Labruyère, B. (2011, May 11). Lost Years.
<https://vimeo.com/23611157>

References

- Adams, R. A., Stephan, K. E., Brown, H. R., Frith, C. D., & Friston, K. J. (2013). The computational anatomy of psychosis. *Frontiers in Psychiatry*, 4, 47. <https://doi.org/10.3389/fpsyt.2013.00047>
- American Psychiatric Association. (2006). American Psychiatric Association Practice Guidelines for the treatment of psychiatric disorders: compendium 2006.
- American Psychiatric Association. (2013). Diagnostic and Statistical Manual of Mental Disorders, 5th Edition: DSM-5 (5 edition). Washington, D.C: American Psychiatric Publishing.
- Benkert, O., Hippus, H., & Angelescu, I.-G. (2014). Kompendium der Psychiatrischen Pharmakotherapie (10., vollst. überarb. und aktualisierte Aufl). Springer.
- Bleuler, E. (1908). Die Prognose der Dementia praecox (Schizophreniegruppe). *Allgemeine Zeitschrift für Psychiatrie und Psychisch-Gerichtliche Medizin*, 65, 436–464.
- Corlett, P. R., Frith, C. D. & Fletcher, P. C. From drugs to deprivation: a Bayesian framework for understanding models of psychosis. *Psychopharmacology (Berl)* 206, 515–530 (2009).
- Crow, T. J. (2000). Schizophrenia as the price that Homo sapiens pays for language: A resolution of the central paradox in the origin of the species. *Brain Research Reviews*, 31(2), 118–129. [https://doi.org/10.1016/S0165-0173\(99\)00029-6](https://doi.org/10.1016/S0165-0173(99)00029-6)
- DGPPN, & Falkai, P. (Eds.). (2012). S3-Leitlinie Psychosoziale Therapien bei schweren psychischen Erkrankungen: S3-Praxisleitlinien in Psychiatrie und Psychotherapie. Springer-Verlag.
- Drake, R. J., Dunn, G., Tarrier, N., Bentall, R. P., Haddock, G., & Lewis, S. W. (2007). Insight as a predictor of the outcome of first-episode nonaffective psychosis in a prospective cohort study in England. *The Journal of Clinical Psychiatry*, 68(1), 81–86.
- Feldman, R. S., Jerrold S. Meyer, & Linda F. Quenzer. (1997). Principles of neuropsychopharmacology. Sunderland, Mass: Sinauer.
- Frith, C. D., & Friston, K. J. (2013). False Perceptions & False Beliefs: Understanding Schizophrenia. In A. Battro, S. Dehaene, & W. Singer (Eds.), *Neurosciences and the Human Person: New Perspectives on Human Activities*. Retrieved from <http://www.casinapiov.va/content/dam/accademia/pdf/sv121/sv121-frithc.pdf>
- Friston, K., Brown, H. R., Siemer, J., & Stephan, K. E. (2016). The dysconnection hypothesis (2016). *Schizophrenia Research*, 176(2–3), 83–94. <https://doi.org/10.1016/j.schres.2016.07.014>
- Fusar-Poli, P., Borgwardt, S., Bechdolf, A., Addington, J., Riecher-Rössler, A., Schultze-Lutter, F., ... Yung, A. (2013). The Psychosis High-Risk State: A Comprehensive State-of-the-Art Review. *JAMA Psychiatry*, 70(1), 107–120. <https://doi.org/10.1001/jamapsychiatry.2013.269>
- Greenwood, T. A., Light, G. A., Swerdlow, N. R., Radant, A. D., & Braff, D. L. (2012). Association Analysis of 94 Candidate Genes and Schizophrenia-Related Endophenotypes. *PLOS ONE*, 7(1), e29630. <https://doi.org/10.1371/journal.pone.0029630>
- Goldman, D., & Ducci, F. (2007). Deconstruction of Vulnerability to Complex Diseases: Enhanced Effect Sizes and Power of Intermediate Phenotypes [Research article]. <https://doi.org/10.1100/tsw.2007.210>
- Holzman, P. S. (2000). Eye movements and the search for the essence of schizophrenia. *Brain Research Reviews*, 31(2), 350–356. [https://doi.org/10.1016/S0165-0173\(99\)00051-X](https://doi.org/10.1016/S0165-0173(99)00051-X)
- Huhn, M. et al. Comparative efficacy and tolerability of 32 oral antipsychotics for the acute treatment of adults with multi-episode schizophrenia: a systematic review and network meta-analysis. *The Lancet* 394, 939–951 (2019).
- Jaspers, K. (1973). *Allgemeine Psychopathologie* (9., unveränderte Auflage). Berlin, Heidelberg, New York: Springer.
- Jones, C., Hacker, D., Meaden, A., Cormac, I., Irving, C. B., Xia, J., ... Chen, J. (2018). Cognitive behavioural therapy plus standard care versus standard care plus other psychosocial treatments for people with schizophrenia. *Cochrane Database of Systematic Reviews*, (11). <https://doi.org/10.1002/14651858.CD008712.pub3>
- Kaiser, S., Berger, G., Conus, P., Kawohl, W., Müller, T. J., Schimmelmann, B. G., ... Seifritz, E. (2016). SGPP Behandlungsempfehlungen Schizophrenie - Version Februar 2016.
- Koethe, D., Gerth, C. W., Neatby, M. A., Haensel, A., Thies, M., Schneider, U., ... Lewke, F. M. (2006). Disturbances of visual information processing in early states of psychosis and experimental delta-9-tetrahydrocannabinol altered states of consciousness. *Schizophrenia Research*, 88(1–3), 142–150.
- Leucht, S., Busch, R., Kissling, W., & Kane, J. M. (2007). Early prediction of antipsychotic nonresponse among patients with schizophrenia. *The Journal of Clinical Psychiatry*, 68(3), 352–360.
- Leucht, S., Cipriani, A., Spinelli, L., Mavridis, D., Örey, D., Richter, F., ... Davis, J. M. (2013). Comparative efficacy and tolerability of 15 antipsychotic drugs in schizophrenia: a multiple-treatments meta-analysis. *The Lancet*, 382(9896), 951–962. [https://doi.org/10.1016/S0140-6736\(13\)60733-3](https://doi.org/10.1016/S0140-6736(13)60733-3)
- Keshavan, M. S., Nasrallah, H. A., & Tandon, R. (2011). Schizophrenia, “Just the Facts” 6. Moving ahead with the schizophrenia concept: From the elephant to the mouse. *Schizophrenia Research*, 127(1), 3–13. <https://doi.org/10.1016/j.schres.2011.01.011>
- Keshavan, M. S., Tandon, R., Boutros, N. N., & Nasrallah, H. A. (2008). Schizophrenia, “just the facts”: What we know in 2008: Part 3: Neurobiology. *Schizophrenia Research*, 106(2), 89–107. <https://doi.org/10.1016/j.schres.2008.07.020>
- Menezes, N. M., Arenovich, T., & Zipursky, R. B. (2006). A systematic review of longitudinal outcome studies of first-episode psychosis. *Psychological Medicine*, 36(10), 1349–1362. <https://doi.org/10.1017/S0033291706007951>

References

- Millan, M. J., Andrieux, A., Bartzokis, G., Cadenhead, K., Dazzan, P., Fusar-Poli, P., ... Weinberger, D. (2016). Altering the course of schizophrenia: progress and perspectives. *Nature Reviews Drug Discovery*, 15(7), 485–515. <https://doi.org/10.1038/nrd.2016.28>
- National Collaborating Centre for Mental Health (UK). (2014). *Psychosis and Schizophrenia in Adults: Treatment and Management*. Retrieved from <http://www.ncbi.nlm.nih.gov/books/NBK248060/>
- Perkins, D. O., Gu, H., Boteva, K., & Lieberman, J. A. (2005). Relationship between duration of untreated psychosis and outcome in first-episode schizophrenia: a critical review and meta-analysis. *The American Journal of Psychiatry*, 162(10), 1785–1804. <https://doi.org/10.1176/appi.ajp.162.10.1785>
- Powers, A. R., Mathys, C., & Corlett, P. R. (2017). Pavlovian conditioning–induced hallucinations result from overweighting of perceptual priors. *Science*, 357(6351), 596–600. <https://doi.org/10.1126/science.aan3458>
- Saha, S., Chant, D., Welham, J., & McGrath, J. (2005). A Systematic Review of the Prevalence of Schizophrenia. *PLOS Medicine*, 2(5), e141. <https://doi.org/10.1371/journal.pmed.0020141>
- Saklad, S. R. (2018). Graphic representation of pharmacology: Development of an alternative model. *The Mental Health Clinician*, 7(5), 201–206. <https://doi.org/10.9740/mhc.2017.09.201>
- Schizophrenia Working Group of the Psychiatric Genomics Consortium. (SWGGPGC, 2014). Biological insights from 108 schizophrenia-associated genetic loci. *Nature*, 511(7510), 421–427. <https://doi.org/10.1038/nature13595>
- Schwartz, T. L., Sachdeva, S., & Stahl, S. M. (2012). Glutamate Neurocircuitry: Theoretical Underpinnings in Schizophrenia. *Frontiers in Pharmacology*, 3. <https://doi.org/10.3389/fphar.2012.00195>
- Sifakis, S., Tzachanis, D., Samara, M., & Papazisis, G. (2018). Antipsychotic Drugs: From Receptor-binding Profiles to Metabolic Side Effects. *Current Neuropharmacology*, 16(8), 1210–1223. <https://doi.org/10.2174/1570159X15666170630163616>
- Stephan, K. E., Diaconescu, A. O., & Iglesias, S. (2016). Bayesian inference, dysconnectivity and neuromodulation in schizophrenia. *Brain*, 139(7), 1874–1876. <https://doi.org/10.1093/brain/aww120>
- Stephan, K. E., Iglesias, S., Heinze, J., & Diaconescu, A. O. (2015). Translational Perspectives for Computational Neuroimaging. *Neuron*, 87(4), 716–732. <https://doi.org/10.1016/j.neuron.2015.07.008>
- Stephen M. Stahl. (2013). *Stahl's essential psychopharmacology: neuroscientific basis and practical application* (4th ed.). Cambridge: Cambridge University Press.
- Sterzer, P., Adams, R. A., Fletcher, P., Frith, C., Lawrie, S. M., Muckli, L., ... Corlett, P. R. (2018). The predictive coding account of psychosis. *Biological Psychiatry*. <https://doi.org/10.1016/j.biopsych.2018.05.015>
- Tandon, R., Keshavan, M. S., & Nasrallah, H. A. (2008a). Schizophrenia, “Just the Facts”: What we know in 2008: Part 1: Overview. *Schizophrenia Research*, 100(1), 4–19. <https://doi.org/10.1016/j.schres.2008.01.022>
- Tandon, R., Keshavan, M. S., & Nasrallah, H. A. (2008b). Schizophrenia, “Just the Facts” What we know in 2008. 2. Epidemiology and etiology. *Schizophrenia Research*, 102(1), 1–18. <https://doi.org/10.1016/j.schres.2008.04.011>
- Tandon, R., Nasrallah, H. A., & Keshavan, M. S. (2009). Schizophrenia, “just the facts” 4. Clinical features and conceptualization. *Schizophrenia Research*, 110(1), 1–23. <https://doi.org/10.1016/j.schres.2009.03.005>
- Tandon, R., Nasrallah, H. A., & Keshavan, M. S. (2010). Schizophrenia, “Just the Facts” 5. Treatment and prevention Past, present, and future. *Schizophrenia Research*, 122(1), 1–23. <https://doi.org/10.1016/j.schres.2010.05.025>
- Umbricht, D., & Krljes, S. (2005). Mismatch negativity in schizophrenia: a meta-analysis. *Schizophrenia Research*, 76(1), 1–23. <https://doi.org/10.1016/j.schres.2004.12.002>
- van Os, J. (2009). A salience dysregulation syndrome. *The British Journal of Psychiatry*, 194(2), 101–103. <https://doi.org/10.1192/bjp.bp.108.054254>

Media References

- Canada, S. (2018T06:29). Today is #NSPAD18! #Recovery is possible, and expected, with the right treatments. Effective treatment can include counselling, medication, meaningful activity, & support from loved ones --> [#https://www.earlypsychosisintervention.ca/psychosis](https://www.earlypsychosisintervention.ca/psychosis) #SupportSZ #CMHA100 #Schizophrenia #EndStigmatic.twitter.com/cTseJJT9Yd [Tweet]. Retrieved August 23, 2019, from @schizophreniac website: <https://twitter.com/schizophreniac/status/999643428605526016>
- JanBaby. (n.d.). Free Image on Pixabay - Signpost, Road Signs, Sign, Post. Retrieved August 23, 2019, from <https://pixabay.com/illustrations/signpost-road-signs-sign-post-2030780/>
- Labruyère, B. (2011, May 11). Lost Years. <https://vimeo.com/23611157>
- mcmurryjulie | Pixabay. (n.d.). Retrieved August 30, 2019, from <https://pixabay.com/users/mcmurryjulie-2375405/>
- Netherlands, S. van der W. from. (2010). In order to reach the highest mountains, you shouldn't be afraid of the deepest valleys. Retrieved from [https://commons.wikimedia.org/wiki/File:Depressed_\(4649749639\).jpg](https://commons.wikimedia.org/wiki/File:Depressed_(4649749639).jpg)
- Munch, E. (1893). English: The Scream [Oil, tempera & pastel on cardboard]. National Gallery of Norway. https://commons.wikimedia.org/wiki/File:Edvard_Munch,_1893,_The_Scream,_oil,_tempera_and_pastel_on_cardboard,_91_x_73_cm,_National_Gallery_of_Norway.jpg
- Open-Clipart-Vectors. (n.d.). Free Image on Pixabay - Biology, Brain, Cortex, Holiday. Retrieved September 1, 2019, from <https://pixabay.com/vectors/biology-brain-cortex-holiday-1295897/>
- OpenIcons. (n.d.). Free Image on Pixabay - World, Globe, Global, Earth. Retrieved September 1, 2019, from <https://pixabay.com/vectors/world-globe-global-earth-97864/>
- Pablo Picasso, 1901-02, Femme aux Bras Croisés, Woman with Folded Arms (Madchenbildnis), oil on canvas, 81 × 58 cm (32 × 23 in).jpg. (2020). In Wikipedia. [https://en.wikipedia.org/w/index.php?title=File:Pablo_Picasso,_1901-02,_Femme_aux_Bras_Crois%C3%A9s,_Woman_with_Folded_Arms_\(Madchenbildnis\),_oil_on_canvas,_81_%C3%97_58_cm_\(32_%C3%97_23_in\).jpg&oldid=974424175](https://en.wikipedia.org/w/index.php?title=File:Pablo_Picasso,_1901-02,_Femme_aux_Bras_Crois%C3%A9s,_Woman_with_Folded_Arms_(Madchenbildnis),_oil_on_canvas,_81_%C3%97_58_cm_(32_%C3%97_23_in).jpg&oldid=974424175)
- User:Fvasconcellos, U. P. J. L. (2015). English: The main dopaminergic pathways of the human brain: the mesocortical pathway, connecting the ventral tegmental area (VTA) with the frontal cortex; the mesolimbic pathway, connecting the VTA with the nucleus accumbens; the nigrostriatal pathway, connecting the substantia nigra with the dorsal striatum; and the tuberoinfundibular pathway, connecting the hypothalamus with the pituitary. Retrieved from https://commons.wikimedia.org/wiki/File:Dopaminergic_pathways.svg; edited
- James, D. (2019). English: Two doses of intravenous ketamine, 50mg/ml and 10mg/ml. Retrieved from https://commons.wikimedia.org/wiki/File:Two_doses_of_iv_ketamine.jpg, edited
- Unknown. (1962). English: Advert from ca. 1962 for Thorazine (trade-name of chlorpromazine in the U.S.). An antipsychotic (neuroleptic, major tranquilizer, antischizophrenic, actaractic). In Europe it is known as Largactil. Retrieved from https://commons.wikimedia.org/wiki/File:Thorazine_advert.jpg