

Assignment 3

Peter Lorenz

Infimal Convolution: Several lambdas= $\{0.1, 0, 10\}$ are used to show the infimal convolutions.




Figure 1: $f(y) = \max(0, 1-y)$




Figure 2: $f(y) = |y| + \frac{1}{3}|y|^3$




Figure 3: $f(y) = \begin{cases} 0 & \text{if } y \in [a, b] \\ +\infty & \text{else} \end{cases}$




Figure 4: $f(y) = \max(|y|, y^2)$

Proximal Operators: Beside the original function (blue), we plotted the proximal of the function (orange).




Figure 5: $f(x) = \begin{cases} -a \ln(x) + \frac{x^2}{2} & x > 0 \\ +\infty & x \leq 0 \end{cases}$




Figure 6: $f(x) = \max(|x| - a, 0)$




Figure 7: $f(x) = \begin{cases} \ln(a - |x|) & \text{for } |x| < a \\ +\infty & \text{otherwise} \end{cases}$

(F)ISTA: The followig plots show the comparison of the ISTA and to the fast one FISTA. In the end, the Energy of ISTA and FISTA will be equal.




Figure 8: Energy Plot. 16 Dim




Figure 9: Energy Plot. 32 Dim




Figure 10: Energy Plot. 64 Dim





Figure 11: Energy Plot. 128 Dim


Reconstructed Images: We want to reconstruct images where have the an original data base and a D with parts of B and C that we want to find. D multiplied with C gives us the reconstructed image of matrix B. The higher the dimension= $\{16,32,64,128\}$ of D the better the reconstruction.




Figure 12: Original Data Base B




(a) Reconstructed DC 16



(b) Reconstructed DC 32




(c) Reconstructed DC 64



(d) Reconstructed DC 128


Figure 13: ISTA reconstructed




(a) Reconstructed DC 16



(b) Reconstructed DC 32



(c) Reconstructed DC 64



(d) Reconstructed DC 128

Figure 14: FISTA reconstructed