

Pure Number

Time Limit : 1 Second

Memory Limit : 512 MB

Let's start with some basic definitions. A **prime number** is a positive integer greater than 1 whose divisor are 1 and itself, e.g., 2, 3, 5, 7. Any other positive integers greater than 1 which are not prime are called **composite number**, e.g., 4, 6, 8, 9, 10.

Now, let's define a pure number. A **pure number** of degree K is a positive integer which can be formed by multiplying exactly K unique prime numbers. As you might have noticed, a pure number of degree 1 is also a prime number. For example,

- 2 is a pure number of degree 1, i.e. 2;
- 6 is a pure number of degree 2, i.e. 2×3 ;
- 35 is a pure number of degree 2, i.e. 5×7 ;
- 66 is a pure number of degree 3, i.e. $2 \times 3 \times 11$;
- 12 is not a pure number as it cannot be formed by multiplying unique prime numbers.

By definition, 1 is not a pure number.

Given three integers L, R, and K, your task in this problem is to determine how many pure number of degree K between L and R (inclusive).

Input

Input begins with an integer: T ($1 \leq T \leq 100$) denoting the number of cases.

Each case contains the following input block: Each case contains three integers: L R K ($1 \leq L \leq R \leq 10^6$; $1 \leq K \leq 1000$) in a line.

Output

For each case, output in a line "Case #X: Y" where X is the case number (starts from 1) and Y is the output for the respective case.

Examples

input	Example #1
3 1 10 1 1 10 2 10 14 3	
output	

Case #1: 4

Case #2: 2

Case #3: 0

explanation

Case 1: 2, 3, 5, and 7 are pure numbers of degree 1

Case 2: 6 and 10 are pure numbers of degree 2.

Case 3: There is no pure number of degree 3 between 10 and 14.

End of Problem