




Annelida — cnetplot: top-10 terms × top-19 foreground genes (fold ≥ 3)










Bilateria — cnetplot: top-10 terms × top-11 foreground genes (fold ≥ 3)




Bivalvia — cnetplot: top-10 terms x top-11 foreground genes (fold ≥ 3)







Chordata — cnetplot: top-10 terms x top-26 foreground genes (fold ≥ 3)




Clitellata — cnetplot: top-10 terms x top-18 foreground genes (fold ≥ 3)




Cnidaria — cnetplot: top-10 terms × top-22 foreground genes (fold ≥ 3)




Coleoidea — cnetplot: top-10 terms × top-12 foreground genes (fold ≥ 3)



Ctenophora — cnetplot: top-10 terms x top-23 foreground genes (fold ≥ 3)



Decapodiformes — cnetplot: top-10 terms x top-14 foreground genes (fold ≥ 3)



Deuterostomia — cnetplot: top-10 terms × top-20 foreground genes (fold ≥ 3)

blood vessel development

BMPR2
RARGF2

RNA processing

RPS7
DROSHA
NOP58
PRKX1

translation initiation factor activ...

eukaryotic translation initiation f...

eukaryotic 48S preinitiation complex

EIF4E
MCT51
EIF2B4
COP55
EIF3E
EIF3I
EIF3J
EIF3H

mitochondrial DNA replication

RNASEH1
POLG


maintenance of DNA repeat elements

somatic recombination of immunoglob...


negative regulation of DNA recomb...

guanine/thymine mispair binding


MSH2




Diptera — cnetplot: top-10 terms × top-14 foreground genes (fold ≥ 3)






Echinodermata — cnetplot: top-10 terms x top-16 foreground genes (fold ≥ 3)






Gastropoda — cnetplot: top-10 terms x top-19 foreground genes (fold ≥ 3)




Hexapoda — cnetplot: top-10 terms x top-7 foreground genes (fold ≥ 3)




Insecta — cnetplot: top-10 terms × top-7 foreground genes (fold ≥ 3)




Mammalia — cnetplot: top-10 terms × top-14 foreground genes (fold ≥ 3)






Mollusca — cnetplot: top-10 terms x top-14 foreground genes (fold ≥ 3)




Nematoda — cnetplot: top-10 terms × top-27 foreground genes (fold ≥ 3)










Platyhelminthes — cnetplot: top-10 terms x top-19 foreground genes (fold ≥ 3)





Porifera — cnetplot: top-10 terms x top-4 foreground genes (fold ≥ 3)



Protostomia — cnetplot: top-10 terms x top-5 foreground genes (fold ≥ 3)



Scaphopoda — cnetplot: top-10 terms × top-20 foreground genes (fold ≥ 3)




namespace





BP

CC

Spiralia — cnetplot: top-10 terms × top-34 foreground genes (fold ≥ 3)



Teleostei — cnetplot: top-10 terms x top-12 foreground genes (fold ≥ 3)



Vertebrata — cnetplot: top-10 terms × top-31 foreground genes (fold ≥ 3)

