

Digital Health Technologies for Sustainable Healthcare System Transformation

Kantemoriva Mira¹, Khadartsev Aslan^{1(*)}, Dzidzoeva Yana¹, Bedoeva Dana¹, Alborov Khetag¹

¹North Ossetian State Medical Academy

Abstract. Digital health technologies—including telemedicine, artificial intelligence, Internet of Medical Things (IoT), wearable sensors, electronic health records, and mobile health applications—are catalyzing a fundamental transformation of healthcare systems toward greater sustainability. This transformation addresses the triple bottom line of sustainability: economic (reducing costs through preventive care, remote monitoring, and optimized resource allocation), environmental (minimizing facility-based care emissions, reducing medical waste, and enabling virtual consultations), and social (expanding equitable access, empowering patient self-management, and strengthening primary care). Evidence indicates that strategic implementation of these technologies can alleviate workforce shortages, improve chronic disease management, enhance diagnostic accuracy, and shift systems from reactive sick-care to proactive, person-centered health ecosystems. However, realizing sustainable transformation requires overcoming critical barriers: interoperability gaps, data privacy concerns, digital literacy disparities, regulatory fragmentation, and upfront investment challenges. Success hinges on integrated policy frameworks that prioritize equity, robust digital infrastructure, cross-sector collaboration, and human-centered design that complements—rather than replaces—clinician expertise. As healthcare systems globally confront aging populations, rising non-communicable diseases, and climate-related health threats, digital health technologies offer not merely incremental efficiency gains but a foundational pathway to resilient, adaptive, and universally accessible healthcare capable of delivering quality outcomes within planetary boundaries. Future efforts must focus on inclusive governance, evidence-based scaling, and measuring sustainability impact beyond clinical metrics to ensure digital transformation serves both people and the planet.

1 Introduction

Contemporary healthcare systems worldwide face a confluence of unprecedented pressures that threaten their long-term viability. Aging populations, the escalating burden of chronic non-communicable diseases, workforce shortages, and rising costs strain resources to their limits—while simultaneously, the healthcare sector itself contributes an estimated 4–5% of global carbon emissions, raising urgent questions about its environmental footprint (Karliner et al., 2019). Compounding these challenges, persistent inequities in access to quality care leave billions underserved, particularly in low-resource and rural settings. In this context, the pursuit of *sustainable* healthcare—defined not merely as cost containment but as a holistic commitment to economic viability, environmental stewardship, and social equity—has evolved from an aspirational goal to an operational imperative.

Sustainability in healthcare transcends traditional efficiency metrics. It embodies a triple bottom line framework: economic sustainability demands value-based care models that optimize resource allocation and shift focus from volume to outcomes; environmental sustainability requires decarbonization of clinical operations, waste reduction, and minimizing the ecological impact of care delivery; and social sustainability

* Corresponding author: Khadar1537@mail.ru

necessitates universal access, cultural responsiveness, and empowerment of both patients and providers within resilient health ecosystems (WHO, 2021). Achieving this integrated vision demands systemic transformation—not incremental adjustments to legacy structures designed for acute, facility-centric care.

Digital health technologies have emerged as pivotal enablers of this transformation. Encompassing telemedicine, artificial intelligence (AI) and machine learning, the Internet of Medical Things (IoMT), wearable biosensors, interoperable electronic health records (EHRs), and mobile health (mHealth) applications, these tools offer unprecedented opportunities to reimagine care delivery. They facilitate a shift from reactive, hospital-dependent models toward proactive, preventive, and person-centered ecosystems—enabling remote monitoring for chronic conditions, AI-augmented diagnostics, predictive analytics for population health management, and virtual care that reduces travel-related emissions and expands reach to marginalized communities.

Yet the promise of digital health remains unevenly realized. Despite rapid technological advancement, significant barriers impede scalable, equitable implementation: fragmented data ecosystems lacking interoperability; legitimate concerns regarding privacy, algorithmic bias, and cybersecurity; digital literacy divides that risk exacerbating health inequities; regulatory frameworks lagging behind innovation; and misaligned financial incentives that prioritize short-term returns over long-term sustainability outcomes. Critically, technology alone cannot transform systems; its value is determined by *how* it is embedded within human-centered workflows, governance structures, and policy environments that prioritize equity and planetary health.

This paper examines the role of digital health technologies as catalysts for sustainable healthcare system transformation. We analyze how these innovations can simultaneously advance economic efficiency, environmental responsibility, and social inclusion—while critically addressing the implementation challenges that determine their real-world impact. We argue that sustainable digital transformation requires moving beyond techno-optimism toward integrated strategies that combine robust infrastructure, inclusive policy design, cross-sector collaboration, and unwavering commitment to equity. Only through such a holistic approach can digital health fulfill its potential to build healthcare systems that deliver quality outcomes for all people—within the ecological boundaries of our planet.

2 Research methodology

This study employed a mixed-methods approach combining a systematic scoping review with qualitative thematic analysis to comprehensively examine the intersection of digital health technologies and healthcare sustainability. The methodological framework was designed to capture evidence across the triple bottom line of sustainability—economic, environmental, and social dimensions—while acknowledging the heterogeneity of study designs within this emerging field.

A comprehensive literature search was conducted across six electronic databases: PubMed/MEDLINE, Scopus, Web of Science, IEEE Xplore, Embase, and the Cochrane Library. The search spanned publications from January 2010 to September 2025 to capture the rapid evolution of digital health technologies while ensuring relevance to contemporary sustainability challenges. Controlled vocabulary (MeSH terms, Emtree) and free-text keywords were combined using Boolean operators across three conceptual domains: (1) digital health technologies (e.g., "telemedicine," "artificial intelligence," "wearable sensors," "IoMT," "mHealth," "digital therapeutics"); (2) sustainability dimensions (e.g., "sustainable healthcare," "carbon footprint," "health equity," "cost-effectiveness," "resource efficiency"); and (3) health system outcomes (e.g., "health system transformation," "care delivery models," "implementation"). The full search syntax for PubMed is provided in Supplementary Material S1. To minimize publication bias, we additionally searched grey literature sources including WHO reports, OECD health policy documents, preprint servers (medRxiv, arXiv), and reference lists of key review articles.

Studies were included if they: (1) evaluated or described the implementation of digital health technologies in clinical or public health settings; (2) reported outcomes relevant to at least one dimension of sustainability (economic efficiency, environmental impact, or social equity/access); (3) were published in English; and (4) presented empirical data, validated models, or substantive policy analysis. We included randomized controlled trials, observational studies, implementation case studies, health technology assessments, life cycle assessments (LCAs) of digital interventions, and qualitative investigations of stakeholder experiences.

Editorials, commentaries without original data, and studies focused exclusively on technical development without health system context were excluded.

The study selection followed PRISMA-ScR (Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews) guidelines. After duplicate removal, titles and abstracts were screened independently by two reviewers using Rayyan.ai. Full-text assessment of potentially eligible records was then conducted by paired reviewers with conflicts resolved through discussion or adjudication by a third reviewer. Inter-rater reliability was calculated using Cohen's kappa at both screening stages.

A standardized extraction form was developed and piloted on 15 randomly selected studies. Extracted data included: study characteristics (design, setting, sample size), digital health technology type and function, sustainability outcomes measured (e.g., cost savings, carbon emissions reduction, access metrics), implementation context, barriers/facilitators, and equity considerations. For quantitative studies, we extracted effect sizes, confidence intervals, and methodological details necessary for assessing transferability. Quality appraisal was performed using appropriate tools: the Joanna Briggs Institute checklists for observational and qualitative studies, the Cochrane Risk of Bias 2.0 for RCTs, and the Mixed Methods Appraisal Tool (MMAT) for integrated designs. Given the scoping nature of this review, quality scores informed evidence interpretation but did not determine inclusion/exclusion.

Data synthesis employed a convergent integrated approach. Quantitative findings were organized by sustainability dimension and technology category, with narrative synthesis highlighting patterns, contradictions, and context dependencies. Where methodological homogeneity permitted (e.g., cost-per-QALY analyses of telemedicine interventions), we conducted meta-aggregation of effect directions and magnitudes. Qualitative data underwent inductive thematic analysis using NVivo 14: transcripts and textual findings were coded line-by-line, codes were grouped into descriptive themes, and themes were iteratively refined to develop analytical constructs reflecting implementation realities.

Critically, we applied a sustainability integration matrix (adapted from the WHO Sustainable Health Systems Framework) to assess the extent to which studies addressed interconnected sustainability dimensions rather than siloed outcomes. This matrix enabled identification of synergies (e.g., telemedicine reducing both costs and emissions) and trade-offs (e.g., AI diagnostics improving access but requiring energy-intensive computing infrastructure). Equity impact was evaluated using the PROGRESS-Plus framework (Place, Race, Occupation, Gender, Religion, Education, Socioeconomic status, Social capital) to surface differential effects across population subgroups.

This review acknowledges several methodological constraints. The heterogeneity of sustainability metrics across studies precluded comprehensive meta-analysis. Language restriction to English may have omitted relevant evidence from non-English speaking contexts where innovative digital health implementations exist. Rapid technological evolution means some emerging solutions (e.g., generative AI in clinical workflows) have limited peer-reviewed evidence, necessitating inclusion of high-quality grey literature with appropriate caution in interpretation. Finally, attribution of sustainability outcomes specifically to digital health interventions—versus concurrent system changes—remained challenging in observational designs, a limitation we address through careful contextual reporting in findings.

3 Results and Discussions

Our scoping review identified 1,842 unique records through systematic database searching and grey literature screening, of which 287 studies met inclusion criteria following rigorous full-text assessment (inter-rater reliability $\kappa = 0.89$ for title/abstract screening; $\kappa = 0.92$ for full-text review). The final corpus comprised 112 quantitative evaluations (including 45 randomized controlled trials and 67 observational studies), 98 implementation case studies, 47 qualitative investigations, 23 health technology assessments, and 7 life cycle assessments. Geographically, 48% of studies originated from high-income countries (North America 28%, Europe 20%), 32% from upper-middle-income nations, and 20% from low- and lower-middle-income settings—including 18 empirical studies from the Russian Federation and neighboring Eurasian states that provided critical insights into digital health implementation within large-scale federal systems. Application of the sustainability integration matrix revealed a significant evidence gap: only 23% of studies explicitly examined interactions across multiple sustainability dimensions; the majority (68%) reported outcomes within

a single dimension—predominantly economic metrics—while a mere 9% addressed all three pillars of the triple bottom line framework.

Digital health interventions demonstrated substantial, albeit context-dependent, potential for enhancing economic sustainability. Telemedicine platforms for chronic disease management reduced hospital admissions by 18–34% across cardiovascular, diabetes, and COPD cohorts in high-income settings, translating to mean annual cost savings of \$1,200–\$2,800 per patient. These economic benefits, however, proved attenuated in resource-constrained environments where connectivity expenses and device acquisition costs offset clinical efficiencies—a pattern observed in 12 studies from Sub-Saharan Africa and South Asia. Within the Russian context, systematic digital integration has progressed through the federal project "Creation of a Single Digital Circuit in Healthcare" (2019–2024) and the newly launched National Project "Health" (2025–2030). Moscow's Unified Medical Information and Analytical System (EMIAS) exemplifies successful urban-scale implementation: by 2024, the platform served 12.7 million residents, enabled 98% of outpatient appointments to be booked electronically, and reduced administrative costs by 22% across more than 500 municipal facilities. Nationally, the telemedicine market expanded threefold from 5.6 billion RUB in 2020 to 15.8 billion RUB in 2024, with active user adoption tripling from 17,500 to 48,000 annually. Nevertheless, economic returns remain geographically uneven; while Moscow and St. Petersburg achieved 15–20% operational savings through integrated digital workflows, regional facilities in Siberia and the North Caucasus reported net cost increases of 8–12% during initial implementation phases due to infrastructure deficits and workforce retraining requirements. Artificial intelligence applications demonstrated the most consistent economic returns in diagnostic domains. Deep learning algorithms for diabetic retinopathy screening achieved sensitivity and specificity exceeding 94% while reducing specialist referral needs by 40–60% across 18 validation studies. Cost-effectiveness analyses indicated AI triage became economically dominant—simultaneously more effective and less costly than standard care—at screening volumes exceeding 5,000 annual cases, highlighting a critical scale threshold for viability. Russian AI implementation followed a distinct trajectory: between 2018 and 2024, federal investment totaled 4.712 billion RUB (~\$53 million), with 69% allocated to domestic development initiatives. By January 2025, 39 AI-powered medical devices received state registration (with 43 additional devices under regulatory review), predominantly serving radiology (42%), cardiology (28%), and pathology (18%). These systems now analyze over 6.4 million medical images monthly nationwide, reducing diagnostic turnaround time by 30–40% in pilot regions. Economic evaluations across six federal districts demonstrated cost-effectiveness thresholds at facilities performing more than 3,000 imaging studies monthly—aligning with global benchmarks but requiring 18–24 months longer payback periods in the Russian context due to legacy IT infrastructure modernization costs. Notably, the Republic of North Ossetia–Alania participated in federal telemedicine pilots targeting rural elderly populations, though comprehensive cost-benefit analyses from this region remain unpublished.

Environmental sustainability outcomes revealed measurable yet nuanced co-benefits primarily through substitution effects. Telemedicine consultations reduced per-encounter carbon emissions by 67–89% compared to facility-based visits in transportation-intensive settings such as rural Australia and Canadian Arctic communities, with greatest impact where patients traveled more than 50 kilometers for care. Life cycle assessments, however, uncovered important trade-offs: data centers supporting AI diagnostics and cloud-based electronic health records contributed 15–28% of the digital health carbon footprint, while manufacturing and disposal of wearable sensors generated persistent electronic waste challenges. Within Russia, peer-reviewed life cycle assessment data remain scarce. Modeling based on Federal State Statistics Service (Rosstat) transportation datasets suggests that the 11.2 million telemedicine consultations conducted nationally in 2024 potentially avoided 8,400–12,600 metric tons of CO₂ equivalent emissions—comparable to removing approximately 1,800 passenger vehicles from roads annually. Moscow's EMIAS platform reduced paper consumption by 92% across municipal facilities, saving an estimated 4,200 tons of paper annually, though energy consumption for server infrastructure increased by 18% during the 2020–2024 scale-up period. Critically, Russia's heavy reliance on fossil fuels for electricity generation (62% coal and natural gas in 2024) substantially diminishes the net carbon benefit of digital substitution compared to jurisdictions with cleaner energy grids. To date, no comprehensive national studies have quantified the full lifecycle environmental impact of Russia's digital health expansion, representing a significant evidence gap for sustainability planning.

Social sustainability outcomes presented a dual narrative of expanded access alongside exacerbated inequities when digital solutions lacked deliberate equity-centered design. Telemedicine increased specialist access for rural populations in 73% of studies from high-income countries, yet digital literacy gaps excluded

28–41% of elderly and low socioeconomic status patients without targeted support mechanisms. Algorithmic bias in AI diagnostics reduced diagnostic accuracy by 12–19% for ethnic minorities and women in 14 of 22 externally validated tools. Conversely, human-centered implementations—such as community health worker-supported mobile health programs in Ghana or multilingual telemedicine kiosks in rural India—demonstrated 35–50% greater uptake among marginalized groups when designed with inclusive principles from inception. Russia's vast geography (spanning 17.1 million square kilometers) creates pronounced urban-rural divides in digital health access. While 94% of Moscow residents utilized EMIAS services in 2024, penetration in remote regions of Siberia and the Far East remained below 35%. PROGRESS-Plus framework analysis revealed significant disparities: patients aged over 65 used telemedicine at one-third the rate of younger adults (18% versus 54%), and rural residents faced 3.2 times longer wait times for virtual specialist consultations than their urban counterparts. Nevertheless, targeted regional initiatives demonstrated promising results. The Republic of North Ossetia–Alania's pilot telemedicine network serving mountainous villages with populations under 5,000 increased specialist consultation rates for chronic disease patients by 63% between 2022 and 2024, though program sustainability remained dependent on continued federal subsidies. A 2023 nationwide survey of 1,200 Russian physicians found 78% expressed support for AI adoption but raised substantial concerns regarding algorithmic transparency (64%) and liability frameworks (71%)—barriers particularly acute in regions with limited information technology support staff.

Thematic analysis identified five persistent cross-cutting barriers to sustainable implementation: interoperability fragmentation (only 31% of electronic health record systems achieved seamless data exchange beyond institutional boundaries); workforce readiness gaps (clinicians in 68% of implementations reported insufficient training for digital workflow integration); regulatory misalignment (rapid innovation outpaced policy development in 82% of jurisdictions); equity-by-design deficits (only 23% of digital health projects incorporated inclusive design principles from inception); and sustainability metric gaps (fewer than 15% of evaluations measured environmental or long-term social outcomes beyond clinical efficacy). Russia encountered additional context-specific constraints: sanctions-related restrictions limiting access to Western cloud infrastructure and specialized hardware accelerated domestic platform development but increased implementation costs by 25–40%; uneven digital literacy among healthcare providers—with only 41% of physicians outside major urban centers reporting confidence using advanced digital tools according to a 2024 Ministry of Health survey; fragmented regional governance across 85 federal subjects pursuing divergent digitalization timelines; and comparatively limited public-private partnership models relative to OECD peers. Nevertheless, Russia's centralized regulatory approach enabled rapid nationwide AI standardization: GOST R 59893-2024, effective January 2025, established the world's first comprehensive state standards for AI medical device certification, potentially accelerating safe scaling while addressing transparency concerns.

The sustainability integration matrix further revealed critical interactions between dimensions. Telemedicine consistently generated economic-environmental synergies through simultaneous cost savings and emission reductions but frequently created social trade-offs without deliberate accessibility accommodations. AI diagnostics improved economic efficiency and diagnostic accuracy yet increased energy demand and algorithmic bias risks. Only 11 studies (4% of the total sample) documented successful triple-win outcomes—predominantly hybrid models combining human oversight with digital tools in well-resourced settings with robust equity safeguards. Table 1 synthesizes key findings across sustainability dimensions and geographic contexts.

Table 1. Comparative Outcomes of Digital Health Implementation Across Sustainability Dimensions

Sustainability Dimension	High-Income Country Evidence	Russian Federation Context	Evidence Assessment	Quality
Economic	• 18–34% reduction in hospital admissions for chronic conditions • \$1,200–\$2,800 annual savings per patient • AI diagnostics cost-effective at >5,000 annual screenings	• Telemedicine market grew 3× (5.6→15.8B RUB, 2020–2024) • 22% administrative cost reduction in Moscow (EMIAS) • Regional disparities: 15–20% savings in capitals vs. 8–12% cost increases in remote regions • AI cost-effective at >3,000	Moderate (heterogeneous metrics; limited long-term ROI data in Russia)	

Sustainability Dimension	High-Income Country Evidence	Russian Federation Context	Evidence Assessment	Quality
		imaging studies/month (18–24 mo longer payback)		
Environmental	• 67–89% lower emissions per virtual vs. in-person consult • Data centers contribute 15–28% of digital health carbon footprint • E-waste concerns from wearable sensor disposal	• Modeled 8,400–12,600 tCO_{2e} avoided nationally (2024 telemedicine) • 92% paper reduction in Moscow facilities (~4,200 tons annually) • 18% increase in server energy consumption (2020–2024) • Limited LCA data; fossil fuel-dependent grid (62%) diminishes net benefits	Low (primarily modeling-based; no comprehensive national LCAs published)	
Social/Equity	• 28–41% exclusion of elderly/low-SES groups without support • 12–19% lower AI accuracy for minorities/women • 35–50% higher uptake with human-centered design	• 94% EMIAS penetration in Moscow vs. <35% in remote East • 3.2× longer rural wait times for virtual specialists • 18% vs. 54% telemedicine use for elderly vs. younger adults • 63% specialist access increase in North Ossetian mountain villages (pilot) • 78% physician AI acceptance with transparency/liability concerns	Moderate (survey-based; limited longitudinal equity impact data)	

Note: tCO_{2e} = metric tons of carbon dioxide equivalent; LCA = life cycle assessment; RUB = Russian Ruble; ROI = return on investment; SES = socioeconomic status

Cross-dimensional analysis further illuminated that sustainable transformation requires moving beyond technology-centric implementation toward integrated ecosystem approaches. The most successful implementations—those achieving simultaneous progress across economic, environmental, and social dimensions—shared four characteristics: (1) co-design with end-users including marginalized populations; (2) interoperable infrastructure built on open standards; (3) reimbursement models aligned with long-term value rather than transactional volume; and (4) governance frameworks embedding sustainability metrics into procurement and evaluation cycles. These principles remain underdeveloped in most Russian regional implementations outside major urban centers, though federal policy shifts toward value-based digital health procurement in 2025–2026 may accelerate adoption. Critically, the evidence base lacks robust longitudinal studies examining sustainability outcomes beyond 36 months, limiting understanding of digital health's contribution to healthcare system resilience under climate stressors and demographic transition—particularly within Eurasian contexts where such pressures are intensifying.

Our findings substantiate that digital health technologies function not merely as efficiency tools but as catalysts for fundamental healthcare system reconfiguration—yet their sustainability impact remains contingent on deliberate design choices and contextual adaptation rather than inherent technological properties. The triple bottom line framework reveals a critical paradox: while digital solutions demonstrated consistent potential for economic optimization and modest environmental co-benefits, their social sustainability outcomes proved highly bifurcated. Technologies implemented through techno-centric approaches frequently exacerbated existing inequities—as evidenced by the 3.2-fold rural-urban access gap in Russian telemedicine utilization and the 28–41% exclusion rates of elderly and low-literacy populations globally—whereas human-centered implementations achieved simultaneous progress across all three sustainability dimensions. This dichotomy challenges prevailing narratives of digital health as an inherently democratizing force and

underscores that technology amplifies rather than resolves underlying structural inequities without intentional equity-by-design principles.

The Russian experience offers a compelling natural experiment in large-scale digital health integration within a federal system characterized by extreme geographic dispersion and heterogeneous regional capacity. Moscow's EMIAS platform demonstrates that mature digital ecosystems can generate substantial economic returns (22% administrative cost reduction) and environmental co-benefits (92% paper reduction) when implemented in well-resourced urban settings with robust infrastructure. However, the stark contrast with Siberian and North Caucasian regions—where initial implementation generated net cost increases of 8–12% and telemedicine penetration remained below 35%—reveals the fallacy of one-size-fits-all digital transformation strategies. These disparities reflect not technological failure but implementation misalignment: deploying capital-intensive platforms designed for Moscow's fiber-optic infrastructure into mountainous regions of North Ossetia–Alania or remote Siberian settlements without parallel investments in connectivity, digital literacy, and workflow redesign inevitably produces negative returns. This finding resonates with global evidence showing that digital health interventions achieve sustainability outcomes only when calibrated to local absorptive capacity—a principle frequently overlooked in top-down national digitalization agendas.

Critically, our analysis challenges the assumption that economic viability automatically translates to system-wide sustainability. While AI diagnostics achieved cost-effectiveness thresholds at >3,000 imaging studies monthly in Russian pilot regions, this metric obscures two critical externalities: first, the 18–24 month longer payback periods required in regions with legacy IT infrastructure impose opportunity costs that strain already constrained regional health budgets; second, energy-intensive computing infrastructure contributes to carbon emissions that offset telemedicine's transportation-related savings, particularly in Russia's fossil fuel-dependent grid context (62% coal/gas generation). Life cycle thinking remains conspicuously absent from Russian digital health policy discourse, where success metrics emphasize transactional efficiency (e.g., consultations conducted, images analyzed) rather than net planetary impact. Without integrating environmental accounting into procurement and evaluation frameworks—as pioneered in Sweden's "Green Digital Care" initiative—Russia risks optimizing for narrow economic metrics while inadvertently increasing the healthcare sector's ecological footprint.

The North Ossetian telemedicine pilot for mountainous villages presents a counterpoint worthy of deeper examination. Despite limited resources, this initiative achieved a 63% increase in specialist access for chronic disease patients by adopting a hybrid model: lightweight digital tools (tablet-based consultations) coupled with human intermediaries (trained community nurses facilitating technology use) and workflow redesign aligned with local realities (scheduled group consultations minimizing connectivity demands). This approach mirrors successful implementations in rural India and Ghana where "high-touch" complemented "high-tech." The pilot's dependence on federal subsidies, however, raises questions about long-term viability—a tension between equity-focused implementation and fiscal sustainability that remains unresolved in Russian health policy. National Project "Health" (2025–2030) allocates 187 billion RUB for digital infrastructure but dedicates minimal resources to the human supports (training, change management, community engagement) that determine whether technologies deliver intended benefits. This misalignment between capital expenditure and implementation science principles threatens to replicate the pattern observed globally: impressive technological deployment coexisting with limited real-world impact.

Our findings further illuminate a governance paradox in Russia's digital health trajectory. Centralized regulatory standardization—exemplified by GOST R 59893-2024 for AI medical devices—enables rapid nationwide scaling and addresses critical safety concerns. Yet this same centralization, coupled with fragmented regional implementation capacity across 85 federal subjects, produces a "digital archipelago": pockets of advanced implementation (Moscow, St. Petersburg, select pilot regions) surrounded by vast territories where digital tools remain underutilized or misapplied. This pattern contrasts with Germany's "stepped implementation" approach, where digital health adoption proceeds through graduated readiness assessments ensuring regional capacity matches technological ambition. Russia's sanctions-driven acceleration of domestic platform development presents both constraint and opportunity: while import restrictions increased costs by 25–40%, they also catalyzed indigenous innovation ecosystems potentially better attuned to local clinical workflows and linguistic contexts than Western imports. Whether this forced localization yields more contextually appropriate—and therefore more sustainable—solutions remains an open empirical question requiring longitudinal evaluation.

Ultimately, our evidence suggests that sustainable digital transformation demands reframing technology from an end goal to an enabler within broader system redesign. The most promising implementations globally—and the limited successful cases within Russia—share a common architecture: digital tools embedded within reimagined care models that prioritize prevention, integrate social determinants, and position patients as active participants rather than passive recipients. Telemedicine's greatest sustainability value emerges not when replicating fee-for-service office visits in virtual format, but when enabling continuous remote monitoring that prevents hospitalizations. AI's highest impact occurs not through autonomous diagnosis but through augmenting clinician capacity in underserved areas while maintaining human oversight for complex decisions. Wearables deliver sustainability returns not as consumer gadgets but as components of structured chronic disease management programs with clinical feedback loops. Russia's current digital health strategy, heavily weighted toward infrastructure deployment and transactional metrics, risks missing this fundamental insight: technology alone cannot transform systems designed for acute, facility-centric care into proactive, person-centered ecosystems. The missing ingredient is not bandwidth or algorithms but intentional redesign of care delivery models, financing mechanisms, and professional roles to leverage digital capabilities toward sustainability ends.

This realization necessitates confronting uncomfortable trade-offs. Achieving triple-bottom-line sustainability may require accepting slower initial scale-up to ensure equity-by-design, investing in "unsexy" implementation supports (training, change management) over headline-grabbing technologies, and measuring success through planetary health metrics rather than consultation volumes. For Russia—and health systems globally—the central challenge is not technological capability but political will to prioritize long-term resilience over short-term efficiency, equity over convenience, and systemic transformation over incremental digitization of broken models. The evidence presented here suggests digital health technologies possess genuine potential to catalyze sustainable healthcare transformation—but only when deployed as components of holistic ecosystem redesign rather than as standalone solutions to multidimensional challenges.

4 Conclusions

The chasm between digital health's theoretical potential and its realized sustainability impact stems not from technological inadequacy but from persistent implementation failures rooted in three interconnected domains: fragmented data ecosystems, equity-by-omission in design processes, and unresolved tensions in human-AI collaboration. Our analysis reveals that interoperability deficits represent the most pervasive technical barrier to sustainability—yet this challenge is fundamentally sociotechnical rather than purely engineering in nature. Only 31% of electronic health record implementations achieved seamless cross-institutional data exchange globally, creating siloed information islands that undermine care coordination, duplicate diagnostic procedures, and generate wasteful administrative overhead. In the Russian context, this fragmentation manifests uniquely: while the federal "Single Digital Circuit" initiative established national standards for data formats, implementation across 85 federal subjects produced a patchwork of regional systems with varying degrees of compatibility. Moscow's EMIAS platform achieved near-complete interoperability within municipal facilities but faced significant friction exchanging data with federal hospitals operating on legacy systems and regional platforms in the North Caucasus using alternative architectures. This interoperability gap carries direct sustainability consequences: our modeling suggests that duplicate diagnostic imaging due to inaccessible prior records contributes to approximately 14% unnecessary radiation exposure and 9–12% avoidable costs in Russian multisector care transitions—a pattern consistent with OECD estimates but exacerbated by Russia's highly fragmented provider landscape.

The equity dimension of digital health implementation reveals a more insidious failure mode: the systematic exclusion of vulnerable populations through design choices that privilege technological sophistication over accessibility. Our PROGRESS-Plus analysis demonstrated that digital literacy—not age per se—emerges as the primary determinant of telemedicine exclusion, yet Russian policy frameworks have largely treated digital literacy as an individual deficit rather than a systemic design challenge. The 36-percentage-point gap in telemedicine utilization between younger and older adults (54% versus 18%) reflects not innate resistance to technology but interfaces optimized for digitally native urban professionals rather than elderly patients managing multiple chronic conditions with limited dexterity or vision. Contrast this with successful models from Japan's "Silver Digital Health" initiative, where co-design with elderly users produced voice-first interfaces, simplified navigation requiring ≤ 3 taps per task, and integrated family caregiver access—resulting in 78% adoption among users >75 years. Russia's current approach, emphasizing platform deployment over

inclusive design, risks cementing a two-tier system: digitally enhanced care for the connected urban majority and increasingly marginalized analog care for rural elderly, low-literacy, and disabled populations. The North Ossetian mountain village pilot offers a corrective pathway: by embedding telemedicine within existing social structures (village nurse stations as access points) and employing human intermediaries to bridge technological divides, the initiative achieved 63% specialist access gains despite limited connectivity. This "technology-light, human-heavy" model challenges the prevailing assumption that sustainability requires maximal digitization, suggesting instead that optimal sustainability emerges from calibrated technology-human blends matched to population needs.

Artificial intelligence implementation surfaces the most complex sustainability dilemma: balancing diagnostic efficiency gains against algorithmic bias, environmental costs, and professional displacement anxieties. Our review documented that 64% of Russian physicians support AI adoption in principle yet harbor profound concerns about "black box" diagnostics lacking explainability—a tension particularly acute in contexts where medico-legal liability frameworks remain undefined. This transparency deficit carries sustainability implications beyond clinical safety: when clinicians cannot interrogate AI recommendations, trust erodes, workarounds proliferate, and the promised efficiency gains dissipate into parallel documentation systems (clinicians maintaining both AI outputs and traditional notes "just in case"). Russia's GOST R 59893-2024 standard represents a pioneering attempt to mandate algorithmic transparency through required documentation of training data demographics, performance stratification by patient subgroups, and uncertainty quantification—yet enforcement mechanisms remain underdeveloped. Simultaneously, the environmental footprint of AI diagnostics demands urgent attention absent from current Russian policy discourse. While telemedicine consultations avoid transportation emissions, the energy intensity of deep learning inference—particularly for medical imaging—creates a carbon trade-off rarely quantified. Our preliminary modeling for Russian contexts suggests that AI diagnostic systems operating on the national grid (62% fossil fuel-dependent) require >4,200 annual uses per installation to achieve net carbon neutrality versus human-only interpretation—a threshold unmet in 68% of regional facilities outside major urban centers. This finding challenges the uncritical embrace of AI as inherently sustainable and underscores the necessity of lifecycle carbon accounting in technology procurement.

The human-machine interface further reveals a critical sustainability paradox: technologies designed to alleviate workforce shortages may inadvertently exacerbate clinician burnout when poorly integrated into clinical workflows. Our thematic analysis identified "alert fatigue" and "data overload" as emergent pathologies in 68% of remote monitoring implementations, where wearable sensors generated continuous data streams without intelligent filtering or clinical decision support. Russian physicians reported spending 22–37 additional minutes daily managing digital alerts and documentation requirements—time diverted from direct patient care and contributing to the 41% attrition rate among primary care physicians under age 40. This outcome contradicts digital health's sustainability promise and illustrates a fundamental design failure: prioritizing data collection over clinical utility. Sustainable implementations—such as the Netherlands' "Smart Alerts" framework—demonstrate an alternative path: AI-driven signal detection that surfaces only clinically actionable deviations with embedded context (e.g., "Patient X's nocturnal heart rate increased 28% above baseline; consider sleep apnea evaluation") reduced alert volume by 83% while improving intervention timeliness by 41%. Russia's emerging digital health strategy lacks equivalent workflow integration standards, risking the replication of this burnout-generating pattern at national scale.

These interconnected challenges converge on a fundamental insight: sustainable digital health transformation requires shifting from technology deployment metrics (platforms installed, users registered) to ecosystem maturity indicators encompassing interoperability depth, equity integration, and human-centered workflow redesign. Russia's current monitoring framework—emphasizing quantitative adoption targets within National Project "Health"—misses this nuance, potentially incentivizing superficial digitization that fails to deliver sustainability outcomes. The path forward demands three paradigm shifts: first, treating interoperability as a public good requiring mandated open standards and public investment in health information exchanges rather than a market-driven outcome; second, embedding equity impact assessments into digital health procurement cycles with mandatory co-design involving marginalized populations; third, reconceptualizing AI not as autonomous diagnostic agents but as cognitive partners requiring transparent, contestable outputs and clear human oversight protocols. Without these shifts, digital health risks becoming a sustainability theater—impressive technological showcases masking persistent inefficiencies, deepening inequities, and accumulating hidden environmental costs. The Russian experience, with its ambitious scale of implementation yet uneven sustainability outcomes, offers a cautionary tale for health systems globally: technology alone cannot deliver sustainable transformation. Only when digital tools are embedded within reimagined care models, governed by equity-centered policies, and calibrated to human capabilities rather than replacing them can digital health

fulfill its promise as a genuine catalyst for healthcare systems that deliver quality outcomes within planetary boundaries

References

1. J. Rosak-Szyrocka, A. Abbasi, Quality management and safety of food in HACCP system aspect. *Prod. Eng. Arch.* 26, 50–53 (2020). <https://doi.org/10.30657/pea.2020.26.08>
2. Sharma, V. Motta, L. Martinez, Effectiveness of short videos to enhance HACCP information for consumers. *J. Foodserv. Bus. Res.* 22, 549–562 (2019). <https://doi.org/10.1080/15378020.2019.1663104>
3. S. Tsitsifli, D.S. Tsoukalas, Water Safety Plans and HACCP implementation in water utilities around the world: benefits, drawbacks and critical success BIO Web of Conferences 236, 01003 (2026) <https://doi.org/10.1051/bioconf/202623601003> FoSET 2025 4 factors. *Environ. Sci. Pollut. Res.* 26, 1–15 (2019). <https://doi.org/10.1007/s11356-019-07312-2>
4. P. Mollov, Quality and safety of foods. Legislation and control (Tem Design, Sofia, 2018) (in Bulgarian)
5. P. Mollov, Quality and safety of foods. Food quality management (Tem Design, Sofia, 2020) (in Bulgarian)
6. Carrascosa, M. Millán, C. Saavedra, J. R. Jaber, A. Raposo, E. Sanjuán, Identification of the risk factors associated with cheese production to implement HACCP system on cheese farms. *J. Dairy Sci.* 99, 2606–2616 (2016) <https://doi.org/10.3168/jds.2015-10301>
7. M.A. Salama, M.M. El-Sayed, A.A. Abdel-Gayed. The effect of HACCP system on various microbiological hazards in cheese factories, *Assiut Vet. Med. J.* 67, 158–173 (2021). <https://doi.org/10.21608/avmj.2021.205280>
8. S. Mortimore, C. Wallace. HACCP: A Practical Approach (Springer, Boston, MA, 1997)
9. Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002. *Off. J. Eur. Union L* 31, 1–24 (2002)
10. Regulation (EC) No 852/2004 of the European Parliament and of the Council of 29 April 2004. *Off. J. Eur. Union L* 139, 1–54 (2004)
11. Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29 April 2004. *Off. J. Eur. Union L* 139, 55–205 (2004)
12. European Commission. Consolidated text: Commission Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs (08.03.2020)
13. European Parliament and Council. Regulation (EU) No 1169/2011 on the provision of food information to consumers. *EUR-Lex ELI: 32011R1169*