


Forecasting COVID-19 mortality in the US Midwest

Guido España, Rachel Oidtman*, Sean Cavany*, Alan Costello,
Annaliese Wieler, Anita Lerch, Carly Barbera, Marya Poterek,
Quan Tran, Sean Moore, and T. Alex Perkins
University of Notre Dame, United States

University of Notre Dame


August 4th, 2020

We based our model on the agent-based model platform: FRED



- Developed at the University of Pittsburgh (Grefenstette et al., 2013).
- FRED is an agent-based model framework highly flexible and adaptable.
- We modified FRED to simulate COVID-19, some of the modifications include
 - ▶ natural history of disease parameters,
 - ▶ implemented adaptive non-pharmaceutical interventions.

Agents can transmit the virus in specific places



- Infectious individuals might spread the virus where susceptible individuals visit the same places the same day.

Probability of symptoms and death





Figure: From (Verity et al., 2020; The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team, 2020).

Infectious period and duration of symptoms






Figure: From (He et al., 2020; Lauer et al., 2020; Bi et al., 2020)

We focused on seven Midwest States




- We used deaths to calibrate the model parameters: shelter-in-place compliance, transmission efficiency, importation scaling, and asymptomatic relative infectivity.
- We simulate 2,000 parameter sets for each state.

We estimated the trend in people staying at home from the google mobility trends




- After the data period, we assumed a linear trend for the forecasting horizon.

Calibrated parameters




- Parameters were estimated for each state.

Model calibration to deaths




- Cumulative deaths match the reported number of deaths in each state.

Forecast: we assume shelter in place stays with the current trend, schools reopen.




- Large impact of schools. Currently working on school reopen strategies.


Reproduction number increases after interventions are relaxed and schools are open at full capacity



Evolution of forecast



Evolution of forecast



Evolution of forecast

COVID-19 Forecasts

Week Ahead


WEEK 21 (2020)

Indiana

TARGET

Cumulative Deaths


Cumulative Deaths (Observed)



0.1 1 10 1e+2 1e+3 1e+4

Time Chart

The **ensemble** forecast combines models unconditional on particular interventions being in place with those conditional on certain social distancing measures continuing. To ensure consistency, only models with 4 week-ahead forecasts ahead are included in the ensemble.



Evolution of forecast

Proportion of tests positive

COVID-19 Forecasts

Week Ahead


WEEK 22 (2020)

Indiana



TARGET

Cumulative Deaths

Cumulative Deaths (Observed)



0.1 1 10 1e+2 1e+3 1e+4



conditional on particular interventions being in place with those conditional on certain social distancing only models with 4 week-ahead forecasts ahead are included in the ensemble.

Evolution of forecast

COVID-19 Forecasts

Week Ahead


WEEK 23 (2020)

Indiana

TARGET


Cumulative Deaths

Cumulative Deaths (Observed)




0.1 1 10 1e+21e+31e+4

Daily deaths




unconditional on particular interventions being in place with those conditional on certain social distancing
only models with 4 week-ahead forecasts ahead are included in the ensemble.



Evolution of forecast

Only forecasting the incidence of deaths after the date of submission



Evolution of forecast

After several updates

COVID-19 Forecasts

Week Ahead


WEEK 28 (2020)

Indiana

TARGET


Cumulative Deaths

Cumulative Deaths (Observed)




Time Chart

The **ensemble** forecast combines models unconditional on particular interventions being in place with those conditional on certain social distancing measures continuing. To ensure consistency, only models with 4 week-ahead forecasts ahead are included in the ensemble.




To improve fit in Indiana, we included age-structure data



- Lower susceptibility in <20 based on (Davies et al., 2020).
- We used the number of cases in long-term care facilities and estimated importations to these facilities.
- We calibrated the model to the age structure of deaths in Indiana and validated to serological studies.

To improve fit in Indiana, we included testing and hospitalizations



Future work

- Adding face-masks to improve fit.
- We are using this model to assess school reopening scenarios in Indiana.

Funding statement

This research is being funded by an NSF RAPID grant: Real-time updating of an agent-based model to inform COVID-19 mitigation strategies. Award number: 2027718.

References I

- Grefenstette, J. J., Brown, S. T., Rosenfeld, R., DePasse, J., Stone, N. T. B., Cooley, P. C., ... Sriram, A., et al. (2013). FRED (A Framework for Reconstructing Epidemic Dynamics): An open-source software system for modeling infectious diseases and control strategies using census-based populations. *BMC public health*, 13(1), 940.
- Verity, R., Okell, L. C., Dorigatti, I., Winskill, P., Whittaker, C., Imai, N., ... Ferguson, N. M. (2020). Estimates of the severity of coronavirus disease 2019: A model-based analysis. *The Lancet. Infectious Diseases*.
doi:10.1016/S1473-3099(20)30243-7

References II

- The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. (2020). The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) — China, 2020. *China CDC Weekly*, 2(8), 113–122.
- He, X., Lau, E. H. Y., Wu, P., Deng, X., Wang, J., Hao, X., ... Leung, G. M. (2020). Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nature Medicine*, 1–4. doi:10.1038/s41591-020-0869-5
- Lauer, S. A., Grantz, K. H., Bi, Q., Jones, F. K., Zheng, Q., Meredith, H. R., ... Lessler, J. (2020). The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Annals of Internal Medicine*. doi:10.7326/M20-0504

References III

- Bi, Q., Wu, Y., Mei, S., Ye, C., Zou, X., Zhang, Z., ... Zhang, T., et al. (2020). Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: A retrospective cohort study. *The Lancet Infectious Diseases*.
- Davies, N. G., Klepac, P., Liu, Y., Prem, K., Jit, M., Pearson, C. A. B., ... CMMID COVID-19 working group. (2020). Age-dependent effects in the transmission and control of COVID-19 epidemics. *Nature Medicine*.
doi:10.1038/s41591-020-0962-9