

# Problems

February 9, 2017

1. Why is L2 norm a vector norm?
2. Verify that the Cauchy-Schwarz inequality holds for

$$\mathbf{u} = (1, 3, 5, 2, 0, 1), \mathbf{v} = (0, 2, 4, 1, 3, 5).$$

3. Let  $W_1$  and  $W_2$  be a subspaces of  $\mathbb{R}^n$ . Prove that the intersection  $W_1 \cap W_2$  is a subspace of  $\mathbb{R}^n$ .
4. Prove that if  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is a linearly independent set, then so is the set  $\{k\mathbf{v}_1, k\mathbf{v}_2, k\mathbf{v}_3\}$  for every nonzero scalar  $k$ .
5. Find all values of  $\lambda$  for which  $\det(A) = 0$ .

$$A = \begin{bmatrix} \lambda - 4 & 0 & 0 \\ 0 & \lambda & 2 \\ 0 & 3 & \lambda - 1 \end{bmatrix}$$

6. Find the determinant of the following matrix by inspection, not by calculation.

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 2 & 2 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 4 \end{bmatrix}.$$

7. Find the eigenvalues and corresponding eigenvectors of the matrix

$$A = \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix}.$$

8. (a) Compute the operator norm of the following matrix.

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}.$$

- (b) What is the vector achieving the following maximum?

$$\max_{\|\mathbf{x}\| \neq 0} \frac{\|A\mathbf{x}\|}{\|\mathbf{x}\|}$$

9. Show that the vectors

$$\begin{aligned} \mathbf{v}_1 &= (1, 0, 0), \quad \mathbf{v}_2 = (0, 2, 0), \\ \mathbf{v}_3 &= (0, 0, 3), \quad \mathbf{v}_4 = (1, 1, 1) \end{aligned}$$

span  $\mathbb{R}^3$  but do not form a basis for  $\mathbb{R}^3$ .

10. Prove that if  $\mathbf{u}$  is a nonzero  $n \times 1$  column vector, then the outer product  $\mathbf{u}\mathbf{u}^T$  is a symmetric matrix of rank 1.
11. Prove that if  $V = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$  is a linearly independent set of vectors in  $\mathbb{R}^n$ , and if  $W = \{\mathbf{w}_1, \dots, \mathbf{w}_{n-k}\}$  is a basis for the null space of the matrix  $A$  that has the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  as its successive rows, then  $V \cup W$  is a basis for  $\mathbb{R}^n$ .
12. Find the least squares solution of  $A\mathbf{x} = \mathbf{b}$  by solving the associated normal system, and show that the least square error vector is orthogonal to the column space of  $A$ .

$$A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \\ 4 & 5 \end{bmatrix}; \quad \mathbf{b} = \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix}.$$

13. Find the eigenvalue decomposition of

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

14. Find a singular value decomposition of  $A$ .

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}.$$

15. Suppose that  $A$  has the singular value decomposition

$$A = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 24 & 0 & 0 \\ 0 & 12 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \\ -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \end{bmatrix}.$$

- (a) Find orthonormal bases for the four fundamental spaces of  $A$ .  
 (b) Find the reduced singular value decomposition of  $A$ .
16. Prove that nuclear norm has the following property:

$$\|A\|_* \triangleq \sum_{i=1}^{\min(m,n)} \sigma_i = \text{tr}(\sqrt{A^T A}),$$

for  $A \in \mathbb{R}^{m \times n}$ , where  $\sqrt{A^T A}$  is a matrix  $B$  such that  $B^2 = A^T A$ .