

An "**ENCOUNTER**" of advanced airway management refers to complete sequence of events leading to a placement of an advanced airway

A "**COURSE**" of advanced airway management refers to ONE method or approach to secure an airway **AND ONE** set of medications (including premedication and induction).

Each course may include one or several "attempts" by one or several providers.

An "**ATTEMPT**" is a single advanced airway maneuver (e.g. tracheal intubation, LMA placement), beginning with the insertion of a device, e.g. laryngoscope (or LMA device) into patient's mouth or nose, and ending when the device (laryngoscope), LMA or tube is removed.

COURSE INFORMATION								
--------------------	--	--	--	--	--	--	--	--

Attempts for this COURSE	1	2	3	4	5	6	7	8
Who intubated (Fellow, Resident, etc)								
Discipline (ICU, ENT, Surgery, etc)								
PGY level (3 rd year resident = PL3, 1 st year fellow = PL4, NP=yrs as NP, etc.)								
ETT (or LMA) size								
ETT type: cuffed/uncuffed/ NA								
Immediately prior to this attempt was cricoid pressure/external laryngeal manipulation provided?								
During this attempt, was cricoid pressure/external laryngeal manipulation provided?								
Attempt Successful: Yes / No								

Difficult to Bag – Mask Ventilate? (Circle ONE only):	_____
---	-------

Difficult Airway Evaluations (Choose/Circle one in each category):

1. Evaluation done before or after this course is completed?	
2. Known prior history of difficult airway?	
3. Any Limited Neck Extension or (Maximal with or without sedation/paralytics) Severe Reduction (e.g. trauma patient with collar)	
4. Widest Mouth Opening – How many <u>Patient's</u> fingers between gum/incisors	
5. Thyromental space – Patient's fingers between chin and thyroid cartilage	
6. Evidence of Upper Airway Obstruction or Anatomical Barrier to visualize glottic opening (Subjective assessment before looking)?	
7. Midfacial Hypoplasia?	
8. Any other signs of difficult airway exist?	
If YES Please Explain:	

Known cyanotic heart disease (R to L shunt)? (Circle ONE only)	
--	--

Medications:
☐ NO DRUGS USED (If no drugs used, select box and go to next section)

Pretreatment Dosage	Paralysis Dosage	Induction Dosage
[] mg Atropine (check unit!)	[] mg Rocuronium	[] mg Propofol
[] mcg Glycopyrrolate	[] mg Succinylcholine	[] mg Etomidate
[] mcg Fentanyl	[] mg Vecuronium	[] mg Ketamine
[] mg Lidocaine	[] mg Pancuronium	[] mg Midazolam
[] mg Vecuronium	[] mg Cisatracurium	[] mg Thiopental
Others:	Others:	Others:

Atropine Indication:**Glycopyrrolate Indication:**

Method: Begin NEW course if NEW method / device used (please use new form):	
---	--

--

Apneic Oxygenation Use:

1. Was Oxygen provided DURING any TI attempts for this course?	
2. If Yes, How was the oxygen provided:	

	Liter Flow	FiO2

Device (Check only ONE) Begin NEW course if NEW method / device used.

--

(Not part of medical record call x5193 if found)