

Area of Cyclic Polygons by Elimination of Variables

Tanay Sonthalia, Connor Haynes, Joshua Jayprakash

Georgia Institute of Technology

November 25, 2024

Table of Contents

- 1 Heron's Formula
- 2 Brahmagupta's Formula
- 3 Gauss' Pentagon Formula
- 4 Automatic Theorem Proving

Table of Contents

- 1 Heron's Formula
- 2 Brahmagupta's Formula
- 3 Gauss' Pentagon Formula
- 4 Automatic Theorem Proving

Heron's Formula

Q: Can we calculate the area of a polygon given only its side lengths?

Heron's Formula

Q: Can we calculate the area of a polygon given only its side lengths?

A: Sometimes.

Heron's Formula

Q: Can we calculate the area of a polygon given only its side lengths?

A: Sometimes.

Q: When is “sometimes”?

Heron's Formula

Q: Can we calculate the area of a polygon given only its side lengths?

A: Sometimes.


Q: When is “sometimes”?

A: When it is *rigid* (when its area is fixed uniquely by its side lengths).

Heron's Formula

Example

A triangle is rigid, but a quadrilateral is not.



$$K_1 \neq K_2$$

Heron's Formula

So, because all triangles are rigid, we can show

Heron's Formula (1st Century BC)

Given a triangle T with side lengths a, b, c , the area K of T is given by

$$K = \sqrt{s(s-a)(s-b)(s-c)},$$

where s is the *semiperimeter*, $s = \frac{1}{2}(a + b + c)$.

Table of Contents

- 1 Heron's Formula
- 2 Brahmagupta's Formula
- 3 Gauss' Pentagon Formula
- 4 Automatic Theorem Proving

Brahmagupta's Formula

One way of ensuring rigidity in polygons is requiring that all vertices lie on a circle. These are *cyclic polygons*.

Brahmagupta's Formula

One way of ensuring rigidity in polygons is requiring that all vertices lie on a circle. These are *cyclic polygons*.


Brahmagupta's Formula describes the area of a cyclic quadrilateral.

Brahmagupta's Formula

Given: Sidelengths a, b, c, d


Brahmagupta's Formula

Given: Sidelengths a, b, c, d




Brahmagupta's Formula

Create the circle described by these four points



Brahmagupta's Formula

Inscribe the quadrilateral



Brahmagupta's Formula

Brahmagupta's Formula (7th Century)

Given sidelengths a, b, c, d , the area K of the cyclic convex quadrilateral with these side lengths is given by


$$K = \sqrt{(s-a)(s-b)(s-c)(s-d)},$$

where s is the *semiperimeter*, $s = \frac{1}{2}(a + b + c + d)$.


Table of Contents

- 1 Heron's Formula
- 2 Brahmagupta's Formula
- 3 Gauss' Pentagon Formula**
- 4 Automatic Theorem Proving


Gauss' Pentagon Formula




Gauss' Pentagon Formula




Gauss' Pentagon Formula




Gauss' Pentagon Formula



Gauss' Pentagon Formula



Gauss' Pentagon Formula



Gauss' Pentagon Formula

Gauss' Pentagon Formula (1823)

Given sidelengths a, b, c, d, e , the area K of the cyclic pentagon with these sidelengths is satisfies

$$K^2 - K(b_0 + b_1 + b_2 + b_3 + b_4) + b_0b_1 + b_1b_2 + b_3b_4 + b_4b_0 = 0,$$

where each b_i is the area of some distinct triangle formed by three consecutive vertices.

Table of Contents

- 1 Heron's Formula
- 2 Brahmagupta's Formula
- 3 Gauss' Pentagon Formula
- 4 Automatic Theorem Proving

Automatic Theorem Proving

The goal of this project is to prove Heron-type formulas for the area of cyclic polygons by automatic means.

Automatic Theorem Proving

The goal of this project is to prove Heron-type formulas for the area of cyclic polygons by automatic means.

This can be done by constructing an ideal using polynomial relations given by the cyclic polygon,

Automatic Theorem Proving

The goal of this project is to prove Heron-type formulas for the area of cyclic polygons by automatic means.

This can be done by constructing an ideal using polynomial relations given by the cyclic polygon,

(a) Side lengths: $a_i^2 - (x_{i+1} - x_i)^2 + (y_{i+1} - y_i)^2 = 0$

(b) Circle: $x_i^2 + y_i^2 - r^2 = 0$

(c) Area: $K = \frac{1}{2} \left(\begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \end{vmatrix} + \cdots + \begin{vmatrix} x_n & y_n \\ x_1 & y_1 \end{vmatrix} \right)$

then eliminating all variables except for the area and the side lengths.

Automatic Theorem Proving

Procedure:

(1) Create a polynomial ring over \mathbb{Q}

Automatic Theorem Proving

Procedure:

- (1) Create a polynomial ring over \mathbb{Q}
- (2) Set $(x_1, y_1) = (r, 0)$ and define other vertex points as indeterminates $(x_2, y_2), \dots, (x_n, y_n)$

Automatic Theorem Proving

Procedure:

- (1) Create a polynomial ring over \mathbb{Q}
- (2) Set $(x_1, y_1) = (r, 0)$ and define other vertex points as indeterminates $(x_2, y_2), \dots, (x_n, y_n)$
- (3) Define area and distance relations

Automatic Theorem Proving

Procedure:

- (1) Create a polynomial ring over \mathbb{Q}
- (2) Set $(x_1, y_1) = (r, 0)$ and define other vertex points as indeterminates $(x_2, y_2), \dots, (x_n, y_n)$
- (3) Define area and distance relations
- (4) Define an ideal I generated by these relations

Automatic Theorem Proving

Procedure:

- (1) Create a polynomial ring over \mathbb{Q}
- (2) Set $(x_1, y_1) = (r, 0)$ and define other vertex points as indeterminates $(x_2, y_2), \dots, (x_n, y_n)$
- (3) Define area and distance relations
- (4) Define an ideal I generated by these relations
- (5) Eliminate all variables other than side lengths and area

Automatic Theorem Proving

Procedure:

- (1) Create a polynomial ring over \mathbb{Q}
- (2) Set $(x_1, y_1) = (r, 0)$ and define other vertex points as indeterminates $(x_2, y_2), \dots, (x_n, y_n)$
- (3) Define area and distance relations
- (4) Define an ideal I generated by these relations
- (5) Eliminate all variables other than side lengths and area
- (6) Factor the principal generator of I , set each factor to zero and solve

Automatic Theorem Proving

Procedure:

- (1) Create a polynomial ring over \mathbb{Q}
- (2) Set $(x_1, y_1) = (r, 0)$ and define other vertex points as indeterminates $(x_2, y_2), \dots, (x_n, y_n)$
- (3) Define area and distance relations
- (4) Define an ideal I generated by these relations
- (5) Eliminate all variables other than side lengths and area
- (6) Factor the principal generator of I , set each factor to zero and solve
- (7) All done!

Automatic Theorem Proving

Procedure:

- (1) Create a polynomial ring over \mathbb{Q}
- (2) Set $(x_1, y_1) = (r, 0)$ and define other vertex points as indeterminates $(x_2, y_2), \dots, (x_n, y_n)$
- (3) Define area and distance relations
- (4) Define an ideal I generated by these relations
- (5) Eliminate all variables other than side lengths and area
- (6) Factor the principal generator of I
- (7) All done!

Why??

Automatic Theorem Proving

How could there be multiple distinct “area” functions?

Automatic Theorem Proving

How could there be multiple distinct “area” functions?

We must investigate what assumptions we make that we are not providing the computer.

Automatic Theorem Proving

How could there be multiple distinct “area” functions?

We must investigate what assumptions we make that we are not providing the computer.

In all the proofs of the given theorems, edges are placed to form a *convex* polygon. This information is not encapsulated in the ideal.

Automatic Theorem Proving

How could there be multiple distinct “area” functions?

We must investigate what assumptions we make that we are not providing the computer.

In all the proofs of the given theorems, edges are placed to form a *convex* polygon. This information is not encapsulated in the ideal.

Q: How else might these relations be interpreted?

Automatic Theorem Proving

How could there be multiple distinct “area” functions?

We must investigate what assumptions we make that we are not providing the computer.

In all the proofs of the given theorems, edges are placed to form a *convex* polygon. This information is not encapsulated in the ideal.

Q: How else might these relations be interpreted?

A: These edges may form nonconvex polygons.


Examples

There are two “combinatorial configurations” of a quadrilateral:

Automatic Theorem Proving

Examples

There are two “combinatorial configurations” of a quadrilateral:




Note: the above quadrilateral configurations do not share the same side lengths.

Automatic Theorem Proving

Examples

There are two “combinatorial configurations” of a quadrilateral:




Note: the above quadrilateral configurations do not share the same side lengths.

When factored, the principal generator of the corresponding ideal yields two polynomials: one that agrees with Brahmagupta's Formula and one that does not.

Automatic Theorem Proving

Examples

There are two “combinatorial configurations” of a quadrilateral:



Note: the above quadrilateral configurations do not share the same side lengths.

When factored, the principal generator of the corresponding ideal yields two polynomials: one that agrees with Brahmagupta's Formula and one that does not.

These correspond to the area of the convex and nonconvex quadrilaterals, respectively.

Automatic Theorem Proving

Without restrictions on the order in which vertices are connected, there exist

$$\Delta_n = \frac{1}{2} \left((2(n-3) + 1) \binom{2(n-3)}{n-3} - 2^{2(n-3)} \right)$$

combinatorial configurations of cyclic n -gons, where $n \geq 5$.

Automatic Theorem Proving

Without restrictions on the order in which vertices are connected, there exist

$$\Delta_n = \frac{1}{2} \left((2(n-3) + 1) \binom{2(n-3)}{n-3} - 2^{2(n-3)} \right)$$

combinatorial configurations of cyclic n -gons, where $n \geq 5$.

This corresponds to the number of factors in the principal generator of I , and each of these factors yields the area of a distinct configuration.

Automatic Theorem Proving

Without restrictions on the order in which vertices are connected, there exist

$$\Delta_n = \frac{1}{2} \left((2(n-3) + 1) \binom{2(n-3)}{n-3} - 2^{2(n-3)} \right)$$

combinatorial configurations of cyclic n -gons, where $n \geq 5$.

This corresponds to the number of factors in the principal generator of I , and each of these factors yields the area of a distinct configuration.

The first few terms of this sequence are

$$\Delta_5 = 7, \Delta_6 = 38, \Delta_7 = 187, \Delta_8 = 874, \dots$$

And so computational requirements grow quickly.

Automatic Theorem Proving

With this procedure we were able to verify all three stated formulas, as well as derive formulas in agreement with the findings of Robbins for the area of nonconvex 4- and 5-gons.

1. F. Miller Maley, David P. Robbins, and Julie Roskies, *On the areas of cyclic and semicyclic polygons*, Advances in Applied Mathematics **34** (2005), no. 4, 669-689 (en).
2. D.P. Robbins, *Areas of Polygons Inscribed in a Circle*, Discrete and Computational Geometry **12** (1994), no. 2, 223-236.