

Polytopes, Amoeba, and Tropical Curves

Connections in Tropical Geometry

C. Haynes J. Whidden

GT Directed Reading Program, Fall 2024

Table of Contents

1 Tropical Curves

2 Polytopes

3 Amoebas

4 Conclusion

Definition

The *tropicalization* of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is given by its image in the *tropical semiring* T .

Definition

The *tropicalization* of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is given by its image in the *tropical semiring* T .

Basically, take addition and make it minimization, then take multiplication and make it addition.

Tropical Curves

Definition

The *tropicalization* of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is given by its image in the *tropical semiring* T .

Basically, take addition and make it minimization, then take multiplication and make it addition.

Example

The tropicalization of $f = x_1 + x_2 + 1$ is given by

Tropical Curves

Definition

The *tropicalization* of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is given by its image in the *tropical semiring* T .

Basically, take addition and make it minimization, then take multiplication and make it addition.

Example

The tropicalization of $f = x_1 + x_2 + 1$ is given by

$$\text{trop}(f) = \min\{x_1, x_2, 1\}.$$


Tropical Curves

Example

The tropicalization of $f = x_1 + x_2 + 1$ is given by

$$\text{trop}(f) = \min\{x_1, x_2, 1\}.$$

Its tropical curve is



These are the points where $\text{trop}(f)$ has its minimum achieved twice.


Tropical Curves

Example

The tropicalization of $f = x_1 + x_2 + 1$ is given by

$$\text{trop}(f) = \min\{x_1, x_2, 1\}.$$

Its tropical curve is



These are the points where $\text{trop}(f)$ has its minimum achieved twice.

This constitutes one construction of a *tropical curve*.

Table of Contents

1 Tropical Curves

2 Polytopes

3 Amoebas

4 Conclusion

Definition

A polytope is the convex hull of some finite set of points in \mathbb{R}^n .

Definition

A polytope is the convex hull of some finite set of points in \mathbb{R}^n .

The *Newton Polytope* of a polynomial $f = \sum_k a_k x^{\mathbf{u}_k} \in \mathbb{C}[x_1, \dots, x_n]$ is the polytope in \mathbb{R}^n given by the convex hull of all $\mathbf{u}_k = (u_{k_1}, u_{k_2}, \dots, u_{k_n})$.

Definition

A polytope is the convex hull of some finite set of points in \mathbb{R}^n .


The *Newton Polytope* of a polynomial $f = \sum_k a_k x^{\mathbf{u}_k} \in \mathbb{C}[x_1, \dots, x_n]$ is the polytope in \mathbb{R}^n given by the convex hull of all $\mathbf{u}_k = (u_{k_1}, u_{k_2}, \dots, u_{k_n})$.

Here $x^{\mathbf{u}} = x_1^{u_1} x_2^{u_2} \dots x_n^{u_n}$.

Example

The Newton Polytope $\text{newt}(f)$ of $f = x_1 + x_2 + 1$ is the convex hull of

$$\{(1, 0), (0, 1), (0, 0)\} :$$




Definition

The *inner normal fan* of a polytope Σ is the polyhedral fan containing the normal cones to each face of Σ .


Example

The inner normal fan of $\text{newt}(f)$, where $f = x_1 + x_2 + 1$, is




Example

The inner normal fan of $\text{newt}(f)$, where $f = x_1 + x_2 + 1$, is



Example

The inner normal fan of $\text{newt}(f)$, where $f = x_1 + x_2 + 1$, is



This constitutes another construction of a *tropical curve*.

Table of Contents

1 Tropical Curves

2 Polytopes

3 Amoebas

4 Conclusion

Definition

The *variety* $\mathbb{V}(f)$ of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is the set of points $v \in \mathbb{C}^n$ such that $f(v) = 0$.

Definition

The *variety* $\mathbb{V}(f)$ of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is the set of points $v \in \mathbb{C}^n$ such that $f(v) = 0$.

Varieties contain exactly those points on which f vanishes.

Definition

The *variety* $\mathbb{V}(f)$ of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is the set of points $v \in \mathbb{C}^n$ such that $f(v) = 0$.

Varieties contain exactly those points on which f vanishes.

Definition


The *amoeba* of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is the image of $\mathbb{V}(f)$ under the local logarithmic map

$$\text{Log} : (\mathbb{C}^*)^n \rightarrow \mathbb{R}^n,$$

$$\text{Log}(v_1, \dots, v_n) = (\log |v_1|, \dots, \log |v_n|).$$



Example

The amoeba of $f = x_1 + x_2 + 1$ is the image of $\mathbb{V}(f)$ under Log:





Example

The amoeba of $f = x_1 + x_2 + 1$ is the image of $\mathbb{V}(f)$ under Log :



Example

The amoeba of $f = x_1 + x_2 + 1$ is the image of $\mathbb{V}(f)$ under Log :



The *spine* of this amoeba is another construction of a *tropical curve*.

Table of Contents

1 Tropical Curves

2 Polytopes

3 Amoebas

4 Conclusion



Conclusion

Maybe you've noticed a pattern?

Conclusion

Maybe you've noticed a pattern?

Idea!



Question: Why do these objects coincide?

Question: Why do these objects coincide?

Keywords: Tropical Geometry, Maslov Dequantization, Log Semiring,
Logarithmic Limit Set, Newton Polytope

Book: *Introduction to Tropical Geometry* - Diane Maclagan and
Bernd Sturmfels