

Fulton's *Algebraic Curves*

Connor Haynes

§ 1 Affine Algebraic Sets

1.1 Algebraic Preliminaries

1.4: For k infinite, the only polynomial that is zero everywhere is the zero polynomial.

Proof. Let $F \in k[X_1, \dots, X_n]$ be a polynomial such that $F(a_1, \dots, a_n) = 0$ for all $a_1, \dots, a_n \in k$. We proceed by induction on the number of variables n . For the case $n = 0$ the statement holds trivially. Now suppose the statement holds for $n - 1$. Then we may write $F = \sum F_i X_n^i$ for $F_i \in k[X_1, \dots, X_{n-1}]$. Now suppose that one of the F_i is not equal to zero. Then there are only finitely many roots for the polynomial $F(a_1, \dots, a_{n-1}, X_n)$, as it is univariate. However the field k is infinite so this polynomial cannot possibly vanish everywhere on k . Therefore $F = 0$. $\triangle$

1.2 Affine Space and Algebraic Sets

1.3 The Ideal of a Set of Points

1.21: An ideal generated by linear polynomials is maximal and the quotient is isomorphic to the field of coefficients

Proof. Suppose $I = (X_1 - a_1, \dots, X_n - a_n)$, and let $J \supseteq I$. Then I is the ideal of the point $\mathbf{a} = (a_1, \dots, a_n)$ and $V(J) \subseteq V(I)$. However $V(I) = \{\mathbf{a}\}$, so $J = I$. That is, I is maximal. Now take the quotient $k[X_1, \dots, X_n]/I$. Now the only ideals in the quotient are the whole ring and the zero ideal, which implies that the ring is a field. The only field contained in $k[X_1, \dots, X_n]$ is k , so the quotient must be isomorphic to this field. $\triangle$

Proposition 1: An algebraic set is irreducible if and only if its ideal is prime.

Proof. First the backward direction. Suppose $I(V)$ not prime. Then there is $fg \in I$ such that $f \notin I$ and $g \notin I$. Then $V(f) \cap V \cup V(g) \cap V = V$, and so V is reducible.

Now the forward direction. Suppose $V = U \cup W$ is reducible. Then for $f \in I(U) \setminus I$ and $g \in I(W) \setminus I$ we have that $fg \in I$, so I is not prime. $\triangle$

1.4 The Hilbert Basis Theorem

1.22: Bijections between ideals of R/I and ideals containing I .

Proof. Part (a) asks us to prove the above statement for any ideals. First the forward direction. If J' is an ideal of R/I then $\pi^{-1}(J')$ is in particular closed under multiplication and addition of generators by the fact that π is a homomorphism. That is, it is an ideal. The fact that it contains I follows by the fact that $0 \in J'$ and $\pi^{-1}(0) = I$.

Part (b) asks us to prove that radicality, primality, and maximality are preserved. First radicality. Supposing J' is radical, we have $f^m \in J'$ implies that $f \in J'$. Then taking preimages, $\pi^{-1}(f^m) = f^m + I = (f + I)^m = \pi^{-1}(f)$, so $\pi^{-1}(J')$ is radical. Likewise, if J is radical then $\pi(J)$ is obviously radical because $\pi(f^m) = \pi(f)^m$ and $\pi(f) \in \pi(J)$. Now primality. If J' is prime then $\pi^{-1}(J') = J$ is because $\pi^{-1}(fg) = fg + I = (f + I)(g + I)$ and one of $f + I$ or $g + I$ is in the preimage. Conversely, if J is prime then multiplicativity of the homomorphism shows that J' is as well. For maximality, note that projection preserves containment.

For part (c), note that if J is finitely generated then we may write any element of J' as

$$\pi(x) = \pi\left(\sum c_\alpha y_i\right) = \sum \pi(c_\alpha) \pi(y_i)^\alpha$$

for y_i generators of J . $\triangle$

1.5 Irreducible Components of an Algebraic Set

1.6 Algebraic Subsets of the Plane

1.7 Hilbert's Nullstellensatz

1.38: Bijections between algebraic subsets of an algebraic set $V = V(I)$ and radical ideals of $k[X_1, \dots, X_n]/I$.

Proof. The Nullstellensatz demonstrates bijections between algebraic sets and radical ideals, and 1.22(a-b) (above) demonstrates the necessary bijections once one notes the inclusion-reversing properties of $I(\cdot)$ and $V(\cdot)$ as functors (J contains I exactly when $V(J)$ is a subset of $V(I)$) and recalls that irreducible varieties have prime ideals, while points have maximal ideals. $\triangle$

1.8 Modules: Finiteness Conditions

1.9 Integral Elements

1.10 Field Extensions

§ 2 Affine Varieties

2.1 Coordinate Rings

2.1: Show that the map $k[X_1, \dots, X_n]$ associating to each F a polynomial function in $\mathcal{F}(V, k)$ is a ring homomorphism with kernel $I(V)$.

Proof. Let $\varphi : k[X_1, \dots, X_n] \rightarrow \mathcal{F}(V, k)$ assign each polynomial a polynomial function. This is obviously a ring homomorphism, and the kernel is shown to be $I(V)$ by noting that $f \rightarrow 0 \iff f(v) = 0$ for all $v \in V \iff f \in I(V)$. $\triangle$

2.2: Show that there is a natural one-to-one correspondence between algebraic subsets (resp. subvarieties, resp. points) and radical ideals (resp. prime ideals, resp. maximal ideals) of $\Gamma(V)$.

Proof. This follows by recalling that $\Gamma(V) = k[X_1, \dots, X_n]/I(V)$ and applying 1.22 and 1.38 (above). $\triangle$

2.3: Show that every polynomial function on V restricts to one on $W \subseteq V$.

Proof. For part (a), note that $I_V(W)$ is the image of $I(W)$ under the quotient $k[X_1, \dots, X_n]/I(V)$. Then obviously any polynomial on V restricts to one on W by the fact that polynomials are still polynomials under the quotient map. This map is surjective precisely because it is a quotient map $k[X_1, \dots, X_n]/I(V) \rightarrow k[X_1, \dots, X_n]/I(W)$ and $I(W) \supseteq I(V)$. The isomorphism follows because $I_V(W)$ is approximately $I(W)/I(V)$. $\triangle$

2.4: Show that the following are equivalent: V is a point, $\Gamma(V) = k$, and $\dim_k \Gamma(V) < \infty$.

Proof. The equivalence of (ii) and (iii) are the content of 1.21 (above). Corollary 4 of the Nullstellensatz tells us that V is finite, and because it is a variety it is irreducible and hence a point. (ii) implies (iii) straightforwardly as k is a k -vector space of dimension one. $\triangle$

2.2 Polynomial Maps

2.6: Show that the composition of pullbacks is pullback of the (reversed) composition, and that the composition of polynomial maps is a polynomial map.

Proof. Let $\varphi : V \rightarrow W$ and $\psi : W \rightarrow Z$. Now $\widetilde{\psi \circ \varphi}(F) = F \circ \psi \circ \varphi = \widetilde{\varphi}(F \circ \psi) = (\widetilde{\varphi} \circ \widetilde{\psi})(F)$. If φ and ψ are polynomial maps, then $\varphi = (T_1, \dots, T_n)$ and $\psi = (S_1, \dots, S_m)$, and from this it follows that $\psi \circ \varphi = (S_1(T_1, \dots, T_n), \dots, S_m(T_1, \dots, T_n))$. Now one can see that $\psi \circ \varphi$ is obviously polynomial by expanding. $\triangle$

2.7: Show that the preimage of an algebraic subset is algebraic, and that if the preimage of an algebraic set is irreducible, then so long as the set lives in the image it is also irreducible.

Proof. Let $\varphi : V \rightarrow W$ be a polynomial map and X an algebraic subset of W . Let $I_W(X)$ be the ideal of X in $\Gamma(W)$ and take $Y = V(F \in \Gamma(V) \mid F \circ \varphi \in I_W(X))$. Then $\varphi(Y) = X$. The claim on irreducibility follows by the fact that if Y is reducible then there will be some $F, G \in \Gamma(V)$ such that $FG \circ \varphi \in I_W(X)$ but neither will have pullback lying in the ideal, implying that it is not prime and so X is reducible. $\triangle$

2.9: Show that for a polynomial map of varieties $\varphi : V \rightarrow W$ and subvarieties V' and W' we have $\varphi(V') \subseteq W'$ implies that $\widetilde{\varphi}(I_W(W')) \subseteq I_V(V')$, and that the restriction gives a polynomial map $V' \rightarrow W'$.

Proof. Let $\varphi : V \rightarrow W$ be a polynomial map and $V' \subseteq V$, $W' \subseteq W$ subvarieties and $\varphi(V') \subseteq W'$. For part (a), we have that because $\varphi(V') \subseteq W'$, $V' \subseteq \varphi^{-1}(W')$. Then $\varphi^{-1}(W')$ is a variety by 2.7, and so we conclude that $I_V(V') \supseteq I_V(\varphi^{-1}(W'))$. Now note that any polynomial F vanishing on $\varphi^{-1}(W')$ satisfies $\widetilde{\varphi}(F) \in I_W(W')$. Therefore $I_W(W') \subseteq I_V(\varphi^{-1}(W'))$ and the result holds.

Part (b) seems more or less obvious: $\varphi(V') \subseteq W'$ means that the map is defined on all of V' , and it being polynomial at every point in V means that it is, in particular, polynomial at $V' \subseteq V$. Perhaps one must use Part (a) to check that nothing is vanishing when it shouldn't. $\triangle$

2.10: Show that the projection map $\mathbb{A}^n \rightarrow \mathbb{A}^r$ is polynomial.

Proof. This is the map $\varphi = (x_1, \dots, x_r)$. $\triangle$

2.3 Coordinate Changes

2.14: Show that affine changes of coordinates preserve linear subvarieties of $\mathbb{A}^n$ and that any nonempty linear subvariety may be expressed as the vanishing of some $X_{m+1}, \dots, X_n$. Show also that the m before is invariant of the choice of coordinate change.

Proof. Let $V \subseteq \mathbb{A}^n(k)$ be a linear subvariety (that is, $V = V(F_1, \dots, F_r)$ for F_i linear) and let $I = I(V)$. First we show part (a). Let T be an affine change of coordinates on $\mathbb{A}^n$. Now $V^T = V(F_1^T, \dots, F_r^T) = V(F_i(T_1, \dots, T_n) \mid 1 \leq i \leq r)$. Writing $F = \sum a_i X_i$ by linearity, we now have $F^T = \sum a_i T_i = \sum a_i \sum b_j X_j$ which is obviously a linear polynomial. Implicitly we are using the fact that I^T is generated by F_i^T , which is clear. Therefore V^T is linear.

For part (b) we proceed by induction on r . For $r = 1$ and $V = V(F)$, write $F = \sum a_i X_i$. Let j be the lowest index for which $a_j \neq 0$ so we may write $T_j = (X_j - \sum_{i \neq j} a_i X_i)/a_j$ and $T_i = X_i$ for $i \neq j$. Then $F^T = F(T_1, \dots, T_n) = \sum_{i \neq j} a_i X_i + a_j (X_j - \sum_{i \neq j} a_i X_i)/a_j = X_j$. Now consider S for which $S_j = X_n$ and $S_i = X_i$ for all other $i \neq j$. Now observe that $F^{T \circ S} = F(T_1(S_1, \dots, S_n), \dots, T_n(S_1, \dots, S_n)) = (F^T)(S_1, \dots, S_n) = (F^T)^S = X_j^S = X_n$. Then the base case of induction holds. Now suppose that for $r - 1$ the statement holds and consider $V = V(F_1, \dots, F_r)$. Now $V(F_1, \dots, F_{r-1})^T = V(X_{m+1}, \dots, X_n)$ for some m and some affine coordinate change T . Then F_r^T is a polynomial in $X_{m+1}, \dots, X_n$ and we may apply the above construction to produce a map S that sends F^T to X_m and fixes all other variables. Postcomposition of T with this map then yields the desired map.

For part (c), we suppose for the sake of contradiction that $T : X = V(X_{m+1}, \dots, X_n) \rightarrow Y = V(X_{s+1}, \dots, X_n)$ is an isomorphism for $m < s$. Let $I = (X_{m+1}, \dots, X_n)$ so that $I^T = (X_{s+1}, \dots, X_n)$. Now the induced map on coordinate rings $\widetilde{T} : \Gamma(Y) \rightarrow \Gamma(X)$ is an isomorphism.

However note that $I(X) = (X_{m+1}, \dots, X_n)$ and $I(Y) = (X_{s+1}, \dots, X_n)$, so the dimensions of the two coordinate rings differ as k vector spaces. This implies that there cannot be a bijection between the two, so in particular they are not isomorphic. $\triangle$

2.4 Rational Functions and Local Rings

2.18: Show that there is a natural one-to-one correspondence between prime ideals of $\mathcal{O}_P(V)$ and subvarieties of V passing through P .

Proof. Suppose I is prime in $\mathcal{O}_P(V)$ so that $I \cap \Gamma(V)$ is also prime ($fg \in I \cap \Gamma(V)$ implies that $\text{wlog } f \in I$ but $fg \in I \cap \Gamma(V)$ implies $\text{wlog } f$ is polynomial and clears g 's denominator). By the proof of Proposition 3 we have that $I \cap \Gamma(V)$ generates I . Now apply the bijection between prime ideals in $\Gamma(V)$ and subvarieties so that $I \cap \Gamma(V)$ gives a subvariety, and note that $P \in V(I \cap \Gamma(V))$ by the fact that every $h \in I$ is defined at P . $\triangle$

2.21: Show that a polynomial map of affine varieties induces a unique ring homomorphism $\mathcal{O}_Q(W) \rightarrow \mathcal{O}_P(V)$. Show that the maximal ideal of the codomain contains the image of the domain's maximal ideal.

Proof. Suppose $\varphi : V \rightarrow W$ is a polynomial map such that $P \in V$ has $\varphi(P) = Q$. Now consider the map on coordinate rings $\tilde{\varphi} : \Gamma(W) \rightarrow \Gamma(V)$. We wish to show that $\tilde{\varphi}$ extends uniquely to a homomorphism $\mathcal{O}_Q(W) \rightarrow \mathcal{O}_P(V)$. We define such a homomorphism by $\tilde{\varphi}(f/g) = \tilde{\varphi}(f)/\tilde{\varphi}(g)$. This map respects the equivalence relations as if $f/g \sim p/q$ then $fq - pg = 0$ and so $\tilde{\varphi}(fq - pg) = \tilde{\varphi}(f)\tilde{\varphi}(q) - \tilde{\varphi}(g)\tilde{\varphi}(p) = 0$ and obviously restricts to the map on coordinate rings. The fact that it is a homomorphism follows by the fact that it is a homomorphism on coordinate rings and the definition of the field of fractions. Uniqueness is clear by construction. In particular, $\tilde{\varphi}(\mathfrak{m}_Q(W)) \subseteq \mathfrak{m}_P(V)$ as if $f/g \in \mathfrak{m}_Q(W)$ vanishes at Q then in particular $f(Q) = 0$ implying that $f(\varphi(P)) = 0$ and so $\tilde{\varphi}(f)(P) = 0$. $\triangle$

2.22: Show that for an affine change of coordinates on $\mathbb{A}^n$, the map described above is an isomorphism and induces an isomorphism on varieties before and after the change of coordinates.

Proof. Suppose $T : \mathbb{A}^n \rightarrow \mathbb{A}^n$ is an affine change of coordinate with $T(P) = Q$. Notably the coordinate rings are isomorphic and so by construction of $\tilde{T}$ the local rings are isomorphic as well. If V is a subvariety of $\mathbb{A}^n$ such that $P \in V^T$, then the restriction to $\mathcal{O}_Q(V)$ has image given by F^T/G^T for all $F/G \in \mathcal{O}_Q(V)$. However this is precisely the field of fractions of V^T , so long as $P \in V^T$ to ensure that we restrict to where F^T is defined. $\triangle$

2.5 Discrete Valuation Rings

2.23: Show that the order function on K is independent the choice of a uniformizing parameter.

Proof. If t is a uniformizing parameter for R then any other uniformizing parameter is of the form $s = ut$ for $u \in R^\times$ a unit. Then if the order of $z \in K = \text{Frac}(R)$ is n we have $z = vt^n$ for some $v \in R^\times$. We may then write $z = vt^n = v(u^{-1}s)^n = ws^n$ for $w = vu^{-n}$ a unit. This expression in terms of the uniformizing parameter is unique, and so the order is fixed. $\triangle$

2.24: Show that the affine line has local ring given by a DVR.

Proof. First we show part (a). Let $f/g \in \mathfrak{m}_a(V) \subset \mathcal{O}_a(\mathbb{A}_k^1)$. Then f/g vanishes at a , implying that $f = h \cdot (X - a)$ for some $h \in \Gamma(V)$, and we may write $f = h \cdot (X - a)^n$ with h not divisible by $(X - a)$. In this case f/g permits an expression of the form $f/g = h/g \cdot (X - a)^n$, and h/g is a unit because neither is divisible by $(X - a)$. Importantly this unit is unique by construction, and so $\mathcal{O}_a(V)$ is a DVR.

Now part (b). We let $\mathcal{O}_\infty(V) = \{f/g \mid \deg(g) \geq \deg(f)\}$. Let $f/g \in \mathcal{O}_\infty(V)$ and suppose that we take a representative with such that X is not a factor of both f and g . Note that f/g is a unit if and only if $\deg f = \deg g$, as then $g/f \in \mathcal{O}_\infty(V)$. Now for any f/g we may take $X^n f/g$ to be a unit for n sufficiently large, implying that $f/g = X^n f/g \cdot (1/X)^n$. This expression is unique by construction, and so $\mathcal{O}_\infty(V)$ is a DVR. $\triangle$

2.26: Show that the maximal ideal of a DVR contains any element of R or its inverse. Show that if S is a DVR containing R , contained in the quotient field of R , and whose maximal ideal contains that of R , then $S = R$.

Proof. Part (a): Let $z \in K$ such that $z \notin R$. Then the order of z is negative, implying that $z = ut^n$ for $n < 0$. Now take an inverse to see that $z^{-1} = u^{-1}t^{-n}$, so in particular z has positive order implying that $z \in \mathfrak{m}$.

Part (b): Suppose $\mathfrak{m}_R \subseteq \mathfrak{m}_S$. Suppose $z \in S - R$. Then $z^{-1} \in \mathfrak{m}_R$ by (a), so $z^{-1} \in \mathfrak{m}_S$. However $z \in S$ so $zz^{-1} \in \mathfrak{m}_S$ which is a contradiction. Therefore $S = R$. $\triangle$

2.28: Show that a discrete valuation on a field gives a DVR by the elements of K with nonnegative valuation.

Proof. Let φ be a valuation on K . It is clear that R is a ring and $\mathfrak{m}$ is an ideal. Observe that $\mathfrak{m}$ is the set of non-units in R , and so R has a unique maximal ideal containing every proper ideal of R by the Lemma in 2.4, which is equal to $\mathfrak{m}$. That is, R is local. Suppose $x \in R$. We wish to show that $x = ut^n$ for some irreducible $t \in R$, some unit $u \in R^\times$, and some $n \geq 0$. Fix some irreducible $t \in \mathfrak{m}$ of minimal degree. Then any nonzero $y \in R$ has $0 \leq \varphi(yt^{-m}) < \varphi(t)$ for some m . This implies that yt^{-m} is a unit, as $\varphi(t)$ is minimal. Equivalently, $y = ut^m$ and this is a unique expression by construction.

The converse is obvious. $\triangle$

2.29: Show that if the order of two elements is different, their sum is equal to the minimum of the two orders. Show that if a collection of elements of K has an element of strictly minimal order then their sum is nonzero.

Proof. Part (a): If $\varphi(a) < \varphi(b)$ then $\varphi(a+b) \geq \min(\varphi(a), \varphi(b)) = \varphi(a)$.

Part (b): If there exists i such that $\varphi(a_i) < \varphi(a_j)$ for all $j \neq i$ then $\varphi(a_1 + \dots + a_n) = \varphi(a_i) \neq \infty$, so $a_1 + \dots + a_n \neq 0$. $\triangle$

2.30: Let R be a DVR with quotient field K , $k \subseteq R$ a field, and an isomorphism $k \rightarrow K \rightarrow R/\mathfrak{m}$. Show that for any $z \in R$ there is unique $\lambda \in k$ such that $z - \lambda \in \mathfrak{m}$ and show that any $z \in R$ admits an expression $z = \lambda_0 + \lambda_1 t + \dots + \lambda_n t^n + z_n t^{n+1}$ for $\lambda_i \in k$ and $z_n \in R$.

Proof. Part (a): Let $z \in R$. If $z \in \mathfrak{m}$ then $\lambda = 0$. Suppose $z \notin \mathfrak{m}$ and note that the coset $z + \mathfrak{m} \in k$ has additive inverse $-\lambda$ such that $z - \lambda = 0 \in k$. However $z - \lambda = 0 \in k$ implies that the preimage lives in $\mathfrak{m}$.

Part (b): To establish existence we proceed by induction on n . For the case $n = 0$ we write $z - \lambda \in \mathfrak{m}$ which implies that $z = \lambda \in k$, which in particular implies that $z = \lambda \in R$. Now suppose the result holds for $n = k - 1$, so there are $\lambda_0, \dots, \lambda_{k-1} \in k$ and $z_{k-1} \in R$ for which $z = \lambda_0 + \dots + \lambda_{k-1} t^{k-1} + z_{k-1} t^k$. Now we may write $z_{k-1} - \lambda_k \in \mathfrak{m}$ so $z = \lambda_k \in R$ as in the base case. Now $z = \lambda_0 + \dots + \lambda_k t^k$ with $z_k = 0$. So the inductive step holds.

For uniqueness, consider the difference of two such expressions $0 = z - z = (\lambda_0 - \mu_0) + \dots + (\lambda_n - \mu_n) t^n + (z_n - y_n) t^{n+1}$. Now note that the order of the term $\lambda_0 - \mu_0$ is minimal in this sum, so by 2.29 the sum cannot be zero. However this implies that $0 \neq 0$ so we have a contradiction, establishing uniqueness. $\triangle$

2.6 Forms

2.35: Count the dimension of forms of degree d as a k -vector space and show that sequences of nonzero linear forms (some not scalar multiples of others) generate the space of d -forms on two variables.

Proof. Part (a): A monomial of degree d in $R[X, Y]$ is of the form $X^i Y^j$ such that $i + j = d$. There are $d + 1$ pairs satisfying this, as can be seen by noting that $i = d - j$ is a solution for all $0 \leq j \leq d$. Similarly, in $R[X, Y, Z]$ we have as many monomials as triples $i + j + k = d$ and writing $i = d - j - k$ we see that the number of such triples is given by $\sum_{j=0}^d \sum_{k=0}^i 1 = \sum_{j=0}^d j = (d+1)(d+2)/2$.

Part (b) is obvious by the definition of forms.

Part (c): **Incomplete** $\triangle$

2.7 Direct Products of Rings

2.38: Show that the direct sum adds dimension.

Proof. The direct product is obviously generated as a ring by the generators of each R_i , and so there are $\dim R_i$ generators for each component. Components are independent, so there are $\sum \dim R_i$ generators. Equivalently, the dimension of the product is the sum of dimensions. $\triangle$

2.8 Operations with Ideals

2.39: Show that ideal multiplication is distributive and associative.

Proof. Part (a): $(I_1 + I_2)J = \{a + b \mid a \in I_1, b \in I_2\} \cdot J = \{(a + b)c \mid a \in I_1, b \in I_2, c \in J\} = \{ac + bc \mid \dots\} = I_1J + I_2J$.

Part (b): $(I_1 \dots I_m)^n = (\{a_1 \dots a_m \mid a_i \in I_i\})^n = \{(a_{1,1} \dots a_{1,n}) \dots (a_{1,1} \dots a_{1,m} \mid a_{i,j} \in I_i\} = I_1^n \dots I_m^n$. $\triangle$

2.40: Show that for I, J comaximal we have $I + J^2 = R$, I^m and J^n comaximal, and that products of comaximal ideals agree with their intersections, even when taking powers.

Proof. Part (a): If I, J are comaximal then $I + J = R$. Then $1 = a + b$ for $a \in I, b \in J$. Now $1 = 1^2 = a^2 + 2ab + b^2 = a(a + 2b) + b^2$. Now $a(a + 2b) \in I$ and $b^2 \in J^2$, so in particular $I + J^2 = R$. This naturally generalizes: $1 = 1^m = af + b^m$ for $f = \sum_{i+j=m} a^i b^j$ where the sum ranges over $0 \leq i < j \leq m - 1$. This shows that I and J^m are radical, and because our rings are commutative $I + J = J + I$ so we may perform the same process to obtain that $J^m + I^n = R$.

Part (b): The Lemma states that I, J comaximal implies that $IJ = I \cap J$, and part (a) shows that any powers of I and J are comaximal. Applying this to $J_i = \bigcap_{j \neq i} J_j$ we note that $I_1^n \cap \bigcap_{j=2}^m I_j^n = I_1^n \cdot \bigcap_{j=2}^m I_j^n = \dots = I_1^n \dots I_m^n = (I_1 \dots I_m)^n = (I_1 \cap \dots \cap I_m)^n$. $\triangle$

2.41: Show that if I is in the radical of an ideal, then some power of I is in the ideal (ChatGPT helped solved this one).

Proof. Suppose $I \subseteq \sqrt{J}$. Then for any $a \in I$ there is some minimal $n \geq 0$ such that $a^n \in J$. For $g_1, \dots, g_m$ generators of I we write $a = \sum b_i g_i$. Then $a^n = \sum b_1^{i_1} \dots b_m^{i_m} g_1^{i_1} \dots g_m^{i_m}$ where the sum is over all $i_1, \dots, i_m$ such that $\sum i_j = n$. Let m_i be the minimal power of g_i such that $g_i^{m_i} \in J$. Taking $n = \sum m_i$, we have that a^n is a sum of elements of the form $g_1^{i_1} \dots g_m^{i_m}$. Now because n is sufficiently large we have that, for any such product, $\sum i_j = n$ implying that at least one $i_j \geq m_j$. This implies that $g_j^{m_j} (g_1^{i_1} \dots g_j^{i_j - m_j} \dots g_m^{i_m}) \in J$, as J is an ideal. Therefore every product of the form $g_1^{i_1} \dots g_m^{i_m}$ such that $\sum i_j = n$ is in J , implying that $I^n \subseteq J$. $\triangle$

2.42: Show that there are natural homomorphisms between quotients: $R/I \rightarrow R/J$ for $I \subseteq J$ and $R/I \rightarrow S/IS$.

Proof. Part (a): If $I \subseteq J$, then R/I has a natural map onto R/J given by the identity map defined on cosets of I . For $a, b \in R$, $a \sim b$ in R/I if and only if $a - b \in I$ which implies that $a - b \in J$ so $a \sim b$ in R/J . Then the fact that we have a homomorphism follows by the fact that R/J is a ring.

Part (b): If $R \subseteq S$ is a subring then we have the natural homomorphism $R/I \rightarrow R/IS$ by the fact that $I \subseteq IS$. Now there is another natural injective homomorphism $R/IS \rightarrow S/IS$ given by the identity, and the composition of these two gives the desired natural map. $\triangle$

2.43: Show that the maximal ideal of the local ring of a point $P = (0, \dots, 0)$ is $(X_1, \dots, X_n)\mathcal{O}$ and that $\mathfrak{m}^r = I^r \mathcal{O}$.

Proof. The maximal ideal of a point is given by all non-units of $\mathcal{O}_P$, i.e. those vanishing at P . A polynomial with no constant term must vanish at P , and I consists of precisely these polynomials. Therefore and $f \in I\mathcal{O}$ has numerator vanishing at P , implying that $f \in \mathfrak{m}$. In fact all these statements are “if-and-only-if,” so the result holds. $\triangle$

2.44: Quotients of defining ideals in the local ring of a point.

Proof. **Incomplete** $\triangle$

2.45: Show that the intersection of two ideals is empty if and only if their defining ideals are comaximal.

Proof. Observe: $V(I) \cap V(J) = V(I + J) = \emptyset$ if and only if $I + J = R$. $\triangle$

2.46: Show that $\dim_k(k[X, Y]/(X, Y)^n) = n(n+1)/2$.

Proof. The power ideal I^n consists of all degree n polynomials in X and Y . Then $k[X, Y]/I^n$ is the set of all degree $d < n$ polynomials, which has dimension $n(n+1)/2$. $\triangle$

2.9 Ideals with a Finite Number of Zeros

No starred exercises.

2.10 Quotient Modules and Exact Sequences

2.48: Show that the image and kernel of an R -module homomorphism are R -modules and that they form an exact sequence.

Proof. The kernel and image are, in particular, subgroups for any group homomorphism, and the R -action obviously extends. To see exactness, note that the inclusion $\ker \varphi \subseteq M$ has image equal to the kernel of φ (big surprise). Therefore the sequence has $\text{im} \subseteq \ker \varphi$. $\triangle$

2.49: Show exactness for a number of natural sequences.

Proof. Part (a): Consider the map $\bar{\varphi}$ that takes cosets $a + N \in M/N$ to elements $\varphi(a) \in M'$. This is well-defined because $\varphi(N) = 0$. Then $\bar{\varphi} \circ \pi(a) = \bar{\varphi}(a + N) = \varphi(a)$. Uniqueness follows by construction: Any other map factoring φ must have that $\bar{\varphi}(\pi(a)) = \varphi(a)$, so in particular cosets of a must go to $\varphi(a)$.

Part (b): If $P \subseteq N \subseteq M$ our natural homomorphisms are achieved by the same argument as in 2.42. Now

$$0 \rightarrow N/P \xrightarrow{\subseteq} M/P \xrightarrow{\varphi} M/N \rightarrow 0$$

has $\text{im } \varphi = M/N$. Then $\ker \varphi = \{a \in M/P \mid a \in N\} = \{a \in N/P\} = \text{im } \subseteq$, so the sequence is exact.

Part (c): This is a consequence of (b) when $P \subseteq N \subseteq M$ are finite-dimensional vector spaces and an application of the rank-nullity theorem.

Part (d): Ideals are obviously modules. Then this follows from (b) taking $I = P \subseteq J = N \subseteq R = M$.

Part (e): This is the situation of (d) with $I = \mathfrak{m}^n \supseteq J = \mathfrak{m}^{n+1}$ and $R = \mathcal{O}$. $\triangle$

2.50: Show that a DVR as in 2.30 has $\dim_k(\mathfrak{m}^n/\mathfrak{m}^{n+1}) = 1$, $\dim_k(R/\mathfrak{m}^n) = n$, and $\text{ord } z = \dim_k(R/\mathfrak{m}^n)$.

Proof. Part (a): Recall that $\mathfrak{m} = (t)$. Then $\mathfrak{m}^n = (t^n)$. Now note that $\mathfrak{m}^n/\mathfrak{m}^{n+1} = t$. Therefore $\dim_k(\mathfrak{m}^n/\mathfrak{m}^{n+1}) = 1$.

Part (b): Writing $\mathfrak{m}^n = (t^n)$ we note that any element $z = ut^k$ of order greater than or equal to n lies in $\mathfrak{m}^n$, and so is mapped to zero by the quotient homomorphism. Therefore the only remaining terms are those of order $k < n$, yielding an n -dimensional vector space.

Part (c): If $(z) = \mathfrak{m}^n$ we use the fact that $\mathfrak{m}^n$ is generated by t^n for t a uniformizing parameter to see that $z = xt^n$ for some $x \in R$. However we also have that $zy = t^n$, so $z = xyz$ which implies that $xy = 1$ (recall that R is a domain). Now we see that z is of order n and so the result follows. $\triangle$

2.52: Let N, P be submodules of M . Show that the subgroup $N + P$ is a submodule of M . Show that there is a natural R -module isomorphism of $N/N \cap P$ onto $(N + P)/P$.

Proof. We have a natural action of R on $N + P$ by $r(n + p) = rn + rp$. This gives the subgroup $N + P$ an R -module structure by the fact that each of N and P are R -modules. The natural R -module isomorphism sends a coset $n + N \cap P \mapsto n + P$. $\triangle$

2.11 Free Modules

No starred exercises.

§ 3 Local Properties of Plane Curves

3.1 Multiple Points and Tangent Lines