

## 2E2 Tutorial Sheet 10<sup>1</sup>

15 January 2006

### Useful facts:

- Solving a linear differential equation: for  $A$  a  $2 \times 2$  matrix with eigenvalues  $\lambda_1$  and  $\lambda_2$  and corresponding eigenvectors  $\mathbf{x}_1$  and  $\mathbf{x}_2$  then if

$$\mathbf{y}' = A\mathbf{y} \tag{1}$$

the solution is

$$\mathbf{y} = C_1\mathbf{x}_1e^{\lambda_1 t} + C_2\mathbf{x}_2e^{\lambda_2 t} \tag{2}$$

where  $C_1$  and  $C_2$  are arbitrary constants.

- If initial condition are given, just set  $t = 0$  to find  $y_1(0)$  and  $y_2(0)$  and this should give simultaneous equations for  $C_1$  and  $C_2$ .

---

<sup>1</sup>Conor Houghton, [houghton@maths.tcd.ie](mailto:houghton@maths.tcd.ie), see also <http://www.maths.tcd.ie/~houghton/2E2.html>

## Questions

1. (2) Find the general solution for the system

$$\frac{dy_1}{dt} = 3y_1 + y_2 \quad (3)$$

$$\frac{dy_2}{dt} = y_1 + 3y_2 \quad (4)$$

2. (3) Find the solution of the system

$$\frac{dy_1}{dt} = 3y_1 + 4y_2 \quad (5)$$

$$\frac{dy_2}{dt} = 4y_1 - 3y_2 \quad (6)$$

with  $y_1(0) = 2$  and  $y_2(0) = -1$ .




Figure 1: Two containers with flow between them.

3. (3) As illustrated in Fig. 1, two large containers are connected and American style sandwich spread is pumped between them at a rate of  $1/2m^3s^{-1}$ . One container has volume  $5m^3$ , the other  $7m^3$ . Both are full of spread. Initially the smaller container contains pure jam, the second container has  $5m^3$  of jam and  $2m^3$  of peanut butter. Assume perfect mixing and so on.
- (i) Write down the differential equation for  $y_1(t)$  and  $y_2(t)$ , the amount of peanut butter in the first and second container.
  - (ii) Solve it to find  $y_1(t)$  and  $y_2(t)$  explicitly.
  - (iii) Use the initial data to find the values of the constants in the solution.