

## 2E2 Tutorial Sheet 9 Solutions<sup>1</sup>

8 January 2006

### Questions

1. (3) Find the eigenvectors and eigenvalues of the following matrices

$$(i) \begin{pmatrix} 10 & -4 \\ 18 & -12 \end{pmatrix} \quad (ii) \begin{pmatrix} 4 & 0 \\ 0 & -6 \end{pmatrix} \quad (iii) \begin{pmatrix} 0 & r \\ r & 0 \end{pmatrix} \quad (1)$$

*Solution:* In (i) the characteristic equation is

$$\begin{vmatrix} 4 - \lambda & 0 \\ 0 & -6 - \lambda \end{vmatrix} = 0 \quad (2)$$

so

$$(4 - \lambda)(-6 - \lambda) = 0 \quad (3)$$

So  $\lambda = 4$  or  $\lambda = -6$ . Taking the  $\lambda = 4$  first

$$\begin{pmatrix} 4 & 0 \\ 0 & -6 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = 4 \begin{pmatrix} a \\ b \end{pmatrix} \quad (4)$$

so  $4a = 4a$  and  $-6b = 4b$ , hence  $b = 0$  and  $a$  is arbitrary, taking  $a = 1$  an eigenvalue 4 eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (5)$$

Taking the  $\lambda = -6$

$$\begin{pmatrix} 4 & 0 \\ 0 & -6 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -6 \begin{pmatrix} a \\ b \end{pmatrix} \quad (6)$$

so  $4a = -6a$  and  $-6b = -6b$ , hence  $a = 0$  and  $b$  is arbitrary, taking  $b = 1$  a eigenvalue  $-6$  eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (7)$$

In (ii) the characteristic equation is

$$\begin{vmatrix} 10 - \lambda & -4 \\ 18 & -12 - \lambda \end{vmatrix} = 0 \quad (8)$$

so

$$(10 - \lambda)(-12 - \lambda) - (-4)18 = 0 \quad (9)$$

or

$$\lambda^2 + 2\lambda - 48 = 0 \quad (10)$$

Solve this gives us  $\lambda = 6$  or  $\lambda = -8$ . Taking the  $\lambda = 6$  first

$$\begin{pmatrix} 10 & -4 \\ 18 & -12 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = 6 \begin{pmatrix} a \\ b \end{pmatrix} \quad (11)$$

so the first equation is  $10a - 4b = 6a$  or  $a = b$ , the other equation is  $18a - 12b = 6b$  which is also  $a = b$ . Taking  $a = 1$  an eigenvalue 6 eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (12)$$

Taking the  $\lambda = -8$  next

$$\begin{pmatrix} 10 & -4 \\ 18 & -12 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -8 \begin{pmatrix} a \\ b \end{pmatrix} \quad (13)$$

so the first equation is  $10a - 4b = -8a$  or  $9a = 2b$ , the other equation is  $18a - 12b = -8b$  which is also  $9a = 2b$ . Taking  $a = 2$ ,  $b = 9$  an eigenvalue  $-8$  eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 2 \\ 9 \end{pmatrix} \quad (14)$$

In (iii) the characteristic equation is

$$\begin{vmatrix} -\lambda & r \\ r & -\lambda \end{vmatrix} = 0 \quad (15)$$

so

$$\lambda^2 - r^2 = 0 \quad (16)$$

or

$$\lambda = \pm r \quad (17)$$

Taking the  $\lambda = r$  first

$$\begin{pmatrix} 0 & r \\ r & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = r \begin{pmatrix} a \\ b \end{pmatrix} \quad (18)$$

so the equation is  $rb = ra$  or  $a = b$ . Taking  $a = 1$  an eigenvalue  $r$  eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (19)$$

<sup>1</sup>Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/2E2.html>

Taking the  $\lambda = -r$

$$\begin{pmatrix} 0 & r \\ r & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -r \begin{pmatrix} a \\ b \end{pmatrix} \quad (20)$$

so the equation is  $rb = -ra$  or  $a = -b$ . Taking  $a = 1$  an eigenvalue  $-r$  eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad (21)$$

2. (2) Find the eigenvectors and eigenvalues of the following matrices

$$(i) \begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix} \quad (ii) \begin{pmatrix} 0 & 3 \\ -3 & 0 \end{pmatrix} \quad (22)$$

*Solution:* In (i) the characteristic equation is

$$\begin{vmatrix} 3-\lambda & 4 \\ 4 & -3-\lambda \end{vmatrix} = 0 \quad (23)$$

so

$$(3-\lambda)(-3-\lambda) - 16 = 0 \quad (24)$$

or

$$\lambda^2 + 2\lambda - 48 - 25 = 0 \quad (25)$$

Solve this gives us  $\lambda = \pm 5$ . Taking the  $\lambda = 5$  first

$$\begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = 5 \begin{pmatrix} a \\ b \end{pmatrix} \quad (26)$$

so the first equation is  $3a + 4b = 5a$  or  $a = 2b$ , the other equation is  $4a - 3b = 5b$  which is also  $a = 2b$ . Taking  $a = 2$  an eigenvalue 5 eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad (27)$$

Taking  $\lambda = -5$  next

$$\begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -5 \begin{pmatrix} a \\ b \end{pmatrix} \quad (28)$$

so the first equation is  $3a + 4b = -5a$  or  $2a = -b$ , the other equation is  $4a - 3b = -5b$  which is also  $2a = -b$ . Taking  $a = 1$  an eigenvalue  $-5$  eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1 \\ -2 \end{pmatrix} \quad (29)$$

In (ii) the characteristic equation is

$$\begin{vmatrix} -\lambda & 3 \\ -3 & -\lambda \end{vmatrix} = 0 \quad (30)$$

so

$$\lambda^2 + 9 = 0 \quad (31)$$

or

$$\lambda = \pm 3i \quad (32)$$

Taking the  $\lambda = 3i$  first

$$\begin{pmatrix} 0 & 3 \\ -3 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = 3i \begin{pmatrix} a \\ b \end{pmatrix} \quad (33)$$

so the equation is  $3b = 3ia$  or  $a = -ib$ . Taking  $b = 1$  an eigenvalue  $3i$  eigenvector is,

$$\mathbf{x} = \begin{pmatrix} -i \\ 1 \end{pmatrix} \quad (34)$$

Taking the  $\lambda = -3i$

$$\begin{pmatrix} 0 & 3 \\ -3 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -3i \begin{pmatrix} a \\ b \end{pmatrix} \quad (35)$$

so the equation is  $3b = -3ia$  or  $a = ib$ . Taking  $a = 1$  an eigenvalue  $-3i$  eigenvector is,

$$\mathbf{x} = \begin{pmatrix} i \\ 1 \end{pmatrix} \quad (36)$$

3. (3) Find the solution for the system

$$\frac{dy_1}{dt} = -3y_1 + 2y_2 \quad (37)$$

$$\frac{dy_2}{dt} = -2y_1 + 2y_2 \quad (38)$$

*Solution:* This equation is  $\mathbf{y}' = \mathbf{A}\mathbf{y}$  with

$$\mathbf{A} = \begin{pmatrix} -3 & 2 \\ -2 & 2 \end{pmatrix}$$

We can find the eigenvalues, the characteristic equation is

$$\begin{vmatrix} -3-\lambda & 2 \\ -2 & 2-\lambda \end{vmatrix} = (\lambda+3)(\lambda-2) + 4 = \lambda^2 + \lambda - 2 = 0$$

so that  $\lambda_1 = 1$  and  $\lambda_2 = -2$ .

Next, we need the eigenvectors. First,  $\lambda_1$ :

$$\begin{pmatrix} -3 & 2 \\ -2 & 2 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \quad (39)$$

so  $-3a + 2b = a$  or  $b = 2a$ , hence, choosing  $a = 1$  we get

$$\mathbf{x}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}. \quad (40)$$

For  $\lambda_2$ :

$$\begin{pmatrix} -3 & 2 \\ -2 & 2 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -2 \begin{pmatrix} a \\ b \end{pmatrix} \quad (41)$$

so  $-3a + 2b = -2a$  giving  $a = 2b$ , choosing  $b = 1$  gives

$$\mathbf{x}_2 = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad (42)$$

Now, in general the solution is

$$\mathbf{y} = c_1 \mathbf{x}_1 e^{\lambda_1 t} + c_2 \mathbf{x}_2 e^{\lambda_2 t} \quad (43)$$

so, here,

$$\mathbf{y} = c_1 \begin{pmatrix} 1 \\ 2 \end{pmatrix} e^t + c_2 \begin{pmatrix} 2 \\ 1 \end{pmatrix} e^{-2t} \quad (44)$$